

Variation in risk factors of dementia among four elderly patient cohorts

Baqar Husaini, Van Cain, Meggan Novotny, Zahid Samad, Robert Levine, Majaz Moonis

Baqar Husaini, Van Cain, Meggan Novotny, Zahid Samad, Center For Prevention Research, Tennessee State University, Nashville, TN 37209, United States

Robert Levine, Department of Preventive and Family Medicine, Meharry Medical College, Nashville, TN 37208, United States

Majaz Moonis, Stroke Center, Department of Neurology, University of Massachusetts, Memorial Hospital, Worcester, MA 01655, United States

Author contributions: Husaini B, Levine R and Moonis M wrote the main manuscript text; Cain V and Novotny M prepared Tables and Figure 1; Samad Z was involved with discussion development; all authors reviewed the manuscript.

Supported by Tennessee State University: CDC, No. ED072081100; NCI, No. 5U54CA163066; by a NIH(National Center on Minority Health and Health Disparity to Meharry Medical College), No. P20MD000516

Correspondence to: Baqar Husaini, PhD, Professor Emeritus, Director Emeritus, Center For Preventive Research, Tennessee State University, 3500 John Merritt Blvd, Nashville, TN 37209, United States. bhusaini@tnstate.edu

Telephone: +1-615-2101132 Fax: +1-615-9635068

Received: April 8, 2014 Revised: May 7, 2014

Accepted: June 14, 2014

Published online: June 28, 2014

Abstract

AIM: To examine variation in risk factors that contributed to dementia among four elderly cohorts by race and gender.

METHODS: We examined 2008 Tennessee Hospital Discharged database for vascular factors that play a role in both stroke and dementia. Risk factors for dementia were examined for black and white patients aged 65+. Four race-gender groups of patients-white males (WM), black males (BM), white females (WF), and black females (BF) were compared for prevalence of dementia and stroke. A logistic model predicting dementia in each group separately used several vascular factors affecting

dementia directly or indirectly through stroke.

RESULTS: Three point six percent of patients hospitalized in 2008 had dementia and dementia was higher among females than males (3.9% *vs* 3.2%, $P < 0.001$), and higher among blacks than whites (4.2% *vs* 3.5%, $P < 0.000$). Further, BF had higher prevalence of dementia than WF (4.2% *vs* 3.8%, $P < 0.001$); similarly BM had more dementia than WM (4.1% *vs* 3.1%, $P < 0.001$). In logistic regression models, however, different patterns of risk factors were associated with dementia in four groups: among WF and WM, hypertension, diabetes, congestive heart failure, and stroke predicted dementia. Among BF and BM, only stroke and diabetes were related to dementia.

CONCLUSION: Aggressive management of risk factors (hypertension and diabetes) may subsequently reduce stroke and dementia hospitalization.

© 2014 Baishideng Publishing Group Inc. All rights reserved.

Key words: Dementia; Race; Gender; Stroke; Vascular factors

Core tip: Large medicaid in-hospital database that examines the differences in prevalence of dementia amongst blacks and white population and by gender. Clear differences emerge; blacks have greater burden of dementia including both genders. Risk factors leading to dementia differed between groups. White males and females had a higher association with stroke, hypertension, heart failure and diabetes while blacks had stroke and diabetes only as risk factors. This difference allows us to target these 2 groups with aggressive management early on to reduce the risk of dementia. The strength lies in analyzing a very large database to derive these conclusions.

Husaini B, Cain V, Novotny M, Samad Z, Levine R, Moonis M.

Variation in risk factors of dementia among four elderly patient cohorts. *World J Neurol* 2014; 4(2): 7-11 Available from: URL: <http://www.wjnet.com/2218-6212/full/v4/i2/7.htm> DOI: <http://dx.doi.org/10.5316/wjn.v4.i2.7>

INTRODUCTION

Dementia is among the most common neuropsychiatric disorder, which afflicts 5% to 10% of American elderly populations over the age of 65, and it accounts for a third of all psychiatric diagnoses among the elderly^[1-4].

Previous studies have reported on the role of cardiovascular (CVD) factors that contribute to dementia and dementia differences by race and gender. Dementia is higher among blacks and females^[5-17]. The role of these CVD factors among elderly cohorts by race-gender remains unknown. Since strokes occur more often among older people, older age possibly contributes to both stroke and dementia^[10-15]. Further, older age is also reported to be correlated with higher rates of hypertension (HTN), diabetes mellitus (DM), myocardial infarction (MI), stroke recurrence^[15], and congestive heart failure (CHF)^[16]. The risk of stroke increases with age in the presence of diabetes^[10-19].

While age is related to all cardiovascular risk factors, stroke, and dementia, it remains unclear whether these risk factors are related to dementia across four elderly race-sex cohorts either directly or indirectly through stroke. Thus, in this study, we explore two issues: (1) Do rates of dementia and cardiovascular risk factors vary by race-gender cohorts? (2) Do cardiovascular risk factors that may influence dementia (directly or indirectly) vary by race-gender cohorts?

MATERIALS AND METHODS

Data

We used Tennessee Hospital Discharge Data files on elderly patients (aged 65+; $n = 154945$) discharged in 2008. These files are administrative files submitted for reimbursement; they do not provide clinical data but only ICD-9 diagnoses for which a patient was treated. The attending physicians give diagnoses. Since the Tennessee population is largely composed of whites and blacks, we used four race-sex cohorts of dementia patients ($n = 5556$): White Males ($n = 1778$), White Females ($n = 3069$), Black Males ($n = 253$), and Black Females ($n = 456$). While the dementia patient's sample had only 11% black patients, this proportion is a good representation of black population in Tennessee, which is approximately 14%. The number of black males in the dementia group, though small with limitations on generalizability, is based on those who received the dementia diagnosis by the attending physicians. For the diagnosis of dementia, we used ICD-9 codes of 290.00, 290.20, 290.40-290.42, 291.2, 294.10, 294.11, and 294.20. Data on risk factors hypertension (HTN), diabetes mellitus (DM), hyperlip-

idemia (CHOL), cardiac arrhythmia (CA), stroke, congestive heart failure (CHF), and MI were also extracted for each patient. All discharge diagnoses in our analyses included a combination of both primary and secondary diagnoses. The dementia sample of patients ($n = 5556$) included whites (89%), blacks (11%), females (63%) and males (37%). The mean age of dementia patients was 82.5 years (SD = 9.1). Black males were younger in age (78.4 years, SD = 8.8) compared to other cohorts (Table 1). Finally, age-adjusted (per age 65+) dementia prevalence rate (of 690.6 per 100000 elderly) was developed per CDC methodology^[20].

Statistical analysis

Prevalence of hospitalized dementia patients (per 100000) was directly age-adjusted and indexed to the Year 2000 Census per methodology provided by CDC for the population at risk^[20]. Prevalence of dementia risk factors by race and gender were evaluated with Pearson χ^2 and the Fisher's Exact Tests. We also used multivariate logistic regression models to examine both the direct and indirect effects of all risk factors impacting dementia. We used logistic models (controlling age) for each race-gender cohort separately to examine the likelihood of dementia associated with each risk factor (Figure 1). Estimating separate equations for each race-gender cohort allowed for the effects of each risk factor to vary across four cohorts.

RESULTS

Prevalence of dementia and cardiovascular risk factors

Our analyses indicate that both dementia and stroke prevalence increased significantly ($P < 0.001$) with increasing age: dementia increased from 1.6% among 65-74 years old to 4.2% among 75-84 years old, to 6.6% among elderly aged 85+. Similarly, stroke prevalence increased from 13.1% among 65-74 years old to 17.3% among 75-84 years old, to 18.0% among elderly older than 85 years of age. Further, blacks have higher prevalence of dementia than whites (4.2% *vs* 3.5%, $P < 0.001$) and so do females than males (3.9% *vs* 3.2%, $P < 0.001$) (not shown). Among four cohorts, dementia prevalence was significantly higher ($P < 0.001$) in black females (4.2%), compared to black males (4.1%), white females (3.8%) and white males (3.1%). The overall diagnosis of dementia was 3.6% ($n = 5556$) of all elderly patients hospitalized in 2008 ($n = 154945$).

As a second trend, while one third of dementia patients had DM, CA, CHF, nearly 60% had stroke and 80% had hypertension (Table 1, col. 7). These risk factors varied across four dementia cohorts in that black patients had higher prevalence of HTN, DM, and Stroke (Table 1, col. 6), whereas white patients had higher prevalence of CA than blacks (Table 1, col. 3). Further, prevalence of MI was higher among males (combined cols. 1 + 4) compared to females (combined cols. 2 + 5). Some of these findings are consistent with those reported previously on risk factors.

Table 1 Clinical characteristics of dementia patients by race and gender cohorts

Variables	White males <i>n</i> = 1778	White females <i>n</i> = 3069	All white <i>n</i> = 9847	Black males <i>n</i> = 253	Black female <i>n</i> = 456	All black <i>n</i> = 709	All dementia age 65+ <i>n</i> = 5556
Column→	1	2	3	4	5	6	7
Mean age (SD)	80.6 (8.4)	83.9 (9.0)	82.7 (9)	78.4 (8.8)	82.3 (9.4)	80.9 (9.4)	82.5 (9.1)
HTN (%)	81.8	83.1	82.6	85.8	91.7 ¹	89.6 ²	83.5
DM (%)	38.1	30.6	33.4	47.2	52.6 ¹	50.6 ²	35.6
Chol (%)	11.3 ¹	8.5	9.6	8.7	9.4	9.2	9.5
CA (%)	42.4 ¹	36.3	38.6 ¹	35.6	32.5	33.6	37.9
MI (%)	7.6	5.3	6.2	7.9 ¹	5.7	6.5	6.2
CHF (%)	36.7 ¹	32.6	34.0	32.4	36.2	34.8	34.2
Stroke (%)	63.1	54.9	57.9	72.7 ¹	63.4	66.7 ¹	59.0
Dementia (%)	3.1	3.8	3.5	4.1	4.2 ¹	4.2 ¹	3.6

¹Differences across cohorts significant at $P < 0.001$; ²Differences between two racial groups (cols. 3 and 6) are significant at $P < 0.001$. HTN: Hypertension; DM: Diabetes mellitus; Chol: Hyperlipidemia; CA: Cardiac arrhythmia; MI: Myocardial infarction; CHF: Congestive heart failure; Stroke: Prior stroke.

Table 2 Odds ratios showing direct effect of risk factors on dementia by race-gender cohorts

Risk factors	White males	White females	All white	Black males	Black females	All black	All dementia patients
Column→	1	2	3	4	5	6	7
HTN	1.22 ²	1.10 ¹	1.15 ²	0.72	1.04 ns	0.88	1.13 ³
DM	1.16 ²	1.16 ²	1.16 ²	1.17 ns	1.28 ³	1.24 ³	1.17 ²
Chol	0.78	0.73	0.75	0.81	0.89	0.85	0.76
CA	0.91	0.89	0.89	1.05 ns	1.03 ns	1.04 ns	0.91
CHF	1.44 ³	1.16 ²	1.26 ²	1.14 ns	1.19 ns	1.17 ns	1.25 ³
MI	0.84	0.83	0.83	1.11 ns	0.8	0.92	0.84
Stroke	9.9 ³	7.8 ³	8.23 ³	13.3 ³	9.2 ³	10.39 ³	8.5 ³

¹Each logistic analysis included age in model, OR significant at $P < 0.05$; ²OR significant at $P < 0.01$, ³OR significant at $P < 0.000$; HTN: Hypertension; DM: Diabetes mellitus; Chol: Hyperlipidemia; CA: Cardiac arrhythmia; MI: Myocardial infarction; CHF: Congestive heart failure; Stroke: Prior stroke.

Cardiovascular predictors of dementia (direct and indirect effects)

Table 2 shows direct effects of CVD factors on dementia as odd ratios for each elderly race-sex cohort. Figure 1A shows both direct effects (in black lines) and indirect effects (in red lines) of risk factors through stroke on dementia.

The direct effect of risk factors on dementia show that among whites (Table 2, col. 3; Figure 1A and B black lines), four risk factors, namely HTN [odds ratio (OR), 1.15, 95%CI: 1.06-1.25], DM (OR, 1.16, 95%CI: 1.08-1.23), CHF (OR, 1.26, 95%CI: 1.18-1.35); and Stroke (OR, 8.23, 95%CI: 7.7-8.74) predicted onset of dementia. Among black males (Table 2, col. 4; Figure 1C), only stroke (OR, 13.3, 95%CI: 9.92-17.16) predicted dementia; and among black females (Table 2, col. 5; Figure 1D) dementia was predicted by two factors, namely DM (OR, 1.28, 95%CI: 1.04-1.57) and stroke (OR, 9.2, 95%CI: 7.4-11.22).

For indirect effect through stroke (red lines), Figure 1A and 1B show that among white patients (combined males and females), five risk factors (HTN, DM, Hyperlipidemia, CA, and CHF) were all related to stroke which in turn predicted dementia. Among blacks (both males and females in Figure 1C and D), only two factors (HTN,

DM) were related to stroke, which in turn predicted dementia. Overall, the indirect effects of HTN, DM, CA and HF on dementia through stroke remain intact for the total sample (Figure 1E).

DISCUSSION

Our findings indicate that the risk of both stroke and dementia increases with increasing age. For our study, prevalence of dementia among the elderly (aged 65+) was estimated at the rate of 690.6 per 100000 elderly population and this rate per 100000 elderly varied significantly among four cohorts: black males had the highest rate (902.2), followed by black females (811.4), white males (619.9), and white females (617.5). Further, since diabetes almost doubles the risk of dementia^[21], and stroke were more prevalent (with increasing age) among blacks, it appears that dementia among blacks, (who survived stroke and thus are in the sample) may largely result from a combination of both diabetes^[19,21] and recurring stroke^[15].

The influence of stroke on dementia appears to be consistent with previously reported findings of both increasing age, and stroke associated with HTN and DM^[15-23]. Since hypertension and diabetes are highly prevalent in all cohorts, additional investigation is needed




Figure 1 Odd ratios for risk factors predicting stroke and dementia. A: White males (aged 65+ yr); B: White female (aged 65+ yr); C: Black males (aged 65+ yr); D: Black females (aged 65+ yr); E: All dementia patients (aged 65+ yr). Each logistic analysis included age in model; ^aOR significant at $P < 0.05$; ^bOR significant at $P < 0.01$; ^cOR significant at $P < 0.000$. HTN: Hypertension; CA: Cardiac arrhythmia; MI: Myocardial infarction; CHF: Congestive heart failure.

to examine the role of small versus large vessel strokes in dementia. It is plausible that repeated small vessel infarction among the surviving stroke patients might contribute to the beginning of dementia with large vessel infarcts that may bring dementia to a recognizable level. Further, our data have not supported the previously reported role of hyperlipidemia in elevated levels of dementia. A plausible explanation for the lack of finding regarding hyperlipidemia in our study could be due to an effective treatment of this condition that may have neutralized the effect of hyperlipidemia on dementia.

We recognize that the effect of risk factors on dementia is not as robust as anticipated, particularly among black males. This is possibly due to a small number of black male patients in our sample. We recognize but would remiss, if we do not note that these black males are those who are stroke survivors and thus they show

a strong effect of stroke on dementia. It may also be noted that there is a higher mortality among older stroke male patients, and thus fewer number of surviving males, particularly black males in our sample, appears to be consistent with the longevity of black population in Tennessee (72.5 years) where black females live longer (longevity of 76 years) than black males (longevity of 68.7 years). Hence the number of black males in our dementia sample, though small, appears to be consistent with the longevity data for the African American population.

Finally, since DM, and stroke predict dementia across most race-gender cohorts, both primary and secondary preventive measures need to be aggressively pursued since most of these risk factors are amenable to effective management strategies aimed at reducing hospitalization for both stroke and dementia among the elderly.

ACKNOWLEDGEMENTS

The authors thank the World Congress of Neurology, Vienna, Austria, September 21-26, 2013

COMMENTS

Background

This paper addresses issues of worldwide importance to the field of neurology.

Research frontiers

The methods are sound as they utilize administrative files of discharged patients with dementia.

Innovations and breakthroughs

The paper presents extensive, populationbased information regarding patients with a discharge diagnosis of dementia. It presents evidence to show possible variations in risk factors according to gender and race. This information is highly useful for planning public health responses to this important problem.

Terminology

The paper represents a useful scientific contribution to the neurologic literature.

Peer review

In this paper, the authors report about the risk factors of dementia among four elderly groups. This is an interesting study. The paper is well written.

REFERENCES

- Gorelick PB, Scuteri A, Black SE, Decarli C, Greenberg SM, Iadecola C, Launer LJ, Laurent S, Lopez OL, Nyenhuis D, Petersen RC, Schneider JA, Tzourio C, Arnett DK, Bennett DA, Chui HC, Higashida RT, Lindquist R, Nilsson PM, Roman GC, Sellke FW, Seshadri S. Vascular contributions to cognitive impairment and dementia: a statement for healthcare professionals from the american heart association/american stroke association. *Stroke* 2011; **42**: 2672-2713 [PMID: 21778438 DOI: 10.1161/STR.0b013e3182299496]
- Folstein MF, Bassett SS, Anthony JC, Romanoski AJ, Nestadt GR. Dementia: case ascertainment in a community survey. *J Gerontol* 1991; **46**: M132-M138 [PMID: 2071834 DOI: 10.1093/geronj/46.4.M132]
- Husaini BA, Sherkat DE, Moonis M, Levine R, Holzer C, Cain VA. Racial differences in the diagnosis of dementia and in its effects on the use and costs of health care services. *Psychiatr Serv* 2003; **54**: 92-96 [PMID: 12509673 DOI: 10.1176/appi.ps.54.1.92]
- Kerola T, Kettunen R, Nieminen T. The complex interplay of cardiovascular system and cognition: how to predict dementia in the elderly? *Int J Cardiol* 2011; **150**: 123-129 [PMID: 21094551 DOI: 10.1016/j.ijcard.2010.10.018]
- Baker FM. Dementing illness in African American populations: Evaluation and management for the primary physician. *J Geriatr Psych* 1991; **24**: 73-91
- Etgen T, Sander D, Bickel H, Förstl H. Mild cognitive impairment and dementia: the importance of modifiable risk factors. *Dtsch Arztebl Int* 2011; **108**: 743-750 [PMID: 22163250 DOI: 10.3238/arztebl.2011.0743]
- Kume K, Hanyu H, Sato T, Hirao K, Shimizu S, Kanetaka H, Sakurai H, Iwamoto T. Vascular risk factors are associated with faster decline of Alzheimer disease: a longitudinal SPECT study. *J Neurol* 2011; **258**: 1295-1303 [PMID: 21327852 DOI: 10.1007/s00415-011-5927-y]
- Hall KS, Gao S, Baiyewu O, Lane KA, Gurejo O, Shen J, Ogunniyi A, Murrell JR, Unverzagt FW, Dickens J, Smith-Gamble V, Hendrie HC. Prevalence rates for dementia and Alzheimer's disease in African Americans: 1992 versus 2001. *Alzheimers Dement* 2009; **5**: 227-233
- Corrada MM, Brookmeyer R, Berlau D, Paganini-Hill A, Kawas CH. Prevalence of dementia after age 90: results from the 90+ study. *Neurology* 2008; **71**: 337-343 [PMID: 18596243 DOI: 10.1212/01.wnl.0000310773.65918.cd]
- Arntzen KA, Schirmer H, Wilsgaard T, Mathiesen EB. Impact of cardiovascular risk factors on cognitive function: the Tromsø study. *Eur J Neurol* 2011; **18**: 737-743 [PMID: 21143340 DOI: 10.1111/j.1468-1331.2010.03263.x]
- Rastas S, Pirttilä T, Mattila K, Verkkoniemi A, Juva K, Niinistö L, Lämsimies E, Sulkava R. Vascular risk factors and dementia in the general population aged > 85 years: prospective population-based study. *Neurobiol Aging* 2010; **31**: 1-7 [PMID: 18378359 DOI: 10.1016/j.neurobiolaging.2008.02.020]
- Ligthart SA, Moll van Charante EP, Van Gool WA, Richard E. Treatment of cardiovascular risk factors to prevent cognitive decline and dementia: a systematic review. *Vasc Health Risk Manag* 2010; **6**: 775-785 [PMID: 20859546 DOI: 10.2147/VHRM.S7343]
- Savva GM, Stephan BC. Epidemiological studies of the effect of stroke on incident dementia: a systematic review. *Stroke* 2010; **41**: e41-e46 [PMID: 19910553 DOI: 10.1161/STROKEAHA.109.559880]
- Whisnant JP. Modeling of risk factors for ischemic stroke. The Willis Lecture. *Stroke* 1997; **28**: 1840-1844 [PMID: 9303034 DOI: 10.1161/01.STR.28.9.1840]
- Sheinart KF, Tuhim S, Horowitz DR, Weinberger J, Goldman M, Godbold JH. Stroke recurrence is more frequent in Blacks and Hispanics. *Neuroepidemiology* 1998; **17**: 188-198 [PMID: 9701833 DOI: 10.1159/000026172]
- Husaini BA, Mensah GA, Sawyer D, Cain VA, Samad Z, Hull PC, Levine RS, Sampson UK. Race, sex, and age differences in heart failure-related hospitalizations in a southern state: implications for prevention. *Circ Heart Fail* 2011; **4**: 161-169 [PMID: 21178017 DOI: 10.1161/CIRCHEARTFAILURE.110.958306]
- Flicker L. Cardiovascular risk factors, cerebrovascular disease burden, and healthy brain aging. *Clin Geriatr Med* 2010; **26**: 17-27 [PMID: 20176290 DOI: 10.1016/j.cger.2009.12.005]
- Piguet O, Garner B. Vascular pharmacotherapy and dementia. *Curr Vasc Pharmacol* 2010; **8**: 44-50 [PMID: 19485934 DOI: 10.2174/157016110790226705]
- Khoury JC, Kleindorfer D, Alwell K, Moomaw CJ, Woo D, Adeoye O, Flaherty ML, Khatri P, Ferioli S, Broderick JP, Kissela BM. Diabetes mellitus: a risk factor for ischemic stroke in a large biracial population. *Stroke* 2013; **44**: 1500-1504 [PMID: 23619130 DOI: 10.1161/STROKEAHA.113.001318]
- Klein RJ, Schoenborn CA. Age adjustment using the 2000 projected U.S. population. *Healthy People 2010 Stat Notes* 2001; (20): 1-10 [PMID: 11676466]
- Ott A, Stolk RP, van Harskamp F, Pols HA, Hofman A, Breteler MM. Diabetes mellitus and the risk of dementia: The Rotterdam Study. *Neurology* 1999; **53**: 1937-1942
- Artero S, Ancelin ML, Portet F, Dupuy A, Berr C, Dartigues JF, Tzourio C, Rouaud O, Poncet M, Pasquier F, Auriacombe S, Touchon J, Ritchie K. Risk profiles for mild cognitive impairment and progression to dementia are gender specific. *J Neurol Neurosurg Psychiatry* 2008; **79**: 979-984 [PMID: 18450788 DOI: 10.1136/jnnp.2007.136903]
- Husaini B, Levine R, Sharp L, Cain V, Novotny M, Hull P, Orum G, Samad Z, Sampson U, Moonis M. Depression increases stroke hospitalization cost: an analysis of 17,010 stroke patients in 2008 by race and gender. *Stroke Res Treat* 2013; **2013**: 846732 [PMID: 23555070 DOI: 10.1155/2013/846732]

P- Reviewers: Louboutin JP, Trkulja V S- Editor: Ji FF

L- Editor: A E- Editor: Lu YJ





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

