

Reviewer 00050424

Thank you for your comment.

For the treatment of the bile duct stones, it is necessary to perform papillary treatment, and the treatment used at the time are broadly classified into two groups such as endoscopic papillary balloon dilatation and endoscopic sphincterotomy (EST). Golden standard is EST, however, there are patients with difficulty for EST, thus we must select the procedure based on understanding of the characteristics of the procedure, and patient backgrounds. For the treatment of the bile duct stones, it is necessary to perform papillary treatment, and the treatment used at the time are broadly classified into two groups such as EPBD and EST. Golden standard is EST, however, there are patients with difficulty for EST, thus we must select the procedure based on understanding of the characteristics of the procedure, and patient backgrounds. This report went review based on evidence accumulated. For example, a patient with the bleeding tendency and the patient after the billroth-2 rebuilding stomach list that you should choose one of EPBD as nipple measures within the article.

Reviewer 00053417

Thank you for your comment.

I report this report based on articles more than review reported than a conventional report in the past. The references cited seem to be a maximum in a past report. In addition, originality is thought to be high because the new knowledge such as the convalescence of EPBD of children is included.

Reviewer 025020

Thank you for your comment.

I kept it in the article that I limited to the measures with the endoscope to the last this time if I listed the following sentences in introduction because I did it.

Percutaneous transhepatic approach exists, too, but I have long time for treatment and am not performed very much because a maneuver is complicated.