

## Diagnosis of colonic amebiasis and coexisting signet-ring cell carcinoma in intestinal biopsy

Alexandra Grosse

Alexandra Grosse, Institute of Clinical Pathology, University Hospital Zurich, 8091 Zurich, Switzerland

**Author contributions:** Grosse A designed the report, collected the patient's clinical and histopathological data, analyzed and interpreted the data and wrote the paper.

**Institutional review board statement:** The study was reviewed and approved by the Institutional Review Board (Kantonale Ethikkommission Zürich).

**Informed consent statement:** Written informed consent was obtained from the patient for publication of this Case Report and any accompanying images.

**Conflict-of-interest statement:** The author has no conflicts of interests to declare.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Correspondence to:** Alexandra Grosse, MD, Institute of Clinical Pathology, University Hospital Zurich, Schmelzbergstrasse 12, 8091 Zurich, Switzerland. [alexandra.grosse@usz.ch](mailto:alexandra.grosse@usz.ch)  
Telephone: +41-44-2552637  
Fax: +41-44-2554440

Received: May 2, 2016  
Peer-review started: May 2, 2016  
First decision: June 20, 2016  
Revised: July 7, 2016  
Accepted: July 31, 2016  
Article in press: August 1, 2016  
Published online: September 28, 2016

### Abstract

Amebiasis is uncommon in developed countries. Several case reports in the literature emphasize that both the presenting symptoms and the radiological findings of colonic amebiasis closely resemble more common conditions, such as idiopathic inflammatory bowel disease and gastro-intestinal malignancy. We describe a unique case of colonic amebiasis (amebomas) coexisting with signet-ring cell carcinoma of the ileocecal valve, the cecum and the appendix. Endoscopically, the ulcerated tumor was indistinguishable from the ulcerations and pseudotumors (amebomas) detected in the ascending colon. Histological examination of biopsy specimens revealed the pathognomonic features of protozoa with ingested erythrocytes in combination with signet-ring cell infiltration. The author concludes that amebiasis may not only mimic carcinoma but, rarely, may coexist with carcinoma in the same patient. Clinicians and pathologists should be aware of this possibility in order not to delay diagnosis and treatment of malignant disease.

**Key words:** Colonic amebiasis; Histopathology; Parasitic disease; Colorectal carcinoma; Colonoscopy

© The Author(s) 2016. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** This case report presents the unique case of a patient in whom colonic amebiasis (amebomas) and signet-ring cell carcinoma of the ileocecal valve, the cecum and the appendix were diagnosed in intestinal biopsy specimens. In the setting of amebiasis, malignancy may easily be overlooked. Therefore, clinicians and pathologists should be aware that colonic amebiasis and colorectal carcinoma may coexist in the same patient.

Grosse A. Diagnosis of colonic amebiasis and coexisting signet-ring cell carcinoma in intestinal biopsy. *World J Gastroenterol* 2016; 22(36): 8234-8241 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v22/i36/8234.htm> DOI: <http://dx.doi.org/10.3748/wjg.v22.i36.8234>

## INTRODUCTION

Amebiasis caused by *Entamoeba histolytica* (*E. histolytica*) is one of the most fatal parasitic diseases worldwide. The organism is commonly acquired by ingestion of contaminated food and water and is endemic in developing countries with poor sanitation. Presentation of intestinal disease ranges from asymptomatic infection or acute proctocolitis to acute fulminant colitis with high mortality rates. Both the clinical presentation and the endoscopic appearance of colonic amebiasis can mimic colon carcinoma<sup>[1-8]</sup>. We present an exceedingly rare case of colonic amebiasis (amebomas) with coexisting signet-ring cell carcinoma of the ileocecal valve, the cecum and the appendix diagnosed upon histological and immunohistochemical examination of intestinal biopsy specimens.

## CASE REPORT

A 35-year-old woman presented to our hospital with diarrhea, right-lower-quadrant abdominal pain and cramps of two months' duration. The patient gave a travel history to Nepal where she had studied for one year prior to presentation. Her medical history was otherwise unremarkable; in particular, she had no family history of chronic idiopathic inflammatory bowel disease.

Laboratory tests revealed normocytic anemia (hemoglobin 105 g/L, hematocrit 0.34 L/L, and erythrocytes 3.70 T/L), with low vitamin B-12 (130 ng/L), high serum iron (35.3 µmol/L), significantly increased serum ferritin (1042 µg/L) and normal transferrin, as well as neutrophilic leukocytosis (leukocytes 11.52 G/L and neutrophils 9.30 G/L). Magnetic resonance imaging (MRI) showed mural thickening of the ileocecum and the appendix, which was interpreted as being inflammatory in nature (Figure 1); there was no evidence of an abscess. MRI also disclosed a nonhomogeneous, cystic mass of the left ovary that measured 7.9 cm in diameter. Gastroscopy and proctoscopy including anorectal examination were inconspicuous. Colonoscopy revealed multiple ulcerations and up to 10 well-circumscribed, ulcerated, tumor-like masses, separated by normal mucosa, in the ascending colon (Figure 2A and B). The masses were livid-colored at the periphery and necrotic at the center, and these masses ranged in size from 8 mm to 33 mm in diameter. A similar, but larger (4.5


cm in diameter) and more solid lesion was found at the ileocecal valve involving the adjacent cecum and the appendix (Figure 2C and D). These endoscopic manifestations were considered to be inconclusive. While the ulcerated lesions were found to be atypical of Crohn's disease, differential diagnoses included tuberculosis, lymphoma and carcinoma.

Microscopically, the masses in the ascending colon consisted of necrosis with large amounts of cell debris, desquamated epithelial cells and neutrophils, and extensive inflammatory tissue reaction in the absence of malignancy. Both in the debris and in the submucosal tissues there was evidence of periodic acid-Schiff (PAS)-positive amebas that displayed characteristic features of the invasive pathogen with ingested erythrocytes (Figure 3A-D). Because of the absence of malignancy, the masses in the ascending colon were considered to be amebomas, a complication of chronic amebic disease. Microscopic examination (hematoxylin-eosin and immunohistochemical stains) of biopsy specimens from the ileocecal valve showed amebas intermixed with signet-ring cells infiltrating the submucosa (Figure 3E-J). The CK 7, CK 20 and CDX-2 positive tumor cells, which stained negative with neuroendocrine markers, demonstrated preserved expression of the mismatch-repair proteins MLH1, PMS2, MSH2 and MSH6, indicating a microsatellite stable phenotype of carcinoma.

The pre-operative patient work-up included computed tomographic imaging of the chest and upper abdominal region, which revealed no extraintestinal manifestations of amebiasis, and explorative laparoscopy (Figure 4A and B) along with cytological examination of and biopsies from the peritoneum, which showed peritoneal carcinomatosis. After treatment with metronidazole, the patient underwent chemotherapy. Four months after diagnosis, right-sided hemicolectomy, omentectomy and Douglas-resection with hysterectomy and bilateral salpingo-oophorectomy were performed. Gross pathology of the surgical specimens showed tumor involvement of the cecum, the appendix and the ileocecal valve (Figure 5A). Metastases were found in the regional lymph nodes, the ovaries (Figure 5B), the fallopian tubes, the peritoneum, the omentum and the diaphragm. The biopsy-based diagnosis of signet-ring cell adenocarcinoma was confirmed by histology of the resected specimens (Figure 6A-F).

## DISCUSSION

*E. histolytica* is a major cause of diarrhea in developing countries, primarily in tropical and subtropical regions with poor sanitation and inadequate barriers between food/water and human feces. In developed countries, most infections arise in immigrants and travelers




**Figure 1** Magnetic resonance imaging showed mural thickening of the ileocecum and the appendix, which was interpreted as being inflammatory in nature. A-E: Magnetic resonance images (A: T2-weighted magnetization transfer contrast transversal; B: T2-weighted fat saturated transversal; C: T2-weighted half-Fourier acquired single-shot turbo spin-echo transversal; D: T1-weighted VIBE-Dixon postcontrast transversal, E: T1-weighted VIBE-Dixon postcontrast coronal) showing mural thickening (arrows in A-C) of the ileocecum and the appendix with contrast enhancement (arrow in D, asterisk in E).

from endemic areas. Young infants, pregnant women, malnourished individuals, and immunocompromised individuals are at highest risk for severe disease, which is characterized by complications such as perforation, toxic megacolon, fistulae, liver abscess and hemorrhage. Each year *E. histolytica* causes 40-50 million symptomatic infections worldwide, and approximately 40000 to 100000 people die each year from the disease, making this condition the second leading cause of death among parasitic diseases worldwide (after malaria)<sup>[9-11]</sup>. The majority of the patients (90%) remain asymptomatic<sup>[12,13]</sup>. Symptomatic patients can have simple acute proctocolitis or develop uncommon forms of GI involvement that include toxic megacolon, which is usually associated with corticosteroid treatment, abscess formation,

and acute fulminant colitis with a mortality rate over 40%<sup>[14]</sup>. Clinical symptoms range from mild, with diarrhea, abdominal cramps and right lower quadrant tenderness ("non-dysenteric" infection), to severe, with abdominal cramps, fever and mucoid or bloody diarrhea ("dysenteric or invasive" infection).

The infestation begins with ingestion of *E. histolytica* cysts from food or water contaminated with feces. After digestion of the cysts in the intestinal lumen trophozoites are released. The trophozoites reproduce by clonal expansion and finally form cysts in the colon that are excreted in the feces and into the environment completing the life cycle of the parasite. *E. histolytica* has phagocytic, proteolytic and cytolytic capabilities and is known to secrete metabolic enzymes that facilitate their invasion into the mucosa and submucosa




**Figure 2 Colonoscopy.** A, B: Colonoscopy showing multiple pseudotumors (amebomas) in the ascending colon, mimicking carcinoma; C, D: Endoscopic view of a large ulcerated tumor mass at the ileocecal valve.

causing characteristic flask-shaped intestinal ulcers. The parasites may penetrate through the mucosa and submucosa into the portal circulation and reach internal organs such as the liver, lungs, skin, etc. Amebiasis may involve any part of the bowel, but the cecum and the ascending colon are predilection sites<sup>[9]</sup>. The most common extraintestinal manifestation is liver abscess caused by hematogenous spread from the GI tract.


The pathophysiological mechanism includes adhesion of *E. histolytica* to the intestinal epithelial cells via the Gal/GalNAc lectin (galactose/N-acetylgalactosamine specific lectin) triggering host cell death through phagocytosis, cytotoxicity and caspase activation. Other molecules involved in the disease process are amebapores, a serine-rich *E. histolytica* protein, and cysteine proteases<sup>[15,16]</sup>. The ulcers observed on endoscopy vary in size from pin-head size to large (> 2.5 cm in diameter), coalescing, geographic or serpiginous lesions with intervening normal mucosa. Amebomas, a late complication of the disease, are pseudo-tumoral lesions formed by granulation tissue with peripheral fibrosis and a core of chronic inflammation usually found in the cecum or ascending colon. Amebomas may cause obstructive symptoms and can be confused with malignancy. In addition, both the endoscopic appearance and the histologic features (including crypt architectural distortion, mural fibrosis and scarring) of chronic amebic colitis may resemble

Crohn's disease. In contrast to amebic colitis, Crohn's disease does not spare the upper gastrointestinal tract and shows additional histological features typical of chronic idiopathic inflammatory bowel disease, such as granulomas, crypt abscesses and transmural lymphoid aggregates in the absence of protozoa.


Although a number of antigenic and molecular diagnostic tools have been developed over the years<sup>[17]</sup>, the most commonly used method for diagnosis of intestinal amebiasis remains the examination of stool or intestinal biopsy by microscopy. The reported sensitivity of microscopic stool examination in identifying amebic protozoa ranges from 25% to 60%<sup>[18]</sup>. Histological features of *E. histolytica* include distinct cell membranes, foamy cytoplasm, round and eccentrically located nuclei with margination of nuclear chromatin, and ingested red blood cells. The parasites may resemble macrophages but are CD-68 negative, PAS and trichrome positive and have smaller nuclei. Because of the conversion of glucose and pyruvate to ethanol, these organisms are highly motile, resulting in a pleomorphic shape (with diameters ranging from 10 to 50  $\mu$ m). In the current case, the diagnosis of colonic amebiasis and signet-ring cell carcinoma was established by histological examination of intestinal biopsy specimens showing the pathognomonic features of protozoa with ingested erythrocytes in combination with infiltrating tumor cells. While the findings were consistent with *E. histolytica* amebomas, antigenic or molecular tests for definitive



**Figure 3 Microscopic examination.** A, B: Microscopy (A: HE stain,  $\times 20$ , B: PAS,  $\times 20$ ) of biopsy specimens from the ascending colon showing aggregations of round PAS-positive *Entamoeba histolytica* trophozoites within cell debris; C, D: High power view (C: HE stain,  $\times 40$ , D: PAS,  $\times 20$ ) demonstrating PAS-positive trophozoites containing ingested erythrocytes (arrow in C); E, F: Microscopy (HE stain,  $\times 20$ ) of biopsy specimens from the ileocecal valve showing infiltrating signet-ring cells (arrows in E) in combination with amebic protozoa (arrow in F); G-J: The tumor cells stain positive with pancytokeratin (G:  $\times 20$ ), CK 7 (H:  $\times 20$ ), CK 20 (I:  $\times 20$ ) and CDX-2 (J:  $\times 20$ ).



**Figure 4 Explorative laparoscopy.** A, B: Laparoscopic view of a tumor mass (arrows) at the ileocecum and the appendix.




**Figure 5 Gross pathology of the surgical specimens.** A: Gross image showing a tumor mass (arrows) that involves the appendix, the ileocecal valve and the cecum. Surgical resection was performed after eradication of amebiasis and chemotherapy. B: Gross image of the Douglas-resection showing bilateral ovary metastases (arrows).

species identification were not performed, posing a limitation to the current study. Thus, (co-)infection with other morphologically indistinguishable *E. species* (*E. dispar* and *E. moshkovskii*) cannot definitively be excluded. Several investigations provide evidence that the prevalence of *E. histolytica*, derived from studies using conventional diagnostic tools (microscopy and/or culture), is often overestimated because of its epidemiological overlap with non-pathogenic *E. species*, and only a proportion of microscopically positive stool samples are true *E. histolytica* infections as confirmed by polymerase chain reaction assay<sup>[19-22]</sup>. However, given that *E. dispar* is generally considered non-invasive and thus related to asymptomatic infections (despite emerging evidence that *E. dispar* can acquire pathogenicity in the presence of bacteria and produce significant lesions in the human intestine and liver<sup>[23]</sup>), based on the clinical presentation, *E. histolytica* is likely the causative agent of dysenteric colitis and amebomas in the current case.

Colonic amebiasis can mimic colon carcinoma clinically, radiologically and endoscopically<sup>[1-8]</sup>. Conversely, amebiasis and coexisting carcinoma are exceedingly rare<sup>[24-29]</sup>. In two African studies, colorectal carcinoma was found to be associated with intestinal amebiasis in 6.1% and 6.5% of cases<sup>[24,25]</sup>. Furthermore, five cases of cervical, perineal, sigmoid and pulmonary carcinomas colonized by *E. histolytica* have been described in three case reports<sup>[26-28]</sup>. However, apart from a single case study dated from 1963<sup>[29]</sup>, to our knowledge the unique coincidence of colon carcinoma and amebomas has not been published previously in the literature.

The unusual co-occurrence of amebic infection and carcinoma poses particular challenges for both the pathologist and the clinician. Because coexisting malignancy in the setting of amebic disease can easily be overlooked, careful evaluation of individuals with amebiasis is necessary to confirm or rule out accompanying malignant disease.



**Figure 6** Biopsy-based diagnosis of signet-ring cell adenocarcinoma was confirmed by histology of the resected specimens. A, B: Microscopy (A: HE stain,  $\times 40$ , B: HE stain,  $\times 20$ ) of surgical specimens demonstrating infiltration of the colonic wall by signet-ring cells; C, D: The tumor involves the cecum (C: HE stain,  $\times 4$ ) and the appendix (D: HE stain,  $\times 4$ ); E, F: Microscopy showing metastases to the ovary (D: HE stain,  $\times 10$ ) and lymph nodes (E: HE stain,  $\times 10$ ).

## COMMENTS

### Case characteristics

A 35-year-old woman with no significant medical history, but a travel history to Nepal, presented with diarrhea and right-lower-quadrant abdominal pain and cramps of two months' duration.

### Clinical diagnosis

Right-lower-quadrant tenderness and mucoid diarrhea.

### Differential diagnosis

Carcinoma, lymphoma, tuberculosis or Crohn's disease.

### Laboratory diagnosis

Laboratory tests showed neutrophilic leukocytosis and normocytic anemia, thought to be related to vitamin B-12 deficiency and occult intestinal bleeding.

### Imaging diagnosis

Magnetic resonance imaging showed mural thickening of the cecum, appendix and adjacent terminal ileum. Colonoscopy disclosed multiple ulcerated, tumor-like masses (up to 33 mm in diameter) in the ascending colon and a 4.5 cm solid lesion at the ileocecal valve. Computed tomographic imaging of the chest

and upper abdominal region revealed no extraintestinal mass-like lesions.

### Pathological diagnosis

Amebomas and coexisting signet-ring cell carcinoma of the ileocecal valve, the cecum and the appendix with metastatic spread to the regional lymph nodes, ovaries, fallopian tubes, peritoneum, omentum and diaphragm.

### Treatment

Treatment with metronidazole followed by chemotherapy and right-sided hemicolectomy, omentectomy, hysterectomy and bilateral salpingo-oophorectomy.

### Related reports

Several case reports emphasize that colonic amebiasis can mimic colon carcinoma leading to misdiagnosis. Rarely, colorectal carcinoma may be colonized by *E. dispar* and *E. moshkovskii*. Apart from a single case report, the unique coincidence of colon carcinoma and amebomas has not been published previously in the literature.

### Term explanation

Amebomas, a late manifestation of amebiasis, are pseudo-tumoral lesions. They consist of granulation tissue with a core of chronic inflammation and

typically occur in the cecum or ascending colon.

### Experiences and lessons

Colonic amebiasis can be confused with a neoplastic process. However, rarely, colon carcinoma and amebic disease may coexist in the same patient. Gastroenterologists and pathologists should be aware of this possibility and carefully evaluate patients with amebiasis to rule out concurrent malignancy.

### Peer-review

The paper is well written. A limitation of the current study lies in the fact that definitive species identification was not performed using antigenic or molecular methods. However, the clinical presentation of the patient indicates that *Entamoeba histolytica* is likely the causative agent of dysenteric colitis and amebomas in the current case.

## REFERENCES

- 1 **Moorchung N**, Singh V, Srinivas V, Jaiswal SS, Singh G. Caecal amebic colitis mimicking obstructing right sided colonic carcinoma with liver metastases: a rare case. *J Cancer Res Ther* 2014; **10**: 440-442 [PMID: 25022419 DOI: 10.4103/0973-1482.136684]
- 2 **Lin CC**, Kao KY. Ameboma: a colon carcinoma-like lesion in a colonoscopy finding. *Case Rep Gastroenterol* 2013; **7**: 438-441 [PMID: 24403882 DOI: 10.1159/000355880]
- 3 **Ayari H**, Rebii S, Ghariani W, Daghfous A, Hasni R, Rehaïem R, Rezgui-Marhouf L, Zoghalmi A. [Colonic amoebiasis simulating a cecal tumor: case report]. *Med Sante Trop* 2013; **23**: 274-275 [PMID: 23797923 DOI: 10.1684/mst.2013.0186]
- 4 **Fernandes H**, D'Souza CR, Swethadri GK, Naik CN. Ameboma of the colon with amebic liver abscess mimicking metastatic colon cancer. *Indian J Pathol Microbiol* 2009; **52**: 228-230 [PMID: 19332922 DOI: 10.4103/0377-4929.48927]
- 5 **Hardin RE**, Ferzli GS, Zenilman ME, Gadangi PK, Bowne WB. Invasive amebiasis and ameboma formation presenting as a rectal mass: An uncommon case of malignant masquerade at a western medical center. *World J Gastroenterol* 2007; **13**: 5659-5661 [PMID: 17948943 DOI: 10.3748/wjg.v13.i42.5659]
- 6 **Ng DC**, Kwok SY, Cheng Y, Chung CC, Li MK. Colonic amoebic abscess mimicking carcinoma of the colon. *Hong Kong Med J* 2006; **12**: 71-73 [PMID: 16495594]
- 7 **Simşek H**, Elşürer R, Sökmensür C, Balaban HY, Tatar G. Ameboma mimicking carcinoma of the cecum: case report. *Gastrointest Endosc* 2004; **59**: 453-454 [PMID: 14997156 DOI: 10.1016/S0016-5107(03)02722-6]
- 8 **Ooi BS**, Seow-Choen F. Endoscopic view of rectal amebiasis mimicking a carcinoma. *Tech Coloproctol* 2003; **7**: 51-53 [PMID: 12750955 DOI: 10.1007/s101510300008]
- 9 **Stanley SL**. Amoebiasis. *Lancet* 2003; **361**: 1025-1034 [PMID: 12660071 DOI: 10.1016/S0140-6736(03)12830-9]
- 10 **Walsh JA**. Problems in recognition and diagnosis of amebiasis: estimation of the global magnitude of morbidity and mortality. *Rev Infect Dis* 1986; **8**: 228-238 [PMID: 2871619 DOI: 10.1093/clinids/8.2.228]
- 11 **Walsh J**. Prevalence of *Entamoeba histolytica* infection. In: Ravdin JJ. Amoebiasis: human infection by *Entamoeba histolytica*. New York: John Wiley and Sons, 1988: 93-105
- 12 **Gathiram V**, Jackson TF. A longitudinal study of asymptomatic carriers of pathogenic zymodemes of *Entamoeba histolytica*. *S Afr Med J* 1987; **72**: 669-672 [PMID: 2891197]
- 13 **Jackson TF**, Gathiram V, Simjee AE. Seroepidemiological study of antibody responses to the zymodemes of *Entamoeba histolytica*. *Lancet* 1985; **1**: 716-719 [PMID: 2857995 DOI: 10.1016/S0140-6736(85)91262-0]
- 14 Necrotizing amebic colitis: a frequently fatal complication. *Am J Surg* 1987; **154**: 451, 458, 462 [PMID: 3661852 DOI: 10.1016/0002-9610(86)90131-5]
- 15 **Mortimer L**, Chadee K. The immunopathogenesis of *Entamoeba histolytica*. *Exp Parasitol* 2010; **126**: 366-380 [PMID: 20303955 DOI: 10.1016/j.exppara.2010.03.005]
- 16 **Boettner DR**, Hutson C, Petri Jr WA. Galactose/N-acetylglactosamine lectin: the coordinator of host cell killing. *J Biosci* 2002; **27**: 553-557 [DOI: 10.1007/BF02704847]
- 17 **Flores MS**, Carrillo P, Tamez E, Rangel R, Rodríguez EG, Maldonado MG, Isibasi A, Galán L. Diagnostic parameters of serological ELISA for invasive amoebiasis, using antigens preserved without enzymatic inhibitors. *Exp Parasitol* 2016; **161**: 48-53 [PMID: 26684288 DOI: 10.1016/j.exppara.2015.12.006]
- 18 **Haque R**, Huston CD, Hughes M, Houpt E, Petri WA. Amebiasis. *N Engl J Med* 2003; **348**: 1565-1573 [PMID: 12700377 DOI: 10.1056/NEJMra022710]
- 19 **Yimer M**, Zenebe Y, Mulu W, Abera B, Saugar JM. Molecular prevalence of *Entamoeba histolytica*/dispar infection among patients attending four health centres in north-west Ethiopia. *Trop Doct* 2016; Epub ahead of print [PMID: 26834117 DOI: 10.1177/004975515627236]
- 20 **Kebede A**, Verweij JJ, Endeshaw T, Messele T, Tasew G, Petros B, Polderman AM. The use of real-time PCR to identify *Entamoeba histolytica* and *E. dispar* infections in prisoners and primary-school children in Ethiopia. *Ann Trop Med Parasitol* 2004; **98**: 43-48 [PMID: 15000730 DOI: 10.1179/000349804225003082]
- 21 **Fotedar R**, Stark D, Marriott D, Ellis J, Harkness J. *Entamoeba moshkovskii* infections in Sydney, Australia. *Eur J Clin Microbiol Infect Dis* 2008; **27**: 133-137 [PMID: 17957394 DOI: 10.1007/s10096-007-0399-9]
- 22 **Nath J**, Ghosh SK, Singha B, Paul J. Molecular Epidemiology of Amoebiasis: A Cross-Sectional Study among North East Indian Population. *PLoS Negl Trop Dis* 2015; **9**: e0004225 [PMID: 26633890 DOI: 10.1371/journal.pntd.0004225]
- 23 **Oliveira FM**, Neumann E, Gomes MA, Caliri MV. *Entamoeba dispar*: Could it be pathogenic. *Trop Parasitol* 2015; **5**: 9-14 [PMID: 25709947 DOI: 10.4103/2229-5070.149887]
- 24 **Ojo OS**, Odesanmi WO, Akinola OO. The surgical pathology of colorectal carcinomas in Nigerians. *Trop Gastroenterol* 1992; **13**: 64-69 [PMID: 1413101]
- 25 **Kenda JF**. Cancer of the large bowel in the African: a 15-year survey at Kinshasa University Hospital, Zaire. *Br J Surg* 1976; **63**: 966-968 [PMID: 188510 DOI: 10.1002/bjs.1800631220]
- 26 **Mhlanga BR**, Lanoie LO, Norris HJ, Lack EE, Connor DH. Amebiasis complicating carcinomas: a diagnostic dilemma. *Am J Trop Med Hyg* 1992; **46**: 759-764 [PMID: 1621901]
- 27 **Arroyo G**, Elgueta R. Squamous cell carcinoma associated with amoebic cervicitis. Report of a case. *Acta Cytol* 1989; **33**: 301-304 [PMID: 2728784]
- 28 **Zhu H**, Min X, Li S, Feng M, Zhang G, Yi X. Amebic lung abscess with coexisting lung adenocarcinoma: a unusual case of amebiasis. *Int J Clin Exp Pathol* 2014; **7**: 8251-8254 [PMID: 25550881]
- 29 **Cannavo L**. Sigmoid carcinoma in a patient with chronic amoebiasis or pseudoneoplastic amoebic granuloma (ameboma). *Panminerva Med* 1963; **5**: 159-162 [PMID: 14018317]

**P- Reviewer:** Magana M, Osawa S, Williams BL **S- Editor:** Qi Y  
**L- Editor:** A **E- Editor:** Zhang FH





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327

