

## Current noninvasive tests for colorectal cancer screening: An overview of colorectal cancer screening tests

Le-Le Song, Yue-Min Li

Le-Le Song, Yue-Min Li, Department of Radiotherapy, the Chinese PLA 309 Hospital, Beijing 100091, China

Le-Le Song, BioChain (Beijing) Science and Technology, Inc., Beijing 100176, China

Author contributions: All authors contributed to this manuscript.

Supported by The Beijing Municipal Science and Technology Project (capital public health project) No. Z151100003915092 sponsored by the Beijing Municipal Science and Technology Commission.

Conflict-of-interest statement: There is no conflict of interest associated with any of the senior author or other coauthors contributed their efforts in this manuscript.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Le-Le Song, MD, PhD, Department of Radiotherapy, the Chinese PLA 309 Hospital, No. 17, Heishanhu Road, Haidian District, Beijing 100091, China. [songlele@sina.com](mailto:songlele@sina.com)  
Telephone: +86-10-66775222

Received: June 12, 2016

Peer-review started: June 17, 2016

First decision: July 30, 2016

Revised: August 1, 2016

Accepted: August 30, 2016

Article in press: August 31, 2016

Published online: November 15, 2016

### Abstract

Colorectal cancer (CRC) has become the third most common cancer in the world. Screening has been shown to be an effective way to identify early CRC and precancerous lesions, and to reduce its morbidity and mortality. Several types of noninvasive tests have been developed for CRC screening, including the fecal occult blood test (FOBT), the fecal immunochemical test (FIT), the fecal-based DNA test and the blood-based DNA test (the SEPT9 assay). FIT has replaced FOBT and become the major screening test due to high sensitivity, specificity and low costs. The fecal DNA test exhibited higher sensitivity than FIT but its current cost is high for a screening assay. The SEPT9 assay showed good compliance while its performance in screening needs further improvements. These tests exhibited distinct sensitivity and specificity in screening for CRC and adenoma. This article will focus on the performance of the current noninvasive *in vitro* diagnostic tests that have been used for CRC screening. The merits and drawbacks for these screening methods will also be compared regarding the techniques, usage and costs. We hope this review can provide suggestions for both the public and clinicians in choosing the appropriate method for CRC screening.

**Key words:** Colorectal cancer; Adenoma; Fecal immunochemical test; Fecal DNA; SEPT9; Septin 9

© **The Author(s) 2016.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** The choice of colorectal cancer (CRC) screening methods is crucial for screening validity and compliance. Currently, the fecal immunochemical test (FIT), fecal DNA and the blood-based SEPT9 assays are the three *in vitro* diagnostic tests for CRC screening. In this article, we reviewed the current application of the three types of assays and compared their performance

in CRC screening. FIT is still the cheapest method with high screening validity, and fecal DNA tests also exhibit high validity but its price is high. In contrast, the SEPT9 assay showed high compliance with acceptable performance. The choice of screening test may depend on the balance of performance, compliance and costs.

---

Song LL, Li YM. Current noninvasive tests for colorectal cancer screening: An overview of colorectal cancer screening tests. *World J Gastrointest Oncol* 2016; 8(11): 793-800 Available from: URL: <http://www.wjgnet.com/1948-5204/full/v8/i11/793.htm> DOI: <http://dx.doi.org/10.4251/wjgo.v8.i11.793>

---

## INTRODUCTION

Colorectal cancer (CRC) has become the second and the third leading cause of new cancer cases in Europe and in the United States, respectively<sup>[1]</sup>. There were approximately 142820 new cases with 50830 deaths in the United States in 2013, and approximately 447000 new cases of CRC and 215000 deaths in European countries in 2012<sup>[1,2]</sup>. The new cases for CRC are approximately 400000 in China in 2012, and it has become the third leading cause of death in the country<sup>[3]</sup>.

Regular screening can achieve early CRC detection and early treatment. However, 60%-70% of patients are found at middle- or late-stage CRC when they are diagnosed<sup>[4]</sup>. Approximately 60% CRC deaths could be avoided and the average 5-year survival rate could be increased from 46% to 73% if healthy people carry out a regular periodic screening each year<sup>[5]</sup>. Therefore, an effective early screening method for CRC can reduce CRC morbidity and mortality.

There are four *in vitro* diagnostic (IVD) screening method currently available for CRC screening, including the fecal occult blood test (FOBT), the fecal immunochemical test (FIT), the fecal DNA test and the plasma SEPT9 gene methylation test. This review will provide a detailed analysis on the performance of these tests, and compare their merits and drawbacks in CRC screening. It is our aim for this review that the public and the professionals can choose the appropriate methods for CRC screening.

## STOOL-BASED TESTS FOR CRC SCREENING

### The FIT test

The guaiac FOBT test (gFOBT) has been used for a long time as a screening test for CRC. It exhibited a sensitivity of 12.9%-79.4% with a specificity of 86.7%-97.7% for CRC screening in many studies<sup>[6-13]</sup>. However, its sensitivity and specificity for CRC detection is lower than the more specific FIT (previous called iFOBT) test. This is because the gFOBT relies on peroxidase-like activity

between heme and guaiac, which can be affected by many factors in daily diet without distinguishment between upper and lower gastrointestinal (GI) tract bleeding, while the FIT test targets the hemoglobin in the lower GI tract, as hemoglobin from upper GI tract will be degraded when it arrives at lower GI tract. This characteristic allows FIT test to specifically detect the bleeding from lower GI tract, and therefore detect the diseases with bleeding, such as adenoma, polyps, inflammatory diseases and CRC, etc. As the gFOBT test has many drawbacks in CRC screening, FIT is used more commonly in current CRC screening. We therefore focus on the performance of FIT test in this review.

The performance of FIT test in CRC screening in asymptomatic, average-risk adults has been listed in Table 1. Data from 19 studies showed that the overall sensitivity for CRC was 0.79 (95%CI: 0.69-0.86) and the overall specificity was 0.94 (95%CI: 0.92-0.95)<sup>[12,14-31]</sup>. This includes a total of 113360 subjects with 437 CRC cases confirmed by colonoscopy or 2-year follow-up. As the overall sensitivity and specificity are satisfactory for a cancer screening test with low costs, FIT is currently the most commonly-used method for CRC screening. The overall CRC positivity rate of 0.39% (437/113360) appeared to be significantly lower than the other two screening reports with asymptomatic, average-risk adults using fecal DNA (0.65%, 65/9989;  $\chi^2 = 15.93$ ,  $P < 0.001$ )<sup>[32]</sup> and SEPT9 gene methylation assay (0.67%, 53/7941;  $\chi^2 = 14.66$ ,  $P < 0.001$ )<sup>[33]</sup>, respectively, indicating that the use of 2-year follow-up as a confirmatory methods may result in underestimation of CRC cases.

The use of qualitative FIT or quantitative FIT has always been an issue in choosing the FIT test for screening. A strip test (colloidal gold immunochromatographic method) is currently the main technique for qualitative FIT. It does not need specific instruments and the interpretation of test results relies on human recognition of test bands, although instruments are available to digitize the chrominance of the bands. In contrast, immunoturbidimetry is the main method for quantitative FIT, and the current devices include automated instrument for samples processing and colorimetry. Therefore, the current qualitative FIT appears to be faster, more convenient, less costly while more subjective than the quantitative FIT.

The performance between the qualitative and quantitative FIT showed significant differences. As shown in Table 1, the overall sensitivity of the qualitative FIT was 0.82<sup>[12,15-21]</sup>, which was significantly higher than that of the quantitative FIT (0.73) ( $\chi^2 = 3.933$ ,  $P = 0.047$ )<sup>[22-31]</sup>, while the qualitative FIT exhibited significantly lower specificity than the quantitative FIT (0.93 vs 0.95) ( $\chi^2 = 81.64$ ,  $P < 0.001$ ), although the difference was small. This comparison needs to be interpreted with caution, as different studies used different cutoff values and resulted in distinct sensitivity and specificity. Ideally, they should be compared under the same cutoff value so that the sensitivity and specificity can be directly

**Table 1** The sensitivity and specificity of qualitative and quantitative fecal immunochemical test

Ref.	Qualitative FIT				Ref.	Quantitative FIT			
	Total cases	CRC cases	Sensitivity	Specificity		Total cases	CRC cases	Sensitivity	Specificity
Allison <i>et al</i> <sup>[12]</sup> , 1996	7493	35	0.69	0.94	Sohn <i>et al</i> <sup>[22]</sup> , 2005	3794	12	0.25	0.99
Allison <i>et al</i> <sup>[15]</sup> , 2007	5356	14	0.86	0.97	Levi <i>et al</i> <sup>[23]</sup> , 2011	1204	6	1.00	0.88
Cheng <i>et al</i> <sup>[16]</sup> , 2002	7411	16	0.88	0.91	Levi <i>et al</i> <sup>[24]</sup> , 2007	80	3	0.67	0.83
Nakama <i>et al</i> <sup>[17]</sup> , 1999	4611	18	0.56	0.97	Morikawa <i>et al</i> <sup>[25]</sup> , 2005	21805	79	0.66	0.95
Nakama <i>et al</i> <sup>[18]</sup> , 1996	3365	12	0.83	0.96	Launoy <i>et al</i> <sup>[26]</sup> , 2005	7421	28	0.86	0.94
Parra-Blanco <i>et al</i> <sup>[19]</sup> , 2010	1756	14	1.00	0.93	Itoh <i>et al</i> <sup>[27]</sup> , 1996	27860	89	0.87	0.95
Chiu <i>et al</i> <sup>[20]</sup> , 2013	8822	13	0.85	0.92	Nakazato <i>et al</i> <sup>[28]</sup> , 2006	3090	19	0.53	0.87
Chiang <i>et al</i> <sup>[21]</sup> , 2011	2796	28	0.96	0.87	Park <i>et al</i> <sup>[29]</sup> , 2010	770	13	0.77	0.94
					de Wijkerslooth <i>et al</i> <sup>[30]</sup> , 2012	1256	8	0.75	0.95
					Brenner <i>et al</i> <sup>[31]</sup> , 2013	2235	15	0.73	0.96
					Brenner <i>et al</i> <sup>[31]</sup> , 2013	2235	15	0.60	0.95
Overall (pooled data)	41610	150	0.82	0.93		71750	287	0.73	0.95

CRC: Colorectal cancer; FIT: Fecal immunochemical test.

**Table 2** Comparison of sensitivity and specificity between Cologuard and fecal immunochemical test

Pathological categories	Cologuard	FIT
Sensitivity <sup>[32]</sup>		
CRC	92.3%	73.8%
Advanced precancerous lesions	42.4%	23.8%
Polyps with high-grade dysplasia	69.2%	46.2%
Serrated sessile polyps	42.4%	5.1%
Specificity <sup>[32]</sup>		
Nonadvanced or negative findings	86.6%	94.9%
Negative results on colonoscopy	89.8%	96.4%

CRC: Colorectal cancer; FIT: Fecal immunochemical test.

compared. The pooled data analyzed here provides a reference for comparing the two types of FIT tests. It can be suggested that the quantitative FIT may be a good choice for CRC screening tests that do not need high accuracy or are performed in hospitals where automated instruments are not available.

However, it should be mentioned that the cutoff value for qualitative FIT is preset, while the cutoff value for quantitative FIT can be adjusted to balance the sensitivity with specificity. Therefore, the data format for qualitative FIT is "positive" or "negative" without traceability, while the results from quantitative FIT are digitized with traceability. This is extremely useful when the relationship between the amount of bleeding in a certain disease and the population/personal information (such as diet, age, habit, sex, *etc.*) is investigated. Future model for predicting CRC incidence might partially relies on the data from quantitative FIT.

### The fecal DNA test

The detection of abnormal DNA or epigenetic markers from colorectal lesions is based on natural exfoliation of cancerous or precancerous cells into the colorectal tract. The fecal DNA test aims at detecting the DNA mutations, microsatellite instability, impaired DNA mismatch repair and abnormal methylation. There are many studies focusing on the detection of CRC by fecal DNA markers<sup>[34,35]</sup>, and the overall sensitivity for CRC

detection by various fecal DNA marker combinations ranged from 53% to 87%, with specificities beyond 76%<sup>[34,35]</sup>. Although there are a large number of fecal DNA markers available in these studies, the first commercial fecal DNA test was not available until the approval of Cologuard (Exact Sciences, Madison, WI, United States) by the United States FDA in 2014. Imperiale *et al*<sup>[32]</sup> published the leading study on Cologuard in 2014. By randomizing subjects to Cologuard or FIT screening, it showed that the sensitivity of Cologuard was superior to that of FIT in CRC, advanced precancerous lesions, polyps with high-grade dysplasia and serrated sessile polyps, while its specificity appeared to be lower than that of FIT (Table 2).

The Cologuard DNA test includes quantitative molecular assays for KRAS mutations, aberrant NDRG4 and BMP3 methylation, and  $\beta$ -actin, plus a hemoglobin immunoassay. As the hemoglobin immunoassay is essentially a FIT test, Cologuard is actually a combination of gene mutation, methylation and occult blood tests. The multitarget stool DNA test provides a new way that combines various detecting technology to detect CRC and early colorectal lesions with high sensitivity and specificity. The high detection of precancerous lesions, HGD and serrated sessile polyps is extremely useful for a screening test, as these lesions may develop into CRC if they are not resected. The only obstacle for broad application of Cologuard is the cost, as the detection of multitargets increased the cost of the test. Its current expense of \$599 per test is high for a routine screening assay.

## BLOOD-BASED TESTS FOR CRC SCREENING

### The plasma SEPT9 gene methylation assay

An ideal screening test for cancer could be a simple blood test in the foreseeable future. The plasma SEPT9 gene methylation test Epi proColon (Epigenomics AG, Berlin, Germany) is currently the only commercially

**Table 3 The reported positive detection rate for each colorectal cancer stage using 1/3 algorithm**

Ref.	Positive detection rate for each colorectal cancer stage			
	I	II	III	IV
deVos <i>et al</i> <sup>[38]</sup> , 2009	52.6% (10/19)	75.0% (30/40)	77.8% (21/27)	100.0% (4/4)
Warren <i>et al</i> <sup>[40]</sup> , 2011	71.4% (5/7)	90.3% (28/31)	100.0% (7/7)	100% (5/5)
Tóth <i>et al</i> <sup>[41]</sup> , 2012	84.0% (21/25)	100.0% (14/14)	100.0% (35/35)	100.0% (18/18)
Lee <i>et al</i> <sup>[47]</sup> , 2013	30.8% (8/26)	36.7% (11/30)	25.0% (7/28)	64.7% (11/17)
Johnson <i>et al</i> <sup>[44]</sup> , 2014	61.5% (16/26)	80.0% (16/20)	65.2% (15/23)	92.3% (12/13)
Pooled positive detection rate	58.3% (60/103)	73.3% (99/135)	70.8% (85/120)	87.7% (50/57)

**Table 4 The reported positive detection rate for each colorectal cancer stage using 2/3 algorithm**

Ref.	Positive detection rate for each colorectal cancer stage			
	I	II	III	IV
Grützmann <i>et al</i> <sup>[37]</sup> , 2008	50.0% (11/22)	69.4% (25/36)	79% (42/53)	91% (10/11)
deVos <i>et al</i> <sup>[38]</sup> , 2009	26.3% (5/19)	60.0% (24/40)	66.7% (18/27)	75.0% (3/4)
Tóth <i>et al</i> <sup>[41]</sup> , 2012	60.0% (15/25)	92.8% (13/14)	81.6% (31/35)	77.8% (14/18)
Jin <i>et al</i> <sup>[46]</sup> , 2015	66.7% (12/18)	82.6% (19/23)	84.1% (37/44)	100.0% (5/5)
Pooled positive detection rate	51.2% (43/84)	71.7% (81/113)	80.5% (128/159)	84.2% (32/38)

available blood-test for CRC early detection and screening, and was approved recently by the United States FDA as a CRC screening test for average-risk population over 50 years old. Many clinical studies have proved the test to be a method with acceptable sensitivity and specificity for CRC detection<sup>[33,36-49]</sup>. The test was firstly developed by Lofton-Day *et al*<sup>[36]</sup> in 2008 as a research kit, and was commercialized by Epigenomics AG as its first generation assay Epi proColon 1.0. At the same time, ARUP lab also developed its SEPT9 assay as a lab-developed test<sup>[40]</sup>. Abbott developed its real-time mS9 CRC assay, but there was only one report on its performance and the sensitivity of 36.3% was much lower than other SEPT9 tests<sup>[47]</sup>. The 2<sup>nd</sup> generation test (Epi proColon 2.0) was launched in 2011-2012 with better performance. Till today, most reports on the SEPT9 assay appeared to be case-control study or cohort study investigating the test performance in selected population, exhibiting a sensitivity of 36.6%-95.6% with a specificity of 81.5%-99.0% using 1/3, 2/3, 1/2 or 1/1 algorithm<sup>[33,36-49]</sup>. In contrast, there is only one study (the PRESEPT trial) investigating the application of the assay in CRC screening in average-risk population, exhibiting a sensitivity of 48.2% and 68.2% with a specificity of 91.5% and 80.0% using 1/2 or 1/3 algorithm, respectively<sup>[33,43]</sup>.

Detection of early stage CRC is crucial for early intervention and reduction of mortality. The positive detection rate (PDR) of the SEPT9 assay for stage I, II, III and IV was 26.3%-84.0%, 36.7%-100.0%, 25.0%-100.0% and 64.7%-100.0%, respectively, depending on different algorithm, exhibiting a huge variation for each stage. As 1/3 and 2/3 algorithm are the most commonly used methods for result interpretation, we calculated the PDR for each stage using the two algorithms. The pooled PDR for stage I, II, III and IV was 58.3%, 73.3%, 70.8% and 87.7%, respectively, using 1/3 algorithm (Table 3), and was

51.2%, 71.7%, 80.5% and 84.2%, respectively, using 2/3 algorithm (Table 4)<sup>[36-47]</sup>. No statistical difference in PDR in any stage between the two algorithms has been found. It can be clearly seen that the PDR for early stage CRC (stage I) was above 50% and fell into 70%-80% for stage II and III, which is acceptable for a blood-based CRC test. However, these PDRs were from case-control or cohort studies, and more studies should be performed at screening settings.

Although the SEPT9 assay was designed for CRC detection, researchers also studied its detection sensitivity for precancerous adenoma. The pooled PDR for non-advanced adenoma and advanced adenoma (AA) was 10.0% and 18.2%, respectively, from six studies, in which the PDR for AA was significantly higher than the PDR of normal control group (11.8%,  $\chi^2$  test,  $P < 0.001$ )<sup>[30,33,37,40,43,46]</sup>. However, as PDR of 18.2% was still too low for an effective test, the SEPT9 assay may not be applicable in adenoma detection.

The SEPT9 assay exhibited high compliance in screening. One recent report showed that 63% of subjects in a CRC screening study refused colonoscopy. 97% of subjects who refused colonoscopy accepted a noninvasive screening test, and 83% chose the SEPT9 test and 15% chose FIT test. The majority of patients who refused colonoscopy chose the SEPT9 assay due to its convenience and less time-consuming procedure<sup>[50]</sup>.

**CEA and other serum glycoprotein markers**

CEA and carbohydrate antigen 199 (CA199) are the two most common serum-based glycoprotein CRC markers, however, they are not appropriate for CRC screening due to their low sensitivity and the lack of CRC specificity, especially for early-stage CRC<sup>[41,51-53]</sup>. For example, CEA test exhibited a sensitivity of 40.9%-51.8% and a specificity of 85.2%-95% for CRC detection in three studies<sup>[41,51,52]</sup>. Therefore, it is more appropriate to be used in monitoring the CRC recurrence or response from

**Table 5** Sensitivity and specificity of fecal immunochemical test, fecal DNA and SEPT9 tests in colorectal cancer and advanced adenoma screening

	FIT <sup>[12,15-31]</sup>	Fecal DNA <sup>[32]</sup>	SEPT9 <sup>[43]</sup>
Sensitivity (CRC)	79%	92%	68%
Specificity	94%	87%	80%
Sensitivity (AA)	24%	42%	18%

FIT: Fecal immunochemical test; CRC: Colorectal cancer; AA: Advanced adenoma.

patients to surgical or systemic therapy, rather than screening<sup>[53]</sup>.

The main drawback of serum glycoprotein markers in CRC screening is that the sensitivity and specificity of any single marker is not high enough to make it a reliable indicator. These markers have been found in various cancers other than CRC with low sensitivity for early stage lesions. Combined use of multiple markers may be a way to achieve diagnostic significance in CRC detection. In one report, five glycoprotein markers, including CEA, CA199, CA242, CA72-4, and CA125, are used together as indicators for CRC. It showed that the sensitivity of any single marker was low (18.8%–52.2%) for detecting CRC in stages I and II, while the combination of the five exhibited a sensitivity of 85.3% at the specificity of 95%<sup>[54]</sup>.

## COMPARISON OF NONINVASIVE TESTS FOR CRC SCREENING

The sensitivity for CRC and AA, and the specificity in asymptomatic average-risk population for FIT, fecal DNA and SEPT9 tests are shown in Table 5. It can be seen that the fecal DNA test exhibited the best performance in terms of sensitivity for CRC and AA, while its specificity was slightly lower than that of the FIT. It is noteworthy that the fecal DNA test can detect 42% AA, which may reduce the number of subjects progressing to CRC, *i.e.*, reducing the CRC morbidity. The SEPT9 assay is the only blood-base CRC screening assay currently. Although its screening performance was not satisfactory at the moment, it showed very high compliance<sup>[50]</sup>. The blood-based CRC screening assay may be more popular if the sensitivity and specificity in screening setting could be improved to the level of those in case-control studies (ideally sensitivity > 70% and specificity > 90% for CRC screening).

The current costs for FIT, fecal DNA and the SEPT9 test are \$10-50, \$599 and approximately \$170, respectively. As the recommended screening frequency for FIT, fecal DNA and the SEPT9 is once per year, once per three years and once per year, respectively, FIT might be the cheapest test considering the balance between performance and costs. However, the quality adjusted life year of the three tests should be compared under the same setting to evaluate the cost-effectiveness of them, although some studies have been performed for each

**Table 6** Positive detection rate of SEPT9, fecal immunochemical test and carcino-embryonic antigen tests and various combined tests

SEPT9 alone	FIT alone	CEA alone	SEPT9 + FIT	SEPT9 + CEA	FIT + CEA	SEPT9 + FIT + CEA
77.00%	74.6%	41.3% <sup>e</sup>	94.4% <sup>c</sup>	86.4% <sup>c</sup>	84.5%	97.2% <sup>e</sup>
(181/235)	(53/71)	(97/235)	(67/71)	(203/235)	(60/71)	(69/71)

<sup>c</sup>*P* < 0.01; <sup>e</sup>*P* < 0.001 *vs* SEPT9 alone. FIT: Fecal immunochemical test; CEA: Carcino-embryonic antigen; NS: Not significant.

individual method in different settings, such as different health systems.

## CRC SCREENING WITH COMBINED TESTS

The combination of fecal DNA (mutation and methylation) with a hemoglobin immunoassay in Cologuard has provided a good example for CRC screening when multiple markers are analyzed together to enhance the detection sensitivity. There are merits and drawbacks for this strategy. First, combination of multiple markers enhances sensitivity at the price of reducing specificity. The number of false positive cases will increase with the increased number of markers. Therefore, to identify the markers with high sensitivity and specificity and to find the best combination of markers remain a challenge for combined screening test development. Ideally, the number of markers should be kept to minimum, while the sensitivity and specificity can be balanced to provide the best performance. Secondly, the detection of multiple markers with distinct methods increases the technical difficulties in an assay. For example, the detection of mutation in Cologuard may use sequencing or PCR method, while the detection of methylation needs to use the methylation specific PCR method containing bisulfite conversion. In contrast, immunoassay is used in the detection of hemoglobin. Furthermore, the sample preparation procedure may also be different for detecting different abnormalities. Therefore, a good combined test needs not only optimization of each individual test, but also an accurate algorithm to maximize the performance of each test. The optimization and interpretation of the combined test must come from clinical trials with large number of cases. Thirdly, a screening test should be accurate, fast, convenient, simple and cheap. These features allow large-scale screening in a certain period of time, and allow easy test in areas where test instruments are not available. In addition, low costs ensure screening tests for average-risk population, in which the CRC incidence could be lower than 1% in people over 50 years old. All the above considerations need to be addressed in future development of combined screening test.

As FIT, SEPT9 and CEA tests are all CRC detection tests with high specificity, the combination of them may provide higher sensitivity with no significant compromise

in specificity. We recently tested this assumption in an opportunistic screening setting, in which blood and stool samples were collected from outpatients and inpatients coming to the GI departments of three Chinese hospitals<sup>[55]</sup>. Table 6 shows the test results from the screening. SEPT9, FIT or CEA alone detected 77.0%, 74.6% and 41.3% of CRC cases, respectively, while the combination of the three increased the sensitivity to 97.2%, and SEPT9 plus FIT exhibited a sensitivity of 94.4%. Since CEA is more sensitive to late-stage CRC than early-stage CRC, and no significant difference was found between SEPT9 + FIT+ CEA and SEPT9 + FIT, we recommend SEPT9 + FIT as a routine method for CRC screening.

## CONCLUSION

The FIT, fecal DNA and the SEPT9 tests are IVD tests currently used for CRC screening. FIT tests exhibited satisfactory sensitivity and specificity with low costs and therefore become the major screening test for CRC at the moment. The sensitivity of the fecal DNA test appeared to be very high due to combination of multiple methods while its high cost is an obstacle preventing the test from broad use. Both sensitivity and specificity for the SEPT9 test in CRC screening were lower than those of the FIT and fecal DNA test, but it showed high compliance with promising future if its accuracy can be improved. Combined tests with multiple markers should be a future direction in CRC screening, however, some hurdles, such as technical integration, test/interpretation optimization, and high costs, etc, need to be overcome before they can be used in large-scale CRC screening aiming at asymptomatic average-risk population.

## REFERENCES

- 1 **American Cancer Society.** Cancer Facts & Figures 2013. American Cancer Society, GA, United States. Available from: URL: <http://www.cancer.org/research/cancerfactsstatistics/cancerfactsfigures2013/index>
- 2 **Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JW, Comber H, Forman D, Bray F.** Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer* 2013; **49**: 1374-1403 [PMID: 23485231 DOI: 10.1016/j.ejca.2012.12.027]
- 3 The National Cancer Center and Disease Prevention and Control Bureau: The 2012 Annual Cancer Report of China. National Cancer Center and Disease Prevention and Control Bureau, Beijing, China
- 4 **American Cancer Society.** Colorectal Cancer Facts & Figures 2011-2013. American Cancer Society, GA, United States. Available from: URL: <http://www.cancer.org/research/cancerfactsstatistics/colorectal-cancer-facts-figures>
- 5 National Cancer Institute, PDQ, Treatment, Health Professionals. Survival Rate for Colorectal Cancer by Stage 1999. National Cancer Institute, Bethesda, MD, United States
- 6 **Mandel JS, Bond JH, Church TR, Snover DC, Bradley GM, Schuman LM, Ederer F.** Reducing mortality from colorectal cancer by screening for fecal occult blood. Minnesota Colon Cancer Control Study. *N Engl J Med* 1993; **328**: 1365-1371 [PMID: 8474513 DOI: 10.1056/NEJM199305133281901]
- 7 **Hardcastle JD, Chamberlain JO, Robinson MH, Moss SM, Amar SS, Balfour TW, James PD, Mangham CM.** Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. *Lancet* 1996; **348**: 1472-1477 [PMID: 8942775 DOI: 10.1016/S0140-6736(96)03386-7]
- 8 **Kronborg O, Fenger C, Olsen J, Jørgensen OD, Søndergaard O.** Randomised study of screening for colorectal cancer with faecal-occult-blood test. *Lancet* 1996; **348**: 1467-1471 [PMID: 8942774 DOI: 10.1016/S0140-6736(96)03430-7]
- 9 **Lieberman DA.** Clinical practice. Screening for colorectal cancer. *N Engl J Med* 2009; **361**: 1179-1187 [PMID: 19759380 DOI: 10.1056/NEJMcp0902176]
- 10 **Ahlquist DA, Wieand HS, Moertel CG, McGill DB, Loprinzi CL, O'Connell MJ, Mailliard JA, Gerstner JB, Pandya K, Ellefson RD.** Accuracy of fecal occult blood screening for colorectal neoplasia. A prospective study using Hemoccult and HemoQuant tests. *JAMA* 1993; **269**: 1262-1267 [PMID: 8437303]
- 11 **Ahlquist DA, Shuber AP.** Stool screening for colorectal cancer: evolution from occult blood to molecular markers. *Clin Chim Acta* 2002; **315**: 157-168 [PMID: 11728417 DOI: 10.1016/S0009-8981(01)00712-4]
- 12 **Allison JE, Tekawa IS, Ransom LJ, Adrain AL.** A comparison of fecal occult-blood tests for colorectal-cancer screening. *N Engl J Med* 1996; **334**: 155-159 [PMID: 8531970 DOI: 10.1056/NEJM199601183340304]
- 13 **Imperiale TF, Ransohoff DF, Itzkowitz SH, Turnbull BA, Ross ME.** Fecal DNA versus fecal occult blood for colorectal-cancer screening in an average-risk population. *N Engl J Med* 2004; **351**: 2704-2714 [PMID: 15616205 DOI: 10.1056/NEJMoa033403]
- 14 **Lee JK, Liles EG, Bent S, Levin TR, Corley DA.** Accuracy of fecal immunochemical tests for colorectal cancer: systematic review and meta-analysis. *Ann Intern Med* 2014; **160**: 171 [PMID: 24658694 DOI: 10.7326/M13-1484]
- 15 **Allison JE, Sakoda LC, Levin TR, Tucker JP, Tekawa IS, Cuff T, Pauly MP, Shlager L, Palitz AM, Zhao WK, Schwartz JS, Ransohoff DF, Selby JV.** Screening for colorectal neoplasms with new fecal occult blood tests: update on performance characteristics. *J Natl Cancer Inst* 2007; **99**: 1462-1470 [PMID: 17895475 DOI: 10.1093/jnci/djm150]
- 16 **Cheng TI, Wong JM, Hong CF, Cheng SH, Cheng TJ, Shieh MJ, Lin YM, Tso CY, Huang AT.** Colorectal cancer screening in asymptomatic adults: comparison of colonoscopy, sigmoidoscopy and fecal occult blood tests. *J Formos Med Assoc* 2002; **101**: 685-690 [PMID: 12517041]
- 17 **Nakama H, Yamamoto M, Kamijo N, Li T, Wei N, Fattah AS, Zhang B.** Colonoscopic evaluation of immunochemical fecal occult blood test for detection of colorectal neoplasia. *Hepatogastroenterology* 1999; **46**: 228-231 [PMID: 10228797]
- 18 **Nakama H, Kamijo N, Abdul Fattah AS, Zhang B.** Validity of immunological faecal occult blood screening for colorectal cancer: a follow up study. *J Med Screen* 1996; **3**: 63-65 [PMID: 8849761]
- 19 **Parra-Blanco A, Gimeno-García AZ, Quintero E, Nicolás D, Moreno SG, Jiménez A, Hernández-Guerra M, Carrillo-Palau M, Eishi Y, López-Bastida J.** Diagnostic accuracy of immunochemical versus guaiac faecal occult blood tests for colorectal cancer screening. *J Gastroenterol* 2010; **45**: 703-712 [PMID: 20157748 DOI: 10.1007/s00535-010-0214-8]
- 20 **Chiu HM, Lee YC, Tu CH, Chen CC, Tseng PH, Liang JT, Shun CT, Lin JT, Wu MS.** Association between early stage colon neoplasms and false-negative results from the fecal immunochemical test. *Clin Gastroenterol Hepatol* 2013; **11**: 832-8.e1-832-8.e2 [PMID: 23376002 DOI: 10.1016/j.cgh.2013.01.013]
- 21 **Chiang TH, Lee YC, Tu CH, Chiu HM, Wu MS.** Performance of the immunochemical fecal occult blood test in predicting lesions in the lower gastrointestinal tract. *CMAJ* 2011; **183**: 1474-1481 [PMID: 21810951 DOI: 10.1503/cmaj.101248]
- 22 **Sohn DK, Jeong SY, Choi HS, Lim SB, Huh JM, Kim DH, Kim DY, Kim YH, Chang HJ, Jung KH, Ahn JB, Kim HK, Park JG.** Single immunochemical fecal occult blood test for detection of colorectal neoplasia. *Cancer Res Treat* 2005; **37**: 20-23 [PMID: 19956505 DOI: 10.4143/crt.2005.37.1.20]
- 23 **Levi Z, Birkenfeld S, Vilkin A, Bar-Chana M, Lifshitz I, Chared M, Maoz E, Niv Y.** A higher detection rate for colorectal cancer and advanced adenomatous polyp for screening with immunochemical

- fecal occult blood test than guaiac fecal occult blood test, despite lower compliance rate. A prospective, controlled, feasibility study. *Int J Cancer* 2011; **128**: 2415-2424 [PMID: 20658527 DOI: 10.1002/ijc.25574]
- 24 **Levi Z**, Rozen P, Hazazi R, Vilkin A, Waked A, Maoz E, Birkenfeld S, Leshno M, Niv Y. A quantitative immunochemical fecal occult blood test for colorectal neoplasia. *Ann Intern Med* 2007; **146**: 244-255 [PMID: 17310048 DOI: 10.7326/0003-4819-146-4-200702200-00003]
- 25 **Morikawa T**, Kato J, Yamaji Y, Wada R, Mitsushima T, Shiratori Y. A comparison of the immunochemical fecal occult blood test and total colonoscopy in the asymptomatic population. *Gastroenterology* 2005; **129**: 422-428 [PMID: 16083699 DOI: 10.1053/j.gastro.2005.05.056]
- 26 **Launoy GD**, Bertrand HJ, Berchi C, Talbourdet VY, Guizard AV, Bouvier VM, Caces ER. Evaluation of an immunochemical fecal occult blood test with automated reading in screening for colorectal cancer in a general average-risk population. *Int J Cancer* 2005; **115**: 493-496 [PMID: 15700317 DOI: 10.1002/ijc.20921]
- 27 **Itoh M**, Takahashi K, Nishida H, Sakagami K, Okubo T. Estimation of the optimal cut off point in a new immunological faecal occult blood test in a corporate colorectal cancer screening programme. *J Med Screen* 1996; **3**: 66-71 [PMID: 8849762]
- 28 **Nakazato M**, Yamano HO, Matsushita HO, Sato K, Fujita K, Yamanaka Y, Imai Y. Immunologic fecal occult blood test for colorectal cancer screening. *JMAJ* 2006; **49**: 203-207
- 29 **Park DI**, Ryu S, Kim YH, Lee SH, Lee CK, Eun CS, Han DS. Comparison of guaiac-based and quantitative immunochemical fecal occult blood testing in a population at average risk undergoing colorectal cancer screening. *Am J Gastroenterol* 2010; **105**: 2017-2025 [PMID: 20502450 DOI: 10.1038/ajg.2010.179]
- 30 **de Wijkerslooth TR**, Stoop EM, Bossuyt PM, Meijer GA, van Ballegooijen M, van Roon AH, Stegeman I, Kraaijenhagen RA, Fockens P, van Leerdam ME, Dekker E, Kuipers EJ. Immunochemical fecal occult blood testing is equally sensitive for proximal and distal advanced neoplasia. *Am J Gastroenterol* 2012; **107**: 1570-1578 [PMID: 22850431 DOI: 10.1038/ajg.2012.249]
- 31 **Brenner H**, Tao S. Superior diagnostic performance of faecal immunochemical tests for haemoglobin in a head-to-head comparison with guaiac based faecal occult blood test among 2235 participants of screening colonoscopy. *Eur J Cancer* 2013; **49**: 3049-3054 [PMID: 23706981 DOI: 10.1016/j.ejca.2013.04.023]
- 32 **Imperiale TF**, Ransohoff DF, Itzkowitz SH, Levin TR, Lavin P, Lidgard GP, Ahlquist DA, Berger BM. Multitarget stool DNA testing for colorectal-cancer screening. *N Engl J Med* 2014; **370**: 1287-1297 [PMID: 24645800 DOI: 10.1056/NEJMoa1311194]
- 33 **Church TR**, Wandell M, Lofton-Day C, Mongin SJ, Burger M, Payne SR, Castaños-Vélez E, Blumenstein BA, Rösch T, Osborn N, Snover D, Day RW, Ransohoff DF. Prospective evaluation of methylated SEPT9 in plasma for detection of asymptomatic colorectal cancer. *Gut* 2014; **63**: 317-325 [PMID: 23408352 DOI: 10.1136/gutjnl-2012-304149]
- 34 **Binefa G**, Rodríguez-Moranta F, Teule A, Medina-Hayas M. Colorectal cancer: from prevention to personalized medicine. *World J Gastroenterol* 2014; **20**: 6786-6808 [PMID: 24944469 DOI: 10.3748/wjg.v20.i22.6786]
- 35 **Dhaliwal A**, Vlachostergios PJ, Oikonomou KG, Moshenyat Y. Fecal DNA testing for colorectal cancer screening: Molecular targets and perspectives. *World J Gastrointest Oncol* 2015; **7**: 178-183 [PMID: 26483873]
- 36 **Lofton-Day C**, Model F, Devos T, Tetzner R, Distler J, Schuster M, Song X, Lesche R, Liebenberg V, Ebert M, Molnar B, Grützmann R, Pilarsky C, Sledziewski A. DNA methylation biomarkers for blood-based colorectal cancer screening. *Clin Chem* 2008; **54**: 414-423 [PMID: 18089654 DOI: 10.1373/clinchem.2007.095992]
- 37 **Grützmann R**, Molnar B, Pilarsky C, Habermann JK, Schlag PM, Saeger HD, Miehlke S, Stolz T, Model F, Roblick UJ, Bruch HP, Koch R, Liebenberg V, Devos T, Song X, Day RH, Sledziewski AZ, Lofton-Day C. Sensitive detection of colorectal cancer in peripheral blood by septin 9 DNA methylation assay. *PLoS One* 2008; **3**: e3759 [PMID: 19018278 DOI: 10.1371/journal.pone.0003759]
- 38 **deVos T**, Tetzner R, Model F, Weiss G, Schuster M, Distler J, Steiger KV, Grützmann R, Pilarsky C, Habermann JK, Flesher PR, Oubre BM, Day R, Sledziewski AZ, Lofton-Day C. Circulating methylated SEPT9 DNA in plasma is a biomarker for colorectal cancer. *Clin Chem* 2009; **55**: 1337-1346 [PMID: 19406918 DOI: 10.1373/clinchem.2008.115808]
- 39 **Tänzer M**, Balluff B, Distler J, Hale K, Leodolter A, Röcken C, Molnar B, Schmid R, Lofton-Day C, Schuster T, Ebert MP. Performance of epigenetic markers SEPT9 and ALX4 in plasma for detection of colorectal precancerous lesions. *PLoS One* 2010; **5**: e9061 [PMID: 20140221 DOI: 10.1371/journal.pone.0009061]
- 40 **Warren JD**, Xiong W, Bunker AM, Vaughn CP, Furtado LV, Roberts WL, Fang JC, Samowitz WS, Heichman KA. Septin 9 methylated DNA is a sensitive and specific blood test for colorectal cancer. *BMC Med* 2011; **9**: 133 [PMID: 22168215 DOI: 10.1186/1741-7015-9-133]
- 41 **Tóth K**, Sipos F, Kalmár A, Patai AV, Wichmann B, Stoehr R, Golcher H, Schellerer V, Tulassay Z, Molnár B. Detection of methylated SEPT9 in plasma is a reliable screening method for both left- and right-sided colon cancers. *PLoS One* 2012; **7**: e46000 [PMID: 23049919 DOI: 10.1371/journal.pone.0046000]
- 42 **Weiss G**, Rösch T. Potential of a new blood test for colorectal cancer screening - the Septin 9 gene biomarker. *Eur Oncol* 2010; **6**: 51-54 [DOI: 10.17925/eoh.2010.06.1.51]
- 43 **Potter NT**, Hurban P, White MN, Whitlock KD, Lofton-Day CE, Tetzner R, Koenig T, Quigley NB, Weiss G. Validation of a real-time PCR-based qualitative assay for the detection of methylated SEPT9 DNA in human plasma. *Clin Chem* 2014; **60**: 1183-1191 [PMID: 24938752 DOI: 10.1373/clinchem.2013.221044]
- 44 **Johnson DA**, Barclay RL, Mergener K, Weiss G, König T, Beck J, Potter NT. Plasma Septin9 versus fecal immunochemical testing for colorectal cancer screening: a prospective multicenter study. *PLoS One* 2014; **9**: e98238 [PMID: 24901436 DOI: 10.1371/journal.pone.0098238]
- 45 **Tóth K**, Wasserkort R, Sipos F, Kalmár A, Wichmann B, Leiszter K, Valcz G, Juhász M, Miheller P, Patai ÁV, Tulassay Z, Molnár B. Detection of methylated septin 9 in tissue and plasma of colorectal patients with neoplasia and the relationship to the amount of circulating cell-free DNA. *PLoS One* 2014; **9**: e115415 [PMID: 25526039 DOI: 10.1371/journal.pone.0115415]
- 46 **Jin P**, Kang Q, Wang X, Yang L, Yu Y, Li N, He YQ, Han X, Hang J, Zhang J, Song L, Han Y, Sheng JQ. Performance of a second-generation methylated SEPT9 test in detecting colorectal neoplasm. *J Gastroenterol Hepatol* 2015; **30**: 830-833 [PMID: 25471329 DOI: 10.1111/jgh.12855]
- 47 **Lee HS**, Hwang SM, Kim TS, Kim DW, Park DJ, Kang SB, Kim HH, Park KU. Circulating methylated septin 9 nucleic acid in the plasma of patients with gastrointestinal cancer in the stomach and colon. *Transl Oncol* 2013; **6**: 290-296 [PMID: 23730408 DOI: 10.1593/tlo.13118]
- 48 **Li Y**, Song L, Gong Y, He B. Detection of colorectal cancer by DNA methylation biomarker SEPT9: past, present and future. *Biomark Med* 2014; **8**: 755-769 [PMID: 25123042 DOI: 10.2217/bmm.14.8]
- 49 **Song L**, Li Y. SEPT9: A Specific Circulating Biomarker for Colorectal Cancer. *Adv Clin Chem* 2015; **72**: 171-204 [PMID: 26471083 DOI: 10.1016/bs.acc.2015.07.004]
- 50 **Adler A**, Geiger S, Keil A, Bias H, Schatz P, deVos T, Dhein J, Zimmermann M, Tauber R, Wiedenmann B. Improving compliance to colorectal cancer screening using blood and stool based tests in patients refusing screening colonoscopy in Germany. *BMC Gastroenterol* 2014; **14**: 183 [PMID: 25326034 DOI: 10.1186/1471-230X-14-183]
- 51 **Wild N**, Andres H, Rollinger W, Krause F, Dilba P, Tacke M, Karl J. A combination of serum markers for the early detection of colorectal cancer. *Clin Cancer Res* 2010; **16**: 6111-6121 [PMID: 20798228 DOI: 10.1158/1078-0432.CCR-10-0119]
- 52 **Chen JS**, Chen KT, Fan WC, Yu JS, Chang YS, Chan EC. Combined analysis of survivin autoantibody and carcinoembryonic antigen biomarkers for improved detection of colorectal cancer. *Clin Chem Lab Med* 2010; **48**: 719-725 [PMID: 20178447 DOI: 10.1515/CCLM.2010.123]
- 53 **Yu H**, Son GM, Joh YG. The clinical significance of preoperative serum levels of carbohydrate antigen 19-9 in colorectal cancer.

- J Korean Surg Soc* 2013; **84**: 231-237 [PMID: 23577318 DOI: 10.4174/jkss.2013.84.4.231]
- 54 **Fu H**, Wang W, Cai Q. The application of combined detection of serum tumor markers levels of patients with colon cancer. *Chin J Clin* 2012; **6**: 5087-5090
- 55 **Wu D**, Zhou G, Jin P, Zhu J, Li S, Wu Q, Wang G, Sheng J, Wang J, Song L, Han X, Qian J. Detection of Colorectal Cancer Using a Simplified SEPT9 Gene Methylation Assay Is a Reliable Method for Opportunistic Screening. *J Mol Diagn* 2016; **18**: 535-545 [PMID: 27133379 DOI: 10.1016/j.jmoldx.2016.02.005]
- P- Reviewer:** Guo Q, Jeong KY, Noshiro H **S- Editor:** Kong JX  
**L- Editor:** A **E- Editor:** Wu HL





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

