

Dear Editor,

Please find enclosed the edited manuscript in Word format.

Title: *Comparison of Analgesic Effects between Electro-acupuncture and Moxibustion in Rats with Visceral Hyperactivity*

Author: Ji-Meng Zhao, Liang Li, Liu Chen, Yin Shi, Yu-Wei Li, Hai-Xia Shang, Lu-Yi Wu, Zhi-Jun Weng, Chun-Hui Bao, Huan-Gan Wu

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 30662

1. Revision has been made according to the suggestions of the reviewers:

Reviewer's code: 00058573

COMMENTS TO AUTHORS

Well conducted and quite meticulous study. However the major drawback is that along with 4 intervention groups, a NORMAL control group would have added much quality to the study. How can one be sure that whether the placebo control group might have shown better results than all the 4 intervention groups.

Response: We thank very much the reviewer for the suggestion. Actually, there are six groups in the study, (1) EA 1 mA group, (2) EA 3 mA group, (3) Mox 43°C group, (4) Mox 46°C group, (5) Model Control group, (6) Normal Control group; and all the data of the other groups were have compared with the Normal Control group.

Sincerely yours,

Ji-Meng Zhao

Shanghai Institute of Acupuncture-Moxibustion and Meridians

No. 650 South Wanping Road, Xuhui District, Shanghai 200030, China

Tel:+86-21-64395973