

**Name of Journal:** *World Journal of Hepatology*

**ESPS Manuscript NO:** 30229

**Manuscript Type:** Original Article

*Prospective Study*

**Is hepatic steatosis associated with left ventricular mass index increase in the general population?**

**Katharina Piontek, Carsten O Schmidt, Sebastian E Baumeister, Markus M Lerch, Julia Mayerle, Marcus Dörr, Stephan B Felix, Henry Völzke**

**Abstract**

## Match Overview

- Crossref** 77 words  
Katharina Lau. "The association between fatty liver diseases ... and blood pressure in a population-based prospective longit" **2%**
- Internet** 36 words  
crawled on 10-Aug-2016  
[respiratory-research.biomedcentral.com](http://respiratory-research.biomedcentral.com) **1%**
- Internet** 29 words  
crawled on 23-Feb-2017  
[spandidos-publications.com](http://spandidos-publications.com) **1%**
- Internet** 23 words  
crawled on 01-Jun-2010  
[pt.wkhealth.com](http://pt.wkhealth.com) **<1%**
- Crossref** 20 words  
"Liver", *The American Journal of Gastroenterology*, 2015. **<1%**



Is hepatic steatosis associated with left ventricular mass index increase in t



All

News

Images

Shopping

Videos

More ▾

Search tools

About 2,770,000 results (1.02 seconds)

## Association between Hepatic Triglyceride Content and Left Ventricular ...

[pubs.rsna.org/doi/full/10.../radiol.2015150035?...](https://pubs.rsna.org/doi/full/10.../radiol.2015150035?...) Radiological Society of North America ▾

We hypothesized that hepatic triglyceride content is **associated with LV** ..... are reflected by decreased LV end-diastolic **index** and **increased LV mass** in obesity. .... triglyceride content: prevalence of **hepatic steatosis** in the **general population**.

## Full Text - Arteriosclerosis, Thrombosis, and Vascular Biology

[atvb.ahajournals.org/.../1690.full](https://atvb.ahajournals.org/.../1690.full) ▾ Arteriosclerosis, Thrombosis, and Vascular Biology ▾

by MRP Markus - 2013 - Cited by 19 - Related articles

May 16, 2013 - Individuals with **hepatic steatosis** have **increased** intima media ... left atrial diameter and **left ventricular** mass than individuals without **hepatic steatosis** (Table 1). ... body **mass index** as a potential cofounder instead of waist-to-height ratio. .... In a sample from the **general population** (n=2212), we observed an ...

## Hepatic Rather Than Cardiac Steatosis Relates to Glucose Intolerance ...

[www.ncbi.nlm.nih.gov/.../PMC3951459/](https://www.ncbi.nlm.nih.gov/.../PMC3951459/) National Center for Biotechnology Information ▾

by Y Winhofer - 2014 - Cited by 3 - Related articles

Mar 12, 2014 - **Increased** myocardial lipid content (cardiac **steatosis**) is regarded as an ... was found to be **associated** with elevated **left ventricular mass** [7] and diastolic ... display a relatively young **population** at high risk of developing diabetes. .... In order to evaluate the accuracy of the **fatty liver index** in our study group, ...

## Non-alcoholic fatty liver disease may develop in individuals with ...

[www.ncbi.nlm.nih.gov/.../PMC39593...](https://www.ncbi.nlm.nih.gov/.../PMC39593...) ▾ National Center for Biotechnology Information ▾

by E Margariti - 2012 - Cited by 23 - Related articles

NAFLD diagnosis required elevated ALT and/or GGT, **hepatic steatosis** on .... non-alcoholic fatty liver disease and normal or **increased** body **mass index** (BMI, kg/m<sup>2</sup>) ... heart disease are strongly **associated** with obesity in the **general population** ... NAFLD patients have been reported to have early **left ventricular** diastolic ...

## Risk of cardiovascular, cardiac and arrhythmic complications in ... - NCBI

[All](#)[News](#)[Images](#)[Shopping](#)[Videos](#)[More](#)[Settings](#)[Tools](#)

About 121,000 results (1.08 seconds)

[Non-alcoholic fatty liver disease and its relationship with ... - Gut](#)[gut.bmj.com/content/early/2017/03/17/gutjnl-2017-313884.full.pdf](https://gut.bmj.com/content/early/2017/03/17/gutjnl-2017-313884.full.pdf)

17 Mar 2017 - **left ventricular** morphology and/or diastolic dysfunction,36–39 ... **Population-** based cohort studies, such as the National Health ... **general** NAFLD is **associated** with an **increased** risk of CVD, ..... BMI, body **mass index**; MELD, model for end-stage **liver** disease; MetS, metabolic syndrome; NA, not applicable;....

[Association of nonalcoholic fatty liver disease with subclinical ... - NCBI](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4549239/>by LB VanWagner - 2015 - [Cited by 37](#) - [Related articles](#)

26 Jun 2015 - Subclinical **left ventricular** (LV) systolic dysfunction was defined using ... When adjusted for HF risk factors or body **mass index**, NAFLD remained **associated** with subclinical ... and function apply to the **general** adult **population** is unknown. .... **Hepatic steatosis** is **associated** with an **increased** risk of carotid ...

[Non-alcoholic fatty liver disease may develop in individuals with ...](#)<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#)by E Margariti - 2012 - [Cited by 26](#) - [Related articles](#)

NAFLD diagnosis required elevated ALT and/or GGT, **hepatic steatosis** on .... non-alcoholic fatty liver disease and normal or **increased** body **mass index** (BMI, kg/m<sup>2</sup>) ... heart disease are strongly **associated** with obesity in the **general population** ... NAFLD patients have been reported to have early **left ventricular** diastolic ...

[Nonalcoholic fatty liver disease and vascular disease: State-of-the-art](#)<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#)by S Fargion - 2014 - [Cited by 66](#) - [Related articles](#)

7 Oct 2014 - Nonalcoholic fatty **liver** disease (NAFLD), the most common of ... endothelial dysfunction, arterial stiffness, impaired **left ventricular** ... NAFLD affects about 20%-30% of the **general population** and is closely **related** to insulin resistance and ... but not simple



[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 121,000 条结果 (用时 0.74 秒)

### Non-alcoholic fatty liver disease - the heart of the matter - NCBI

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral\(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) - 翻译此页

作者: H Azzam - 2015 - 被引用次数: 10 - 相关文章

2015年6月8日 - MS is **associated** with a risk of cardiovascular disease and is a common early ... to those with **steatosis** alone independent of age or body **mass index** (BMI)[13]. ... The NAFLD patients had altered **left ventricular** (LV) geometry and early ... with prevalence estimates of 2% to 4% in the **general population**.

### Association of nonalcoholic fatty liver disease with subclinical ... - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4549239/> - 翻译此页

作者: LB VanWagner - 2015 - 被引用次数: 40 - 相关文章

2015年6月26日 - Subclinical **left ventricular** (LV) systolic dysfunction was defined using ... When adjusted for HF risk factors or body **mass index**, NAFLD remained **associated** with subclinical ... and function apply to the **general adult population** is unknown. .... **Hepatic steatosis** is **associated** with an **increased** risk of carotid ...

### Fatty liver linked to heart failure in obese patients - AuntMinnie.com

[www.auntminnie.com/Redirect/Redirect.aspx?ItemId=113247](http://www.auntminnie.com/Redirect/Redirect.aspx?ItemId=113247) - 翻译此页

2016年1月27日 - Fatty liver is an independent risk factor for heart failure in obese patients -- that is, those with a body **mass index** of 30 or higher, according to a new ... nonalcoholic fatty liver disease, also called **hepatic steatosis**, could ... **associated** with a decrease in mean **left ventricular** diastolic function in obese patients.

### Association between Hepatic Triglyceride Content and Left Ventricular ...

[pubs.rsna.org/doi/full/10.1148/radiol.2015150035?mobileUi=0](http://pubs.rsna.org/doi/full/10.1148/radiol.2015150035?mobileUi=0) - 翻译此页

NAFLD covers a spectrum of simple **steatosis** to nonalcoholic steatohepatitis and ... We hypothesized that **hepatic** triglyceride content is **associated with** LV ... content and diastolic function in a large sample