

Dear Yuan Qi

Thank you for accepting our letter for publication subject to changes. This letter details the changes that have been made. These are highlighted in the text which has been uploaded in its modified form.

The text has been amended to contain full names of all authors and we have provided the full address of our institution.

Campbell Jennifer Anne, Irvine Andrew, Hopper Andrew Derek

**Department of Gastroenterology, Royal Hallamshire Hospital, Glossop Road,
Sheffield, S10 2JF, United Kingdom**

The conflict of interest statement has been inserted and also signed and submitted online.

Conflict-of-interest statement: All authors are employed by Sheffield Teaching Hospitals NHS Foundation Trust. No funding was required for this study and there are no competing interests to declare.

The references have been amended to contain PMID and DOI in the format requested. The first page of reference 4 has been uploaded as it is a guideline and does not have a PMID.

1. **Cheriyana DG, Byrne MF.** Propofol use in endoscopic retrograde cholangiopancreatography and endoscopic ultrasound. *World J Gastroenterol* 2014; 20: 5171-6 [PMID: 4017032 DOI: 10.3748/wjg.v20.i18.5171]
2. **Fatima H, DeWitt J, LeBlanc J, Sherman S, McGreevy K, Imperiale TF.** Nurse-administered propofol sedation for upper endoscopic ultrasonography. *Am J Gastroenterol* 2008; 103: 1649-56 [PMID: 18557709 DOI: 10.1111/j1572-0241.2008.01906.x]
3. **Nayar DS, Guthrie WG, Goodman A, Lee Y, Feuerman M, Scheinberg L, Gress FG.** Comparison of propofol deep sedation versus moderate sedation during endoscopy. *Dig Dis Sci* 2010; 55: 2537-44 [PMID 20635148 DOI: 10.1007/s10620-010-1308-0]
4. **Tomlinson A, Green J, Cairns SR, Cressey D, Smith, I, Peacock J.** Guidance for the use of propofol sedation for adult patients undergoing Endoscopic Retrograde Cholangiopancreatography (ERCP) and other complex upper GI endoscopic procedures. On behalf of the Joint Royal College of Anaesthetists (RCoA) and British Society of Gastroenterology (BSG) Working Party
<http://www.bsg.org.uk/clinical-guidance/endoscopy/guidance-for-the-use-of-propofol-sedation-for-adults-undergoing-endoscopic-retrograde-cholangiopancreatography-ercp.html>

5. **Ko** HH, Zhang H, Telford JJ, Enns R. Factors influencing patient satisfaction when undergoing endoscopic procedures. *Gastrointest Endosc* 2009 ;69 :883-91, quiz 91 e1. [PMID: 19152911 DOI: 10.1016/j.gie.2008.06.024]
6. **Pena** LR, Mardini HE, Nickl NJ. Development of an instrument to assess and predict satisfaction and poor tolerance among patients undergoing endoscopic procedures. *Dig Dis Sci* 2005 ;50 :1860-71. [PMID 16187188 DOI: 10.1007/s10620-005-2952-7]
7. **Salmon** P, Shah R, Berg S, Williams C. Evaluating customer satisfaction with colonoscopy. *Endoscopy* 1994; 26: 342-6. [PMID: 8076565 DOI: 10.1055/s-2007-1008988]
8. **Dewitt** J, McGreevy K, Sherman S, Imperiale TF. Nurse-administered propofol sedation compared with midazolam and meperidine for EUS: a prospective, randomized trial. *Gastrointest Endosc* 2008 ;68: 499-509. [PMID 18561925 DOI: 10.1016/j.gie.2008.02.092]
9. **Mortensen** MB, Fristrup C, Holm FS, Pless R, Durup J, Ainsworth AP, Nielsen HO, Hovendal C. Prospective evaluation of patient tolerability, satisfaction with patient information, and complications in endoscopic ultrasonography. *Endoscopy* 2005 ;37 :146-53. [PMID 15692930 DOI: 10.1055/s-2005-861142]

We thank you again for your comments and are looking forward to seeing our letter in press.

Yours sincerely

Jennifer Anne Campbell