

ESPS PEER-REVIEW REPORT

Name of journal: World Journal of Gastrointestinal Endoscopy

ESPS manuscript NO: 31806

Title: Endoscopic resolution and recurrence of Gastric Antral Vascular Ectasia after serial treatment with argon plasma coagulation

Reviewer's code: 03397142

Reviewer's country: Malaysia

Science editor: Ya-Juan Ma

Date sent for review: 2016-12-09 14:43

Date reviewed: 2016-12-28 16:04

CLASSIFICATION	LANGUAGE EVALUATION	SCIENTIFIC MISCONDUCT	CONCLUSION
<input type="checkbox"/> Grade A: Excellent	<input type="checkbox"/> Grade A: Priority publishing	Google Search:	<input type="checkbox"/> Accept
<input type="checkbox"/> Grade B: Very good	<input checked="" type="checkbox"/> Grade B: Minor language polishing	<input type="checkbox"/> The same title	<input type="checkbox"/> High priority for publication
<input checked="" type="checkbox"/> Grade C: Good	<input type="checkbox"/> Grade C: A great deal of language polishing	<input type="checkbox"/> Duplicate publication	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade D: Fair	<input type="checkbox"/> Grade D: Rejected	<input checked="" type="checkbox"/> No	<input checked="" type="checkbox"/> Minor revision
<input type="checkbox"/> Grade E: Poor		BPG Search:	<input type="checkbox"/> Major revision
		<input type="checkbox"/> The same title	
		<input type="checkbox"/> Duplicate publication	
		<input type="checkbox"/> Plagiarism	
		<input checked="" type="checkbox"/> No	

COMMENTS TO AUTHORS

The authors have looked into the outcomes of 20 patients with Gastric Antral Vascular Ectasia (GAVE) who underwent endoscopic Argon Plasma Coagulation (APC) treatment and concluded that endoscopic resolution rate of GAVE was low (40%) with a 25% recurrence rate after treatment with APC. The manuscript is interesting and provided some insight on the treatment of GAVE using APC. Can the authors please kindly clarify the results section in the abstract where you state 28 patients met study criteria but only 20 were included into the final analysis. 'Twenty-eight patients met study criteria and 20 (71%) patients had GAVE related symptoms.' Thank you for sharing your work.