

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 31900

Manuscript Type: Original Article

Retrospective Study

S-1 induced hepatic steatosis in patients with pancreatic cancer: retrospective analysis

Tsuji K et al. S-1 induced hepatic steatosis

Kunihiro Tsuji, Hisashi Doyama

Abstract

AIM

To determine whether S-1 induces hepatic steatosis in patients being treated for pancreatic cancer.

Match Overview

1	Internet 61 words crawled on 07-Jan-2017 bmjopenbeta.bmj.com	3%
2	Publications 35 words "Recent studies from Sungkyunkwan University, Department of Internal Medicine add new data to obesity", Hep	2%
3	Crossref 22 words Tanigawa, Nobuhiko, Hiroki Yamaue, Shigekazu Ohyama, Shinichi Sakuramoto, Takao Inada, Yasuhiro Kadera,	1%
4	Internet 20 words crawled on 24-Mar-2016 www.dovepress.com	1%
5	Crossref 19 words Hyodo, I. "A phase I study of S-1 combined with weekly cisplatin for metastatic gastric cancer in an outpatient setting"	1%
6	Publications 19 words "Low-carbohydrate diet burns more excess liver fat than low-calorie diets.", Nutrition Health Review, Spring 2010	1%
7	Crossref 13 words PD Peppercorn. "Demonstration of hepatic steatosis by computerized tomography in patients receiving 5-fluorouracil"	1%



S-1 induced hepatic steatosis in patients with pancreatic cancer: retrospec



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 40,800 条结果 (用时 0.97 秒)

Google 学术: S-1 induced hepatic steatosis in patients with pancreatic cancer: retrospective analysis

... pancreas: Usefulness in characterizing solid pancreatic ... - Lee - 被引用次数: 165

... and leucovorin for metastatic colorectal cancer - Saltz - 被引用次数: 3222

Adverse effects of reverse transcriptase inhibitors: ... - Brinkman - 被引用次数: 873

Pancreatic cancer and FOLFIRINOX: a new era and new questions

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: RDW Marsh - 2015 - 被引用次数: 15 - 相关文章

2015年2月19日 - With a median follow-up of 26.6 months and with 171 patients in each arm ... Table 1.

Management of FOLFIRINOX toxicity. If platelet counts are problematic (GTX) regimen for metastatic pancreatic cancer: a retrospective analysis. S. Development of hepatic steatosis after pancreatoduodenectomy.

Long-term influence of chemotherapy on steatosis-associated ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: SK Reddy - 2014 - 被引用次数: 4 - 相关文章

2014年5月6日 - On multivariable analysis, chemotherapy treatment was a negative predictive ... Among cancer patients with radiologic evidence of hepatic steatosis without ... These criteria included (1) the presence of other chronic diseases of the D. Gemcitabine-induced liver fibrosis in a patient with pancreatic cancer.

Pancreatic cancer: chemotherapy and radiotherapy - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页



S-1 induced hepatic steatosis in patients with pancreatic cancer: Retrospe



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 39,600 条结果 (用时 0.68 秒)

Google 学术: S-1 induced hepatic steatosis in patients with pancreatic cancer: Retrospective analysis

... **pancreas**: Usefulness in characterizing solid **pancreatic** ... - Lee - 被引用次数: 165

... and leucovorin for metastatic colorectal **cancer** - Saltz - 被引用次数: 3238

Adverse effects of reverse transcriptase inhibitors: ... - Brinkman - 被引用次数: 876

S-adenosylmethionine (SAME) therapy in liver disease: A review of ...

www.sciencedirect.com/science/article/pii/S0168827812004096 ▼ 翻译此页

作者: QM Anstee - 2012 - 被引用次数: 92 - 相关文章

2012年5月30日 - S-adenosyl-L-methionine (SAME; AdoMet) is an important, metabolically pleiotropic ... in the **liver** and is also present in the exocrine **pancreas** and prostate. ... animals are more sensitive to choline-deficient diet **induced steatosis**; and, 1). Confirming this, **liver** biopsies from **patients** with alcoholic hepatitis ...

Bortezomib and gemcitabine in relapsed or refractory Hodgkin's ...

<https://academic.oup.com/annonc/article-pdf/19/10/1759/.../mdn365.pdf> - 翻译此页

作者: JH Mendler - 2008 - 被引用次数: 39 - 相关文章

2008年5月25日 - J. H. Mendler, J. Kelly, S. Voci, D. Marquis, L. Rich, R. M. Rossi, S. H. ...

Department of Internal Medicine, James P. Wilmot **Cancer** Center, University of ... **Patients** received 3-week cycles of bortezomib **1** mg/m² ... **induction** therapy for high-risk **patients**, as well as for salvage **Retrospective analysis of.**

Pancreatic cancer and FOI FIRINOX: a new era and new questions

S-1 induced hepatic steatosis in patients with pancreatic cancer: Retrospective a

全部

图片

视频

新闻

地图

图书

找到约 38,000 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Google 学术: S-1 induced hepatic steatosis in patients with pancreatic cancer: Retrospective analysis](#)

[... pancreas: Usefulness in characterizing solid pancreatic ...](#) - Lee - 被引用次数: 165

[... and leucovorin for metastatic colorectal cancer](#) - Saltz - 被引用次数: 3242

[Adverse effects of reverse transcriptase inhibitors: ...](#) - Brinkman - 被引用次数: 878

[S-adenosylmethionine \(SAME\) therapy in liver disease: A review of ...](#)

www.sciencedirect.com/science/article/pii/S0168827812004096

30 May 2012 ... S-adenosyl-L-methionine (SAME; AdoMet) is an important, metabolically pleiotropic ... in the liver and is also present in the exocrine pancreas and prostate. ... animals are more sensitive to choline-deficient diet induced steatosis; and, 1). Confirming this, liver biopsies from patients with alcoholic hepatitis ...

[\[PDF\] Bortezomib and gemcitabine in relapsed or refractory Hodgkin's ...](#)

<https://academic.oup.com/annonc/article-pdf/19/10/1759/.../mdn365.pdf>

25 May 2008 ... J. H. Mendler, J. Kelly, S. Voci, D. Marquis, L. Rich, R. M. Rossi, S. H. ... Department of Internal Medicine, James P. Wilmot Cancer Center, University of ... Patients received 3-week cycles of bortezomib 1 mg/m2 ... induction therapy for high-risk patients, as well as for salvage Retrospective analysis of.

[Pancreatic cancer and FOLFIRINOX: a new era and new questions](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4472208/>

19 Feb 2015 ... With a median follow-up of 26.6 months and with 171 patients in each arm ... Table 1. Management of FOLFIRINOX toxicity. If platelet counts are A Whipple operation, in and of itself, leads to an increase in hepatic steatosis 32. (GTX) regimen for metastatic pancreatic cancer: a retrospective analysis.

[Pancreatic cancer: chemotherapy and radiotherapy - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3336926/>