

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2017 August 28; 23(32): 5829-6008



**EDITORIAL**

- 5829** Role of tissue microenvironment resident adipocytes in colon cancer

*Tabuso M, Homer-Vanniasinkam S, Adya R, Arasaraadnam RP*

**REVIEW**

- 5836** Ophthalmic manifestations in patients with inflammatory bowel disease: A review

*Troncoso LL, Biancardi AL, de Moraes Jr HV, Zaltman C*

- 5849** Laparoscopic appendectomy for acute appendicitis: How to discourage surgeons using inadequate therapy

*Hori T, Machimoto T, Kadokawa Y, Hata T, Ito T, Kato S, Yasukawa D, Aisu Y, Kimura Y, Sasaki M, Takamatsu Y, Kitano T, Hisamori S, Yoshimura T*

- 5860** Long non-coding RNAs in hepatocellular carcinoma: Potential roles and clinical implications

*Niu ZS, Niu XJ, Wang WH*

**MINIREVIEWS**

- 5875** Nano albumin bound-paclitaxel in pancreatic cancer: Current evidences and future directions

*Giordano G, Pancione M, Olivieri N, Parcesepe P, Velocci M, Di Raimo T, Coppola L, Toffoli G, D'Andrea MR*

**ORIGINAL ARTICLE****Basic Study**

- 5887** Comparison between tocotrienol and omeprazole on gastric growth factors in stress-exposed rats

*Nur Azlina MF, Qodriyah HMS, Chua KH, Kamisah Y*

- 5895** (-)-Epigallocatechin-3-gallate enhances poly I:C-induced interferon- $\lambda$ 1 production and inhibits hepatitis C virus replication in hepatocytes

*Wang YZ, Li JL, Wang X, Zhang T, Ho WZ*

- 5904** Effects and mechanism of adenovirus-mediated phosphatase and tension homologue deleted on chromosome ten gene on collagen deposition in rat liver fibrosis

*Xie SR, An JY, Zheng LB, Huo XX, Guo J, Shih D, Zhang XL*

**Retrospective Study**

- 5913** Integrating *TYMS*, *KRAS* and *BRAF* testing in patients with metastatic colorectal cancer

*Ntavatzikos A, Spathis A, Patapis P, Machairas N, Peros G, Konstantoudakis S, Leventakou D, Panayiotides IG, Karakitsos P, Koumariannou A*

**Clinical Trials Study**

- 5925 Characterizing gastrointestinal stromal tumors and evaluating neoadjuvant imatinib by sequencing of endoscopic ultrasound-biopsies

Hedenström P, Nilsson B, Demir A, Andersson C, Enlund F, Nilsson O, Sadik R

**Observational Study**

- 5936 Novel predictors for lymph node metastasis in submucosal invasive colorectal carcinoma

Yim K, Won DD, Lee IK, Oh ST, Jung ES, Lee SH

- 5945 Changes with aging in gastric biomarkers levels and in biochemical factors associated with *Helicobacter pylori* infection in asymptomatic Chinese population

Shan JH, Bai XJ, Han LL, Yuan Y, Sun XF

**Prospective Study**

- 5954 Modified *Helicobacter* test using a new test meal and a <sup>13</sup>C-urea breath test in *Helicobacter pylori* positive and negative dyspepsia patients on proton pump inhibitors

Tepeš B, Malfertheiner P, Labenz J, Aygen S

- 5962 Real time endoscopic ultrasound elastography and strain ratio in the diagnosis of solid pancreatic lesions

Okasha H, Elkholy S, El-Sayed R, Wifī MN, El-Nady M, El-Nabawi W, El-Dayem WA, Radwan MI, Farag A, El-sherif Y, Al-Gemeie E, Salman A, El-Sherbiny M, El-Mazny A, Mahdy RE

- 5969 Efficacy and safety of sofosbuvir and daclatasvir in treatment of kidney transplantation recipients with hepatitis C virus infection

Xue Y, Zhang LX, Wang L, Li T, Qu YD, Liu F

**Randomized Controlled Trial**

- 5977 New botanical drug, HL tablet, reduces hepatic fat as measured by magnetic resonance spectroscopy in patients with nonalcoholic fatty liver disease: A placebo-controlled, randomized, phase II trial

Jeong JY, Sohn JH, Baek YH, Cho YK, Kim Y, Kim H

**Randomized Clinical Trial**

- 5986 Randomized clinical trial comparing fixed-time split dosing and split dosing of oral Picosulfate regimen for bowel preparation

Jun JH, Han KH, Park JK, Seo HL, Kim YD, Lee SJ, Jun BK, Hwang MS, Park YK, Kim MJ, Cheon GJ

**META-ANALYSIS**

- 5994 Systematic review and meta-analysis of colon cleansing preparations in patients with inflammatory bowel disease

Restellini S, Kherad O, Bessissow T, Ménard C, Martel M, Taheri Tanjani M, Lakatos PL, Barkun AN

**CASE REPORT**

- 6003** Postoperative inflammation as a possible cause of portal vein thrombosis after irreversible electroporation for locally advanced pancreatic cancer

*Su JJ, Su M, Xu K, Wang PF, Yan L, Lu SC, Gu WQ, Chen YL*

**LETTERS TO THE EDITOR**

- 6007** Comment on "Efficacy and adverse events of cold *vs* hot polypectomy: A meta-analysis"

*Sun HH, Huang SL, Bai Y*

**ABOUT COVER**

Editorial board member of *World Journal of Gastroenterology*, Mitsushige Sugimoto, MD, PhD, Associate Professor, Division of Digestive Endoscopy, Shiga University of Medical Science Hospital, Otsu 520-2192, Japan

**AIMS AND SCOPE**

*World Journal of Gastroenterology* (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, and 28<sup>th</sup> each month. The *WJG* Editorial Board consists of 1375 experts in gastroenterology and hepatology from 68 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

**INDEXING/ABSTRACTING**

*World Journal of Gastroenterology* (*WJG*) is now indexed in Current Contents<sup>®</sup>/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch<sup>®</sup>), Journal Citation Reports<sup>®</sup>, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports<sup>®</sup> cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29<sup>th</sup> among 79 journals in gastroenterology and hepatology (quartile in category Q2).

**FLYLEAF**

**I-IX Editorial Board**

**EDITORS FOR THIS ISSUE**

**Responsible Assistant Editor:** *Xiang Li*  
**Responsible Electronic Editor:** *Fen-Fen Zhang*  
**Proofing Editor-in-Chief:** *Lian-Sheng Ma*

**Responsible Science Editor:** *Yuan Qi*  
**Proofing Editorial Office Director:** *Jin-Lei Wang*

**NAME OF JOURNAL**  
*World Journal of Gastroenterology*

**ISSN**  
ISSN 1007-9327 (print)  
ISSN 2219-2840 (online)

**LAUNCH DATE**  
October 1, 1995

**FREQUENCY**  
Weekly

**EDITORS-IN-CHIEF**  
**Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon**, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

**Stephen C Strom, PhD, Professor**, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

**Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology**, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

**EDITORIAL BOARD MEMBERS**  
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

**EDITORIAL OFFICE**  
Jin-Lei Wang, Director  
Yuan Qi, Vice Director  
Ze-Mao Gong, Vice Director  
*World Journal of Gastroenterology*  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

**PUBLISHER**  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpoffice@wjgnet.com](mailto:bpoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>

<http://www.wjgnet.com>

**PUBLICATION DATE**  
August 28, 2017

**COPYRIGHT**  
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

**SPECIAL STATEMENT**  
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

**INSTRUCTIONS TO AUTHORS**  
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

**ONLINE SUBMISSION**  
<http://www.f6publishing.com>



## Basic Study

# Comparison between tocotrienol and omeprazole on gastric growth factors in stress-exposed rats

Mohd Fahami Nur Azlina, Hj Mohd Saad Qodriyah, Kien Hui Chua, Yusof Kamisah

Mohd Fahami Nur Azlina, Hj Mohd Saad Qodriyah, Yusof Kamisah, Department of Pharmacology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur 56000, Malaysia

Kien Hui Chua, Department of Physiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur 56000, Malaysia

**Author contributions:** Nur Azlina MF, Qodriyah HMS, Chua KH and Kamisah Y designed the research; Nur Azlina MF performed the research and wrote the paper.

**Supported by** Universiti Kebangsaan Malaysia Research Fund, No. UKM-DLP-066-2011 and No. FF-032-2012.

**Institutional animal care and use committee statement:** All procedures involving animals were reviewed and approved by the Institutional Animal Care and Use Committee of Universiti Kebangsaan Malaysia (UKMAEC approval number: 354/2011).

**Conflict-of-interest statement:** The authors have no conflict of interests to declare.

**Data sharing statement:** No additional data is available.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Correspondence to:** Mohd Fahami Nur Azlina, DVM, MMed-SC, PhD, Associate Professor, Department of Pharmacology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur 56000, Malaysia. [nurazlinamf@ukm.edu.my](mailto:nurazlinamf@ukm.edu.my)  
Telephone: +603-91459574  
Fax: +603-91459547

Received: December 23, 2016

Peer-review started: December 28, 2016

First decision: April 21, 2017

Revised: May 8, 2017

Accepted: July 22, 2017

Article in press: July 24, 2017

Published online: August 28, 2017

## Abstract

### AIM

To investigate and compare the effects of tocotrienol and omeprazole on gastric growth factors in rats exposed to water-immersion restraint stress (WIRS).

### METHODS

Twenty-eight male Wistar rats were randomly assigned to four groups of seven rats. The two control groups were administered vitamin-free palm oil (vehicle) and the two treatment groups were given omeprazole (20 mg/kg) or tocotrienol (60 mg/kg) by oral gavage. After 28 d of treatment, rats from one control group and both treated groups were subjected to WIRS one time for 3.5 h. Gastric lesions were measured and gastric tissues were obtained to measure vascular endothelial growth factor (VEGF), epidermal growth factor (EGF), basic fibroblast growth factor (bFGF), and transforming growth factor- $\alpha$  (TGF- $\alpha$ ) mRNA expression.

### RESULTS

Rats exposed to WIRS for 3.5 h demonstrated the presence of considerable ulcers in the form of gastric erosion. The lesion index in the stressed control (S) group was increased ( $P < 0.001$ ) compared to the tocotrienol treated and omeprazole treated groups. Stress led to a decrease in gastric VEGF ( $P < 0.001$ ), bFGF ( $P < 0.001$ ) and TGF- $\alpha$  ( $P < 0.001$ ) mRNA levels and caused an increase in EGF mRNA ( $P < 0.001$ ) that was statistically significant compared to the non-stressed control group. Although both treatment agents

exerted similar ulcer reducing ability, only treatment with tocotrienol led to increased expression of VEGF ( $P = 0.008$ ), bFGF ( $P = 0.001$ ) and TGF- $\alpha$  ( $P = 0.002$ ) mRNA.

### CONCLUSION

Tocotrienol provides gastroprotective effects in WIRS-induced ulcers. Compared to omeprazole, tocotrienol exerts a similar protective effect, albeit through multiple mechanisms of protection, particularly through up-regulation of growth factors that assist in repair of gastric tissue injuries.

**Key words:** Tocotrienol; Omeprazole; Restraint-stress; Gastric ulcers; Growth factors

© The Author(s) 2017. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** During the process of ulcer healing, growth factors such as epidermal growth factor, transforming growth factor- $\alpha$ , basic fibroblast growth factor and vascular endothelial growth factor acts by activating the migration of cells from the edge of the ulcer and cell proliferation together with the formation of granulation tissue and angiogenesis. Rats exposed to stress develop gastric mucosal ulcers and changes in expression of these growth factors surrounding the ulcers had been reported. Tocotrienol effects on gastric mucosal growth factors were compared to those by omeprazole in this study. The findings suggest that in contrast with omeprazole, tocotrienol has a protective effect on the gastric mucosa through its effect on these growth factors.

Nur Azlina MF, Qodriyah HMS, Chua KH, Kamisah Y. Comparison between tocotrienol and omeprazole on gastric growth factors in stress-exposed rats. *World J Gastroenterol* 2017; 23(32): 5887-5894 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i32/5887.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i32.5887>

### INTRODUCTION

Stress ulcers often occur in critically ill patients as a result of major stressful events, such as trauma, shock, surgery, sepsis and burns. The responses to stress are both psychological and physiological. Physiological responses include neurohormonal and immunological activation, which includes release of corticotropin-releasing factor<sup>[1]</sup>, while the psychological responses include anxiety, depression, feeling of helplessness, fear, *etc.*

The pathological basis for the development of stress ulcers is multifactorial and includes changes in gastric acid secretion<sup>[2]</sup>, oxidative stress<sup>[3-5]</sup>, impaired gastric blood flow<sup>[6]</sup>, reduced gastric prostaglandin synthesis<sup>[7,8]</sup>, inflammation<sup>[9,10]</sup>, and inhibition of

mucosal growth and proliferation. Growth factors, by contrast, play pivotal roles in prevention and repair of stress-induced gastric ulcers<sup>[11]</sup>. This is particularly true during recovery of the mucosa after stress-induced injuries. Vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), transforming growth factor- $\alpha$  (TGF- $\alpha$ ) and epidermal growth factor (EGF) are crucial for reconstruction of damaged mucosal structures.

Among the growth factors involved, VEGF and bFGF are important factors because of their effects on angiogenesis. These growth factors are produced by endothelial cells, fibroblasts, macrophages and smooth muscle cells, and are involved in the regulation of physiological and pathological angiogenesis<sup>[12]</sup>. Angiogenesis and growth factors such as bFGF and VEGF play important roles in the repair of gastric ulcers caused by disturbances in the balance between factors that damage and factors that protect the stomach<sup>[13]</sup>.

In animal studies, angiogenesis has been shown to play a role in the process of supplying oxygen and nutrients to ulcers of affected areas<sup>[14]</sup>. Malara *et al.*<sup>[15]</sup> demonstrated a role for VEGF and angiogenesis in the repair of gastric ulcers caused by stress induced in rats. Research has also shown that a significant increase in the expression of VEGF protein followed by the formation of new blood vessels occurred as early as 1 d after formation of the ulcers<sup>[14]</sup>. Exogenous bFGF was found to assist in repair of gastric ulcers and other stress, and angiogenesis was also found to reduce gastric acid secretion<sup>[13]</sup>. Ernst *et al.*<sup>[13]</sup> reported that there was a reduction in stress-induced endogenous bFGF and that this resulted in reduced gastric micro-circulation, which plays an important role in the repair of ulcers. It is unknown if transcription of these growth factors are activated in response to stress-induced gastric injury.

Tocotrienol has been shown to prevent gastric ulcer development in rats exposed to noxious stimuli including ethanol, non-steroidal anti-inflammatory drugs (NSAIDs) and stress<sup>[8,16]</sup>. Tocotrienol, in comparison to tocopherol, was reported to be a more potent antioxidant<sup>[17]</sup>. Other than its antioxidant capabilities, tocotrienol has also been shown to have anti-inflammatory effects<sup>[10]</sup>, which may play an effective role in reducing damage to the gastric mucosa due to stress.

The gastric ulcer formation is complex in nature and involves multiple pathways that play a role in the prevention and repair<sup>[18]</sup>. The ulcer model that was used in this study was water-immersion restraint stress (WIRS). This experimental model was chosen because of its reproducibility, reliability and validity<sup>[15,19]</sup>. This model had been used to mimic clinical acute gastric ulcers formation in critically ill patients and is widely accepted for research involving the mechanism of stress-induced gastric ulcers<sup>[10,12]</sup>. The present study evaluated the limited information on the gastro-protective activity of palm-derived tocotrienol with

relation to anti-ulcer properties and gastric growth factors. The purpose of the study is to contribute a better understanding of the pathophysiology of stress-induced gastric ulcers. In this study, tocotrienol was compared to omeprazole, one of the widely used drugs for peptic-ulcer disease in clinical settings.

## MATERIALS AND METHODS

Male Wistar rats ( $n = 28$ ) were divided into four equally sized groups. Two control groups were fed a normal rat diet (non-stressed: NS and stressed: S), while the treatment groups received the same diet but with oral supplement of tocotrienol or omeprazole at 60 mg/kg or 20 mg/kg body weight, respectively, for 28 d. The tocotrienol dose was chosen based on our previous studies that demonstrated a protective effect on stress-induced gastric lesions<sup>[10,20]</sup>. The tocotrienol and omeprazole were diluted in vitamin-free palm oil, acting as vehicle, and administered by oral gavage using an 18G gavage needle. Both S and NS control groups were administered vitamin-free palm oil.

At the end of the treatment period, rats from the S control group and both treated groups were exposed to WIRS, by placing them in individual plastic restrainers measuring approximately 17 cm × 5 cm and immersing them in water neck deep one time for 3.5 h, as previously described by Aziz Ibrahim *et al.*<sup>[16]</sup>. Following the restraining procedure, rats were sacrificed by exsanguination under anesthesia. Stomachs were then dissected along the greater curvature. The dissected stomachs were taken for evaluation of gastric ulcers and mRNA expression of gastric EGF, bFGF, VEGF, and TGF- $\alpha$ . Gastric tissues were homogenized using an Omni Bead Ruptor machine at 25 °C with the speed of 8 m/s for 20 s. Homogenates were centrifuged at 1500 × *g* for 5 min at 4 °C. Supernatants were then used for mRNA analysis.

All rats were kept on a regular night/day cycle, with natural light for a period of 12 h (0700 to 1900 h). Throughout the feeding period, all rats were habituated to handling to reduce stress-related disturbances. Rats were housed in large cages with wide wire-mesh floors to prevent coprophagy. Food and water were given *ad libitum* throughout the experiment. Ethical approval was obtained from Universiti Kebangsaan Malaysia Animal Ethics Committee (UKMAEC). Humane methods of euthanasia were practiced (*i.e.*, exsanguination under anesthesia) by following guidelines of and with approval from UKMAEC. The anesthetic agent used was a combination of ketamine and xylazine (1:1 ratio).

### Assessment of stress-induced gastric lesions

The macroscopic assessment of stress-induced gastric lesions in the gastric mucosa was performed by two independent examiners who were blinded to the treatment. The assessment of lesions was done according to a semi-quantitative scale. Lesion size

(mm) was determined by measuring each lesion area. Five petechial lesions were equal to 1 mm lesion. The total lesion area in each group of rats were averaged and expressed as the lesion index; this method was modified as previously described by Aziz Ibrahim *et al.*<sup>[16]</sup>.

### Analysis of EGF, bFGF, VEGF and TGF- $\alpha$ mRNA expression

mRNA levels of EGF, bFGF, VEGF and TGF- $\alpha$  from gastric tissues were assayed according to the manufacturer's instructions using the standard QuantiGene Plex 2.0 assay kit (Genospectra, Fremont, CA, United States). Briefly, tissue lysates were transferred to a capture well in the presence of the gene-specific probe set and then hybridized at 53 °C overnight. Wells were washed twice with bDNA wash buffer and then incubated at 46 °C sequentially with an amplifier and an alkaline phosphatase-labeled probe, with a wash step in between incubations. After the final wash step, addition of streptavidin phycoerythrin generated a signal that was proportional to the amount of target RNA present in the sample. The luminescence signal was detected using a Luminex instrument. The protocol was followed as previously described by Zhang *et al.*<sup>[21]</sup>.

### Statistical analysis

Statistical analyses were performed using PRISM software version 6.00 (Graphpad, San Diego, CA, United States). The results are expressed as the mean ± SE of the mean. Statistical significance ( $P < 0.05$ ) was determined by ANOVA and Tukey's post-hoc test.

## RESULTS

### Macroscopic observation

Exposure to WIRS for 3.5 h caused the formation of ulcers in the form of gastric mucosal erosion and ulcers which were confined to the corpus of the stomach. The gastric lesion index (area in mm<sup>2</sup>) in the stressed-exposed (S) group was increased ( $11.92 \pm 2.0$  mm<sup>2</sup>;  $P = 0.001$ ) compared to the tocotrienol-treated group ( $0.94 \pm 0.30$  mm<sup>2</sup>) and the omeprazole-treated group ( $2.44 \pm 0.7$  mm<sup>2</sup>), as shown in Figure 1A. Rats not exposed to stress did not develop any gastric lesions.


### Quantitative changes in gastric VEGF mRNA expression in response to stress

Figure 1B shows that VEGF mRNA expression in stressed control rats was decreased by 45% compared to NS rats ( $P < 0.0001$ ). Pre-treatment with tocotrienol caused a statistically significant increase in VEGF expression compared to the stressed control group ( $P = 0.0075$ ). However, pre-treatment with omeprazole did not enhance VEGF expression compared to the S control group ( $P = 0.0593$ ).

### Quantitative changes in gastric EGF mRNA expression in response to stress

Stress exposure caused an increase in EGF gene





**Figure 1** Effects of pre-treatment with tocotrienol (60 mg/kg body weight) or omeprazole (20 mg/kg body weight) on gastric lesions formation (A), and gastric vascular endothelial growth factor (B), epidermal growth factor (C), transforming growth factor- $\alpha$  (D) and basic fibroblast growth factor (E) mRNA expression in rats exposed to water immersion restraint stress ( $n = 7$ ). <sup>a</sup> $P < 0.05$  vs NS non-stressed control (NS); <sup>c</sup> $P < 0.05$  vs Stressed control (S); <sup>e</sup> $P < 0.05$  vs Omeprazole (OMZ + S). bFGF: Basic fibroblast growth factor; BW: Body weight; EGF: Epidermal growth factor; NS: Non-stressed; OMZ: Omeprazole; S: Stressed; TGF- $\alpha$ : Transforming growth factor-alpha; TT: Tocotrienol; VEGF: Vascular endothelial growth factor.

expression ( $P = 0.0001$ ). Expression was increased in all groups exposed to stress and no differences were observed between the three stressed groups (S, tocotrienol-treated and omeprazole-treated) (Figure 1C), suggesting that tocotrienol and omeprazole had no effect on EGF mRNA levels in stress-exposed rats.

#### Quantitative changes in gastric TGF- $\alpha$ mRNA expression in response to stress

Exposure to immobilization stress caused a decrease in expression of TGF- $\alpha$  in gastric tissue (Figure 1D). Stress caused a significant reduction in the gastric expression as shown. Pre-treatment with tocotrienol caused an increase in TGF- $\alpha$  gene expression compared to the S control group ( $P = 0.0024$ ). Similar levels of TGF- $\alpha$  were observed in the NS control group. This result suggests that tocotrienol had a protective effect by preventing the stress-induced decrease of TGF. By contrast, TGF- $\alpha$  gene expression in the omeprazole-treated rats was decreased to levels similar to those observed in the S control group ( $P > 0.999$ ).

#### Quantitative changes in gastric bFGF mRNA expression in response to stress

Stress caused a 63% decrease in bFGF gene expression ( $P = 0.0022$ ), as compared to NS rats. bFGF expression in the tocotrienol-treated group was increased compared

to stressed control rats ( $P = 0.0013$ ) and there was no statistically significant change in bFGF levels when compared with the NS control rats. This suggests that pre-treatment with tocotrienol protects stress-induced rats by preventing a decrease in the level of bFGF to a level that was similar to that observed in the NS controls. bFGF expression in the tocotrienol-treated group was increased by 43% compared to the omeprazole-treated group ( $P = 0.0492$ ). Omeprazole treatment had no protective effect when the bFGF levels were similar to those observed in S rats, as shown in Figure 1E.

## DISCUSSION

Various genes are regulated at different rates and times during and after gastric mucosal injury. Most data regarding gene expression after mucosal injury models are taken from experimental animals. Together with the formation of granulation tissue, blood vessel and angiogenesis, growth factors act to enable the migration of cells from the edge of the ulcer and induce cell proliferation during the process of ulcer healing<sup>[7,22]</sup>. For example, EGF mRNA is detected immediately after ulcer induction, peaks during day 3, and continues to decrease 10 d after the induction of ulcers, whereas TGF- $\alpha$  mRNA expression increases 6 d after injury<sup>[23]</sup>.

VEGF is a growth factor that helps tissue healing

by stimulating angiogenesis, and is also important for the formation of connective tissue<sup>[24,25]</sup>. Antonisamy *et al.*<sup>[26]</sup> showed that injury to the gastric mucosa through administration of indomethacin caused a striking decrease in VEGF content in the gastric mucosa. Findings from this study also showed that exposure to WIRS caused a significant decrease in VEGF gene expression in gastric tissue compared to rats that were not exposed to stress. In another study, pre-treatment with a single dose of oral VEGF protected the stomach against damage due to acute ethanol administration<sup>[27]</sup>. Furthermore, gastric ulcer healing was prolonged and angiogenesis was decreased in response to a reduction in expression of VEGF<sup>[28]</sup>. Finally, up-regulation of VEGF has been shown to play an important role in the healing of acute gastric injury<sup>[29]</sup>.

Our findings show that tocotrienol led to increased VEGF expression in stress-induced rats. Studies that evaluate the effect of tocotrienols on growth factor expression in gastric ulcers are limited.  $\delta$ -tocotrienol has been shown to decrease the expression of VEGF in tumor cells, thereby reducing angiogenesis in these cells<sup>[30-32]</sup>. The tocotrienol administered in this study contained less than 4%  $\delta$ -tocotrienol and consisted mostly of other isomers, the most being isomeric  $\gamma$ -tocotrienol (approximately 50%). The effect obtained from this study leads us to assume that the other isomers in the tocotrienol mixture used in this study might assist in the healing process of ulcers or provide gastric mucosal protection against injury by promoting the process of granulation tissue formation and angiogenesis through VEGF expression.

Unlike tocotrienol, omeprazole administration did not decrease VEGF expression in stress-induced rats. This is inconsistent with results from a study by Kobayashi *et al.*<sup>[30]</sup> that showed that administration of lansoprazole (another proton pump inhibitor) led to increased expression of the VEGF gene in rats with gastric ulcers induced with acetic acid. These results suggest that proton pump inhibitors have an additional impact on protection of gastric ulcer formation (*i.e.*, regulation of growth factor expression) other than prevention of gastric acid secretion<sup>[30]</sup>. However, this effect was not observed in our study using omeprazole, suggesting that not all proton pump inhibitors have the same effect on growth factors. Abdul-Aziz *et al.*<sup>[27]</sup> reported similar results, *i.e.*, omeprazole reduced gastric ulcers but did not enhance VEGF expression levels.

In addition to VEGF, the growth of the mucosa is under the influence of various other growth factors, such as EGF and polyamines, which play important roles in tissue maintenance and repair. EGF is secreted into the intestinal lumen and into the bloodstream after being produced in saliva and pancreatic gland, and excreted *via* the urine as urogastrone<sup>[33]</sup>. It is well established that EGF is necessary for the maintenance of mucosal integrity. Furthermore, accumulation of EGF in the region of gastric mucosal injury promotes

the local lesion healing process<sup>[1]</sup>.

Milani *et al.*<sup>[24]</sup> showed that expression of growth factors detected in gastric mucosal cells, constantly fluctuates even under normal conditions. In the absence of induction of lesions, vitamin E does not significantly affect the function of growth factors where a decline in immunoreactivity of the EGF receptor (EGF-R) has been reported<sup>[34]</sup>. The results of this study also showed that rats that were not exposed to stress had low levels of EGF gene expression. Exposure to NSAIDs causes gastric ulcers to form and leads to decreased levels of EGF in gastric mucosa<sup>[26]</sup>. Increased expression of both EGF and EGF-R in the ulcer area also contributes to the repair process<sup>[35]</sup>. This reaction may occur in response to the sharp decline of this growth factor during gastric injury.

Exposure to stress led to formation of gastric lesions and caused an increase in gene expression of EGF that was statistically significant compared to the non-stressed control group. Increased expression of EGF has been reported to accelerate healing of gastroduodenal ulcers by increasing gastric mucin production and reducing gastric acid secretion<sup>[26]</sup>. This indicates that expression of EGF levels increase in response to injury of the gastric mucosa in order to restore the tissue back to its original state.

In this study, however, pre-treatment with tocotrienol or omeprazole did not change gastric tissue EGF expression; expression of EGF remained elevated in stress-exposed rats compared to rats that were not exposed to stress. This may have occurred due to the presence of gastric tissue injury in the treatment group that could have led to increased expression of EGF, which accelerates the recovery of gastric tissue caused by stress. Studies by Qodriyah *et al.*<sup>[34]</sup> showed that EGF levels were increased compared to the control group under normal conditions after 8 wk of palm vitamin E (PVE) treatment. The results of their study suggest that, under normal circumstances, vitamin E also enhances expression of EGF. While in a state of gastric injury due to NSAIDs, expression of EGF remained elevated after pre-treatment with PVE<sup>[34]</sup>, consistent with the results of this study.

EGF released from the salivary glands and TGF- $\alpha$  from the gastric mucosa act to maintain mucosal integrity and recovery during gastric mucosal injuries. Both these growth factors produce the same biological activities during recovery. For example, TGF- $\alpha$  and bFGF levels change when an injury occurs in the gastric mucosa. In this study, exposure to stress caused a significant decrease in bFGF gene expression. bFGFs have been shown to play a role in both angiogenesis and recovery of gastric ulcers in rats<sup>[36,37]</sup>. bFGF activation occurs in response to injury of the gastric mucosa, as demonstrated by the increased expression observed near ulcers<sup>[38]</sup>. Administration of bFGF (100 ng) locally into ulcers or systemically caused significant recovery in acetic acid-induced gastric ulcers<sup>[37]</sup>. bFGFs are also known to stimulate synthesis of prostaglandins

locally<sup>[30,39]</sup>, leading to increased formation of blood vessels<sup>[14,40]</sup> as well as proliferation of endothelial cells<sup>[41]</sup>, sustaining and assisting the recovery of gastric tissues in the event of injury.

Pre-treatment with tocotrienol in this study led to increased expression of bFGF in rats that were exposed to stress. Vitamin E at 150 mg/kg has been shown to improve bFGF expression in mice that developed gastric mucosal injury due to NSAID exposure<sup>[34,42]</sup>. Rashid *et al.*<sup>[42]</sup> showed that tocotrienol increased bFGF levels, thus reducing the formation of scar tissue. However, our study found that omeprazole had no effect on bFGF gene expression, which is in contrast to the tocotrienol-treated group. Studies that examine bFGF expression in response to omeprazole treatment are limited. Tsuji *et al.*<sup>[43]</sup> found that administration of lansoprazole, also a proton pump inhibitor, helped repair gastric ulcers by increasing bFGF levels at the edge of the ulcer border. Pantoprazole helped promote angiogenesis in gastric lesions induced by NSAIDs through increased expression of bFGF<sup>[44]</sup>. This effect, however, was not observed with pre-treatment of omeprazole in this study.

Growth factors such as EGF, TGF- $\alpha$ , bFGF and VEGF activate migration and proliferation of cells at the edge of the ulcer and promote the formation of granulation tissue and angiogenesis during the process of ulcer repair<sup>[22]</sup>. EGF is required to maintain the integrity of gastric mucosa and promotes healing of injured tissue<sup>[26]</sup>. TGF- $\alpha$  accelerates replacement of the epithelium and regulates regeneration of epithelial cells in gastric tissues<sup>[33]</sup>. VEGF assists in repair of ulcers by stimulating angiogenesis and remodeling of connective tissues<sup>[24]</sup>, while bFGF has been known to stimulate synthesis of local prostaglandins, which ultimately leads to increased formation of blood vessels<sup>[14,40]</sup> and endothelial cell proliferation<sup>[41]</sup>. This assists in maintenance and recovery of gastric tissue in the event of injury. The results of this study suggest that in contrast with omeprazole, tocotrienol has a protective effect on the gastric mucosa through regulation of these growth factors.

Here, we show that tocotrienol provides a gastro-protective effect in WIRS-induced ulcers and exerts similar effectiveness when compared to omeprazole. However, it displays a more diverse mechanism of protection, particularly through increased expression of bFGF, TGF- $\alpha$  and VEGF in a stress-induced gastric ulcer rat model in comparison to omeprazole. Thus, the effect of tocotrienol might be of therapeutic interest for the prevention and repair of gastric mucosal injuries due to other mechanisms.

## ACKNOWLEDGMENTS

We would like to acknowledge ExcelVite® Sdn. Bhd. for the supply of tocotrienol for this research. The authors also wish to thank En. Muhamad Arizi Aziz, Puan Juliana, Cik Hafizah, Puan Farhana and Puan Nurul

Akmal for their technical assistance.

## COMMENTS

### Background

Stress is well known to induce gastric ulcers. Although the proposed pathogenesis is multifactorial, a common entity observed in peptic ulcer diseases is oxidative stress which overwhelms the endogenous antioxidant system. Thus, prevention and treatment using an antioxidant like tocotrienol is a logical therapeutic approach. This study focuses on the therapeutic ability of tocotrienol on reducing stress-induced gastric ulcers and its effects on gastric growth factors which plays an important role in the prevention and repair of ulcers.

### Research frontiers

Few studies had investigated the effect of tocotrienol from palm source on gastric growth factors.

### Innovations and breakthrough

Tocotrienol provides gastroprotective effect in water-immersion restraint stress-induced ulcers. Although tocotrienol provides similar effectiveness as compared to omeprazole, it has a more diverse mechanism of protection, particularly through up-regulation of basic fibroblast growth factor, transforming growth factor- $\alpha$  and vascular endothelial growth factor in a stress-induced gastric ulcer model.

### Applications

Tocotrienol as therapeutic agent for the prevention and enhancing the repair of gastric mucosa against injuries.

### Terminology

Stress ulcers can occur as a result of major stressful events, such as trauma, shock, surgery, sepsis and burns. Tocotrienol prevents gastric ulcer development in rats exposed to noxious stimuli including ethanol, non-steroidal anti-inflammatory drugs and stress.

### Peer-review

The research is well conducted and the paper is well written. The series of experiments conducted were able to answer the objective of the study and the statistical tests used were scientifically sound.

## REFERENCES

- 1 Guo S, Gao Q, Jiao Q, Hao W, Gao X, Cao JM. Gastric mucosal damage in water immersion stress: mechanism and prevention with GHRP-6. *World J Gastroenterol* 2012; **18**: 3145-3155 [PMID: 22791951 DOI: 10.3748/wjg.v18.i24.3145]
- 2 Dalia M, Abd El Motteleb, Mai MH. Gastroprotective effect of simvastatin against experimentally induced gastric ulcers in rats: Role of ATP-sensitive K<sup>+</sup> channels. *J Am Sci* 2011; **7**: 760-768
- 3 Brzozowski T, Konturek PC, Sliwowski Z, Drozdowicz D, Burnat G, Pajdo R, Pawlik M, Bielanski W, Kato I, Kuwahara A, Konturek SJ, Pawlik WW. Gastroprotective action of orexin-A against stress-induced gastric damage is mediated by endogenous prostaglandins, sensory afferent neuropeptides and nitric oxide. *Regul Pept* 2008; **148**: 6-20 [PMID: 18378017 DOI: 10.1016/j.regpep.2008.02.003]
- 4 Nur Azlina MF, Rubaizah K, Muliana M, Nafeeza MI. Modulation of restraint stress induced gastric oxidative changes in rats by tocotrienol and tocopherol. *Int J Pharmacol* 2009; **5**: 58-64 [DOI: 10.3923/ijp.2009.58.64]
- 5 Mohd Fahami NA, Ibrahim IA, Kamisah Y, Mohd Ismail N. Palm vitamin E reduces catecholamines, xanthine oxidase activity and gastric lesions in rats exposed to water-immersion restraint stress. *BMC Gastroenterol* 2012; **12**: 54 [PMID: 22639913 DOI: 10.1186/1471-230X-12-54]
- 6 Wang YB, Liu J, Yang ZX. Effects of intestinal mucosal blood flow

- and motility on intestinal mucosa. *World J Gastroenterol* 2011; **17**: 657-661 [PMID: 21350716 DOI: 10.3748/wjg.v17.i5.657]
- 7 **Konturek PC**, Brzozowski T, Duda A, Kwiecien S, Löber S, Dembinski A, Hahn EG, Konturek SJ. Epidermal growth factor and prostaglandin E(2) accelerate mucosal recovery from stress-induced gastric lesions via inhibition of apoptosis. *J Physiol Paris* 2001; **95**: 361-367 [PMID: 11595461 DOI: 10.1016/S0928-4257(01)00049-3]
- 8 **Nur Azlina MF**, Kamisah Y, Chua KH, Qodriyah HM. Tocotrienol Attenuates Stress-Induced Gastric Lesions via Activation of Prostaglandin and Upregulation of COX-1 mRNA. *Evid Based Complement Alternat Med* 2013; **2013**: 804796 [PMID: 23970937 DOI: 10.1155/2013/804796]
- 9 **Bregonzio C**, Armando I, Ando H, Jezova M, Baiardi G, Saavedra JM. Anti-inflammatory effects of angiotensin II AT1 receptor antagonism prevent stress-induced gastric injury. *Am J Physiol Gastrointest Liver Physiol* 2003; **285**: G414-G423 [PMID: 12686508 DOI: 10.1152/ajpgi.00058.2003]
- 10 **Nur Azlina MF**, Kamisah Y, Chua KH, Ibrahim IA, Qodriyah HM. Preventive Effects of Tocotrienol on Stress-Induced Gastric Mucosal Lesions and Its Relation to Oxidative and Inflammatory Biomarkers. *PLoS One* 2015; **10**: e0139348 [PMID: 26465592 DOI: 10.1371/journal.pone.0139348]
- 11 **Konturek PC**, Brzozowski T, Konturek SJ, Taut A, Sliwowski Z, Stachura J, Hahn EG. Activation of genes for growth factors and cyclooxygenases in rat gastric mucosa during recovery from stress damage. *Eur J Pharmacol* 1998; **342**: 55-65 [PMID: 9544793]
- 12 **Plate KH**. Mechanisms of angiogenesis in the brain. *J Neuropathol Exp Neurol* 1999; **58**: 313-320 [PMID: 10218626 DOI: 10.1097/00005072-199904000-00001]
- 13 **Ernst M**, Inglese M, Waring P, Campbell IK, Bao S, Clay FJ, Alexander WS, Wicks IP, Tarlinton DM, Novak U, Heath JK, Dunn AR. Defective gp130-mediated signal transducer and activator of transcription (STAT) signaling results in degenerative joint disease, gastrointestinal ulceration, and failure of uterine implantation. *J Exp Med* 2001; **194**: 189-203 [PMID: 11457894]
- 14 **Schmassmann A**, Peskar BM, Stettler C, Netzer P, Stroff T, Flogerzi B, Halter F. Effects of inhibition of prostaglandin endoperoxide synthase-2 in chronic gastro-intestinal ulcer models in rats. *Br J Pharmacol* 1998; **123**: 795-804 [PMID: 9535006 DOI: 10.1038/sj.bjp.0701672]
- 15 **Malara B**, Joško J, Tyrciński M, Malara P, Steplewska K. Dynamics of changes in vascular endothelial growth factor (VEGF) expression and angiogenesis in stress-induced gastric ulceration in rats. *J Physiol Pharmacol* 2005; **56**: 259-271 [PMID: 15985707]
- 16 **Aziz Ibrahim IA**, Kamisah Y, Nafeeza MI, Nur Azlina MF. The effects of palm vitamin E on stress hormone levels and gastric lesions in stress-induced rats. *Arch Med Sci* 2012; **8**: 22-29 [PMID: 22457670 DOI: 10.5114/aoms.2012.27276]
- 17 **Serbinova EA**, Packer L. Antioxidant properties of alpha-tocopherol and alpha-tocotrienol. *Methods Enzymol* 1994; **234**: 354-366 [PMID: 7808307 DOI: 10.1016/0076-6879(94)34105-2]
- 18 **El-Moselhy MA**, Abdel-Hamid NM, Abdel-Raheim SR. Gastroprotective effect of nicorandil in indomethacin and alcohol-induced acute ulcers. *Appl Biochem Biotechnol* 2009; **152**: 449-459 [PMID: 18931948 DOI: 10.1007/s12010-008-8384-z]
- 19 **Konturek SJ**, Brzozowski T, Konturek PC, Zwirska-Korczala K, Reiter RJ. Day/night differences in stress-induced gastric lesions in rats with an intact pineal gland or after pinealectomy. *J Pineal Res* 2008; **44**: 408-415 [PMID: 18086146 DOI: 10.1111/j.1600-079X.2007.00543.x]
- 20 **Kamisah Y**, Ibrahim AAI, Nafeeza MI, Nur Azlina MF. Palm tocotrienol rich fraction supplementation suppressed stress-induced gastric oxidative stress in rats. *J Appl Phar Sci* 2011; **1**: 118-122
- 21 **Zhang A**, Pastor L, Nguyen Q, Luo Y, Yang W, Flagella M, Chavli R, Bui S, Nguyen CT, Zheng Z, He W, McMaster G, Witney F. Small interfering RNA and gene expression analysis using a multiplex branched DNA assay without RNA purification. *J Biomol Screen* 2005; **10**: 549-556 [PMID: 16103413 DOI: 10.1177/1087057105277414]
- 22 **Konturek PC**, Konturek SJ, Brzozowski T, Ernst H. Epidermal growth factor and transforming growth factor-alpha: role in protection and healing of gastric mucosal lesions. *Eur J Gastroenterol Hepatol* 1995; **7**: 933-937 [PMID: 8590137]
- 23 **Pohle T**, Shahin M, Domschke W, Konturek JW. Effect of basic fibroblast growth factor on gastric ulcer healing and its own mRNA expression. *Aliment Pharmacol Ther* 1999; **13**: 1543-1551 [PMID: 10571614 DOI: 10.1046/j.1365-2036.1999.00627x]
- 24 **Milani S**, Calabrò A. Role of growth factors and their receptors in gastric ulcer healing. *Microsc Res Tech* 2001; **53**: 360-371 [PMID: 11376497 DOI: 10.1002/jemt.1104]
- 25 **Szabo S**, Vincze A. Growth factors in ulcer healing: lessons from recent studies. *J Physiol Paris* 2000; **94**: 77-81 [PMID: 10791686 DOI: 10.1016/S0928-4257(00)00146-7]
- 26 **Antonisamy P**, Kannan P, Aravinthan A, Duraipandiyan V, Arasu MV, Ignacimuthu S, Al-Dhabi NA, Kim JH. Gastroprotective activity of violacein isolated from *Chromobacterium violaceum* on indomethacin-induced gastric lesions in rats: investigation of potential mechanisms of action. *ScientificWorldJournal* 2014; **2014**: 616432 [PMID: 25162059 DOI: 10.1155/2014/616432]
- 27 **Abdul-Aziz MA**. Anti-ulcerogenic activity of peel an extracts of *Musa acuminata* against ethanol-induced ulcer in vivo. Bachelor's thesis, Universiti Selangor, Bestari Jaya Campus, Kuala Selangor, Selangor, Malaysia. 2011
- 28 **Harsch IA**, Brzozowski T, Bazela K, Konturek SJ, Kukharsky V, Pawlik T, Pawlowski E, Hahn EG, Konturek PC. Impaired gastric ulcer healing in diabetic rats: role of heat shock protein, growth factors, prostaglandins and proinflammatory cytokines. *Eur J Pharmacol* 2003; **481**: 249-260 [PMID: 14642793 DOI: 10.1016/j.ejphar.2003.09.019]
- 29 **Jones MK**, Kawanaka H, Baatar D, Szabó IL, Tsugawa K, Pai R, Koh GY, Kim I, Sarfeh II, Tarnawski AS. Gene therapy for gastric ulcers with single local injection of naked DNA encoding VEGF and angiopoietin-1. *Gastroenterology* 2001; **121**: 1040-1047 [PMID: 11677194 DOI: 10.1053/gast.2001.29308]
- 30 **Kobayashi S**, Nakajima N, Ito Y, Moriyama M. Effects of lansoprazole on the expression of VEGF and cellular proliferation in a rat model of acetic acid-induced gastric ulcer. *J Gastroenterol* 2010; **45**: 846-858 [PMID: 20333532 DOI: 10.1007/s00535-010-0224-6]
- 31 **Weng-Yew W**, Selvaduray KR, Ming CH, Nesaretnam K. Suppression of tumor growth by palm tocotrienols via the attenuation of angiogenesis. *Nutr Cancer* 2009; **61**: 367-373 [PMID: 19373610 DOI: 10.1080/01635580802582736]
- 32 **Selvaduray KR**, Radhakrishnan AK, Kutty MK, Nesaretnam K. Palm tocotrienols decrease levels of pro-angiogenic markers in human umbilical vein endothelial cells (HUVEC) and murine mammary cancer cells. *Genes Nutr* 2012; **7**: 53-61 [PMID: 21526401 DOI: 10.1007/s12263-011-0223-0]
- 33 **Tétreault MP**, Chailier P, Beaulieu JF, Rivard N, Ménard D. Epidermal growth factor receptor-dependent PI3K-activation promotes restitution of wounded human gastric epithelial monolayers. *J Cell Physiol* 2008; **214**: 545-557 [PMID: 17708540 DOI: 10.1002/jcp.21239]
- 34 **Qodriyah HMS**, Kamsiah J, Kamisah Y, Gapor MT, Nafeeza MI. Kesan palm vitamin E ke atas gatser tikus aruhan dadah anti-inflamasi bukan steroid. Thesis Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia, 2006
- 35 **Al Mofleh IA**. Spices as Alternative Agents for Gastric Ulcer Prevention and Treatment, Peptic Ulcer Disease, Dr. Jianyuan Chai (Ed.), 2011 ISBN: 978-953-307-976-9, InTech, Available from: URL: <http://www.intechopen.com/books/peptic-ulcer-disease/spices-as-alternative-agents-for-gastric-ulcerprevention-and-treatment>
- 36 **Hull MA**, Brough JL, Powe DG, Carter GI, Jenkins D, Hawkey CJ. Expression of basic fibroblast growth factor in intact and ulcerated human gastric mucosa. *Gut* 1998; **43**: 525-536 [PMID: 9824581 DOI: 10.1136/gut.43.4.525]
- 37 **Brzozowski T**, Konturek PC, Konturek SJ, Kwiecień S, Drozdowicz D, Bielanski W, Pajdo R, Ptak A, Nikiforuk A,



- Pawlik WW, Hahn EG. Exogenous and endogenous ghrelin in gastroprotection against stress-induced gastric damage. *Regul Pept* 2004; **120**: 39-51 [PMID: 15177919 DOI: 10.1016/j.regpep.2004.02.010]
- 38 **Tarnawski AS**. Cellular and molecular mechanisms of gastrointestinal ulcer healing. *Dig Dis Sci* 2005; **50** Suppl 1: S24-S33 [PMID: 16184417 DOI: 10.1007/s10620-005-2803-6]
- 39 **Kawaguchi H**, Pilbeam CC, Gronowicz G, Abreu C, Fletcher BS, Herschman HR, Raisz LG, Hurley MM. Transcriptional induction of prostaglandin G/H synthase-2 by basic fibroblast growth factor. *J Clin Invest* 1995; **96**: 923-930 [PMID: 7635987 DOI: 10.1172/JCI118140]
- 40 **Wang CL**, Shi DZ, Yin HJ. [Effect of panax quinquefolius saponin on angiogenesis and expressions of VEGF and bFGF in myocardium of rats with acute myocardial infarction]. *Zhongguo Zhongxiyi Jiehe Zazhi* 2007; **27**: 331-334 [PMID: 17526173]
- 41 **Hull MA**, Hewett PW, Brough JL, Hawkey CJ. Isolation and culture of human gastric endothelial cells. *Gastroenterology* 1996; **111**: 1230-1240 [PMID: 8898637 DOI: 10.1053/gast.1996.v111.pm8898637]
- 42 **Rashid SA**, Halim AS, Muhammad NA. The effect of vitamin E on basic fibroblast growth factor level in human fibroblast cell culture. *Med J Malaysia* 2008; **63** Suppl A: 69-70 [PMID: 19024988]
- 43 **Tsuji S**, Kawano S, Higashi T, Mukuda T, Imaizumi T, Tatsumi T, Miura N, Miyajima K, Fukuda M, Noguchi M. Gastric ulcer healing and basic fibroblast growth factor: effects of lansoprazole and famotidine. *J Clin Gastroenterol* 1995; **20** Suppl 2: S1-S4 [PMID: 7594320]
- 44 **Lee JS**, Lee GH, Hyun KY. Effects of GUF (*Glycyrrhiza uralensis* Fischer) extract in water-immersion restraint stress (wirs)-induced gastric injury in rats. *Adv Sci Technol Lett* 2015; **88**: 247-250 [DOI: 10.14257/astl.2015.88.52]

**P- Reviewer:** Yuen KH    **S- Editor:** Gong ZM    **L- Editor:** Filipodia  
**E- Editor:** Xu XR





Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>



ISSN 1007-9327

