

## Match Overview

1	<b>Crossref</b> 20 words Eiichi Ogawa. "Transient elastography for patients with chronic hepatitis B and C virus infection: Non-invasive, quan ...	1%
2	<b>Crossref</b> 13 words Seung Up Kim. "The usefulness of liver stiffness measurement using FibroScan® in chronic hepatitis C in South Kor	<1%
3	<b>Internet</b> 13 words crawled on 03-Apr-2014 <a href="http://cjasn.asnjournals.org">cjasn.asnjournals.org</a>	<1%
4	<b>Internet</b> 13 words <a href="http://www.sriconference.com">www.sriconference.com</a>	<1%

Name of journal: *World Journal of Hepatology*

ESPS Manuscript NO: 32203

Manuscript Type: Original Article

*Retrospective Study*

Use of Aspartate aminotransferase to platelet ratio to reduce the need for FibroScan in the evaluation of liver fibrosis

Stephanie Wong, Dep Huynh, Frank Zhang, Nam Q Nguyen

### Abstract

#### AIM

To evaluate the performance of Aspartate aminotransferase to platelet ratio (APRI) score against FibroScan in predicting the presence of fibrosis.

Using APRI to reduce the need for FibroScan in liver fibrosis evaluation

全部 图片 视频 新闻 购物 地图 图书

找到约 64,500 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

### [Non-invasive assessment of liver fibrosis](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3959378/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3959378/>

In parallel with this, noninvasive assessment of liver fibrosis has experienced ... Thus, liver biopsy can be poorly tolerated by patients, especially if it needs to be repeated. ... Overview of non-invasive methods for the evaluation of liver fibrosis Q1. ... APRI and Fibrotest are the most extensively studied serum markers.

### [Non-invasive diagnosis of liver fibrosis and cirrhosis - NCBI](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4631961/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4631961/>

7 Nov 2015 ... The evaluation and follow up of liver fibrosis and cirrhosis have been .... It was also comparable to APRI, with AUROCs around 0.8[35], and of 0.73 in a recent ... Reduction in the Fibrotest score was also observed after treatment of ..... Among SWE methods, TE (Fibroscan) is the only non-imaging method, ...

### [Non-invasive assessment of liver fibrosis: Between prediction ... - NCBI](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4722001/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4722001/>

28 Jan 2016 ... Over the past decade, various non-invasive methods have been proposed to detect ... or excluding liver fibrosis, thus reducing the need for liver biopsy. ... The predicted liver fibrosis score was more strongly correlated with fibrosis ... by FibroScan, and it was found superior to liver stiffness alone, APRI, the ...

### [Core Concepts - Evaluation and Staging of Liver Fibrosis ...](http://www.hepatitisc.uw.edu/go/evaluation.../evaluation.../core.../all)

[www.hepatitisc.uw.edu/go/evaluation.../evaluation.../core.../all](http://www.hepatitisc.uw.edu/go/evaluation.../evaluation.../core.../all)

21 Oct 2015 ... Pathogenesis of Fibrosis with Chronic Hepatitis C: Hepatic fibrosis is a ... index ( APRI), FibroSure, and transient elastography (FibroScan), liver biopsy was ... In addition, some studies have evaluated other serum fibrotic markers, ... Hepatic Ultrasound: Hepatic ultrasound is a non-invasive, lower cost, and ...

[PDF] [EASL-ALEH Clinical Practice Guidelines: Non-invasive tests for ...](#)



Use of Aspartate aminotransferase to platelet ratio to reduce the need for FibroS

全部 图片 视频 新闻 地图 图书

找到约 28,300 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

### [Non-invasive assessment of liver fibrosis - NCBI - NIH](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3959378/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3959378/>

In parallel with this, noninvasive assessment of **liver fibrosis** has experienced .... **Aspartate aminotransferase** (AST) to **alanine aminotransferase** (ALT) **ratio** index .... The Fibroindex<sup>®</sup> was developed for patients with CHC and **uses platelet** ..... concluding that this approach may **reduce the need** for **liver biopsy** by 30% [182].

### [Non-invasive assessment of liver fibrosis in chronic hepatitis C](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3090550/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3090550/>

20 Jan 2011 ... In patients with chronic hepatitis C, precise staging of liver fibrosis is ... Proposed serum scores for non-invasive **evaluation of liver fibrosis** in ... To date, FibroTest<sup>®</sup> and **aspartate** to **platelet ratio** index (APRI) **have** been ... One of the **main limitations** to the clinical **use** of "direct" markers of liver fibrosis is that ...

### [Evaluation of Aspartate Aminotransferase-to-Platelet Ratio Index as ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4668451/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4668451/>

1 Nov 2015 ... Cirrhosis is the end-stage consequence of **fibrosis** of the **liver** parenchyma ... infective peritonitis which limit the **use** of **liver** biopsies in all patients [2]. ... **Aspartate aminotransferase: Platelet Ratio** Index (APRI) was .... However, the purpose of this study was to **avoid the need** of biopsy in predicting cirrhosis.

### [\[PDF\] SAFE biopsy: A validated method for largescale staging of liver ...](https://onlinelibrary.wiley.com/doi/10.1002/hep.22859/pdf)

[onlinelibrary.wiley.com/doi/10.1002/hep.22859/pdf](https://onlinelibrary.wiley.com/doi/10.1002/hep.22859/pdf) ▼

described sequential algorithm for **fibrosis evaluation** (SAFE) biopsy, which detects signifi- ... rrosis simultaneously with high accuracy and a 36% **reduction** in the **need** for **liver** biopsy. ... Abbreviations: APRI, AST-to-**platelet ratio** index; AUROC, area under the receiver ... quential **use** of APRI, Fibrotest-Fibrosure, and **liver**.

### [\[PDF\] Diagnostic accuracy of the aspartate aminotransferase - to - platelet ...](#)

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 31,000 条结果 (用时 0.80 秒)

### Non-invasive assessment of liver fibrosis

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: V Papastergiou - 2012 - 被引用次数: 52 - 相关文章

In parallel with this, noninvasive assessment of liver fibrosis has experienced .... Aspartate aminotransferase (AST) to alanine aminotransferase (ALT) ratio index .... The Fibroindex<sup>®</sup> was developed for patients with CHC and uses platelet ..... concluding that this approach may reduce the need for liver biopsy by 30% [182].

### Non-invasive assessment of liver fibrosis in chronic hepatitis C

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: L Castera - 2011 - 被引用次数: 59 - 相关文章

2011年1月20日 - In patients with chronic hepatitis C, precise staging of liver fibrosis is ... Proposed serum scores for non-invasive evaluation of liver fibrosis in ... To date, FibroTest<sup>®</sup> and aspartate to platelet ratio index (APRI) have been ... One of the main limitations to the clinical use of "direct" markers of liver fibrosis is that...

### Evaluation of Aspartate Aminotransferase-to-Platelet Ratio Index as a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4668451/> - 翻译此页

作者: P Jain - 2015 - 相关文章

2015年11月1日 - Cirrhosis is the end-stage consequence of fibrosis of the liver parenchyma ... infective peritonitis which limit the use of liver biopsies in all patients [2]. .... Aspartate aminotransferase: Platelet Ratio Index (APRI) was .... However, the purpose of this study was to avoid the need of biopsy in predicting cirrhosis.

### Non-invasive diagnosis of liver fibrosis in chronic hepatitis C

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: L de Lencastre Schiavon - 2014 - 被引用次数: 24 - 相关文章