Q

学术搜索

Glgoo

获得 9 条结果 (用时0.07秒)

文章

我的图书馆

时间不限

2017以来 2016以来 2013以来

自定义范围....

按相关性排序

按日期排序

搜索所有网页

中文网页 简体中文网页

✔ 包括专利

☑ 包含引用

≥ 创建快讯

小提示: 只搜索中文(简体)结果,可在学术搜索设置。指定搜索语言

Transforming growth factor β in tissue fibrosis

WA Border, NA Noble - New England Journal of Medicine, 1994 - Mass Medical Soc ... in endothelial cells), and tumor necrosis factor and interleukin-1 (in monocytes) by stimulating ... In arthritic joints, anti-TGF-β1 decreased inflammation and bone and synovial destruction. ... Transforming growth factor type β induces monocyte chemotaxis and growth factor production ... 被引用次数: 3344 相关文章 所有 8 个版本 引用 保存

Experimental models of liver fibrosis

SC Yanguas, <u>B Cogliati</u>, J Willebrords, M Maes... - Archives of ..., 2016 - Springer ... 12(11):610–621PubMedCrossRef. French SW (2001) Intragastric ethanol infusion model for ... PubMedCentralCrossRef. Kisseleva T, Uchinami H, Feirt N et al (2006) Bone marrow-derived fibrocytes participate ... KS, Buck M, Houglum K et al (1995) Activation of hepatic stellate cells ... 被引用次数: 4 相关文章 所有 7 个版本 引用 保存

Pathophysiology of progressive renal disease

A Eddy - Pediatric Nephrology, 2009 - Springer

... Circulating monocytes that enter the kidney in response to inflammatory cues undergo distinctive ... through the use of hepatocyte growth factor (HGF) or bone morphogenic protein-7 ... as transferrin can stimulate expression of chemokines such a monocyte chemoattractant protein-1 ... 被引用次数: 2 相关文章 所有 2 个版本 引用 保存

Immune and inflammatory role in renal disease

JD Imig, MJ Ryan - Comprehensive Physiology, 2013 - Wiley Online Library

... by studies showing that macrophage depletion (97) or inhibition of monocyte chemoattractant protein ... Improved renal hemodynamics has been demonstrated to correlate with decreased levels of MCP-1 ... Monocytes that express MCP-1 and CCR2 infiltrate damaged tissue (88, 98 ... 被引用次数: 29 相关文章 所有 6 个版本 引用 保存

Compositions and Methods for Preventing or Treating Diseases, Conditions, or Processes Characterized by Aberrant Fibroblast Proliferation and Extracellular Matrix

C Lander, C Brophy - US Patent 20,160,136,234, 2016 - freepatentsonline.com

[HTML] nih.gov

[HTML] nih.gov

32302-Review
BY VERUSKA CINTIA ALEXANDRINO DE SOUZA

Quotes Excluded Bibliography Excluded

3% SIMILAR

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 32302

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Bone marrow-derived monocyte infusion improves hepatic fibrosis by decreasing osteopontin, TGF- β 1, IL-13 and oxidative stress

Souza VCA et al. Monocyte therapy improves hepatic fibrosis

Veruska Cintia Alexandrino de Souza, Thiago Almeida Pereira, Valéria Wanderley Teixeira, Helotonio Carvalho, Maria Carolina Accioly Brelaz de Castro, Carolline Guimarães D'assunção, Andréia Ferreira de Barros, Camila Lima Carvalho, Virgínia Maria Barros de Lorena, Vláudia Maria Assis Costa, Álvaro Aguiar Coelho Teixeira, Regina Celia Bressan Queiroz Figueiredo, Sheilla Andrade de Oliveira.

Veruska Cintia Alexandrino de Souza, Camila Lima Carvalho, Virgínia Maria Barros de Lorena, Regina Celia Bressan Oueiroz Figueiredo and Sheilla Andrade





Bone marrow-derived monocyte infusion improves hepatic fibrosis by decrea

Q

学术搜索

找到约 33 条结果 (用时0.09秒)

文章

我的图书馆

时间不限

2017以来 2016以来 2013以来 自定义范围...

按相关性排序

按日期排序

不限语言

中文网页 简体中文网页

▼ 包括专利

✔ 包含引用

☑ 创建快讯

小提示: 只搜索中文(简体)结果,可在学术搜索设置。指定搜索语言

The pathogenesis of cardiac fibrosis

P Kong, P Christia, NG Frangogiannis - Cellular and Molecular Life ..., 2014 - Springer ... A growing body of evidence suggests that bone marrow-derived circulating fibrocytes may represent an ... Studies using bone marrow transplantation strategies to generate chimeric mice have ... to the marked upregulation of the CC chemokine monocyte chemoattractant protein ... 被引用次数: 173 相关文章 所有 13 个版本 引用 保存

Mesenchymal stem cell therapy in lung disorders: pathogenesis of lung diseases and mechanism of action of mesenchymal stem cell

AC Inamdar, AA Inamdar - Experimental lung research, 2013 - Taylor & Francis ... The intravenous infusion of MSCs, the common mode of delivery, is followed by ... Specifically, MSCs based cell therapy may hold the potential to improve the quality of ... A2 (TXA2) synthase inhibitor, ozagrel, attenuates lung injury and decreases monocyte chemoattractant protein ... 被引用次数: 24 相关文章 所有 5 个版本 引用 保存

Transforming growth factor β in tissue fibrosis

WA Border, NA Noble - New England Journal of Medicine, 1994 - Mass Medical Soc ... in endothelial cells), and tumor necrosis factor and interleukin-1 (in monocytes) by stimulating ... In arthritic joints, anti-TGF-β1 decreased inflammation and bone and synovial destruction. ... Transforming growth factor type β induces monocyte chemotaxis and growth factor production ... 被引用次数: 3370 相关文章 所有 7 个版本 引用 保存

Cellular interplay between cardiomyocytes and nonmyocytes in cardiac remodeling

N Takeda, I Manabe - International journal of inflammation, 2011 - hindawi.com ... Finally, it has been suggested that monocytes/macrophages represent another potential source of ... In response to fibrogenic stimuli, however, many other cell types, including bone marrow-derived cells, pericytes ... In addition, infusion of synthetic ANP or BNP is useful for treating ... 被引用次数: 45 相关文章 所有 14 个版本 引用 保存 更多

Pathophysiology of progressive renal disease

[PDF] researchgate.net

[HTML] nih.gov

[PDF] hindawi.com



Bone marrow-derived monocyte infusion improves hepatic fibrosis by decreasir



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 42,300 条结果 (用时 0.94 秒)

Google 学术: Bone marrow-derived monocyte infusion improves hepatic fibrosis by decreasing osteopontin, TGF-β1, IL-13 and oxidative stress

The pathogenesis of cardiac fibrosis - Kong - 被引用次数: 190

Mesenchymal stem cell therapy in lung disorders: ... - Inamdar - 被引用次数: 25

Transforming growth factor β in tissue fibrosis - Border - 被引用次数:3397

Liver fibrosis - NCBI - NIH

https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: R Bataller - 2005 - 被引用次数: 3607 - 相关文章

These cells are activated by fibrogenic cytokines such as TGF- β 1, ... The main causes of liver fibrosis in industrialized countries include chronic HCV ... Besides HSCs, portal myofibroblasts and cells of bone marrow origin have been injury (e.g., IL-1 β , IL-6, IL-10, and IL-13, IFN- γ , SOCS-1, and osteopontin) determine the ...

Contribution of bone marrow-derived fibrocytes to liver fibrosis

https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页作者: J Xu - 2015 - 被引用次数: 10 - 相关文章

Contribution of bone marrow-derived fibrocytes to liver fibrosis ... like transforming growth factor β 1 (TGF- β 1), interleukin 6 (IL-6), interleukin 1 β (IL-1 β) and ... It is believed that decreased bile flow causes critical damage to the biliary ... that early activated PFs release factors (such as IL-13) that stimulate activation of HSCs in ...

Pathophysiology of NASH: perspectives for a targeted treatment

https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: F Marra - 2013 - 被引用次数: 59 - 相关文章

Within the liver, hepatocyte apoptosis, ER stress and oxidative stress are key such as