

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 3,780 条结果 (用时 0.95 秒)

## Hep C Articles... - NATAP

[www.natap.org/hcv.htm](http://www.natap.org/hcv.htm) ▼

AASLD: **Effectiveness** and **Safety** of DAA Combination **Therapies** for **Treatment** of HCV in ... AASLD: **Treatment** outcomes for **hepatitis C genotype 1** infection with direct acting .... AASLD: **Real World Effectiveness** of **Ledipasvir/Sofosbuvir** in **Patients** .... to **Sofosbuvir/Velpatasvir** for 12 **Weeks** in DAA-Naïve **Genotype 1-6 HCV** ...

## Role of ledipasvir/sofosbuvir combination for genotype 1 hepatitis C ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: V Sundaram - 2016 - 被引用次数: 1

2016年6月28日 - Among the **six** HCV genotypes, **genotype 1** was significantly more ... or ~2%–3% of the **world's** population, **hepatitis C** virus (HCV) is the ... The recommendation for **treatment-naïve patients** with **genotype 1** infection without **cirrhosis** is the ... **HCV RNA** level <**6 million IU/mL** who received **8 weeks** of therapy ...

## INITIAL TREATMENT OF HCV INFECTION | Recommendations for ...

[www.hcvguidelines.org/full-report/initial-treatment-hcv-infection](http://www.hcvguidelines.org/full-report/initial-treatment-hcv-infection) ▼ 翻译此页

Initial treatment of HCV infection includes **patients** with chronic **hepatitis C** who have **not** been ... **Genotype 1a Treatment-Naïve Patients** Without **Cirrhosis** - Recommended .... **8 weeks** of **ledipasvir/sofosbuvir** who had baseline **HCV RNA** levels below **6 million IU/mL** (2%; 2 of 123), and was the same for **patients** with similar ...

## Ledipasvir-Sofosbuvir (Harvoni) - Treatment - Hepatitis C Online

[www.hepatitisc.uw.edu/page/treatment/drugs/ledipasvir-sofosbuvir](http://www.hepatitisc.uw.edu/page/treatment/drugs/ledipasvir-sofosbuvir) ▼ 翻译此页



32607-Review-check.docx

Quotes Included  
Bibliography Included4%  
SIMILARName of Journal: *World Journal of Gastroenterology*

Manuscript NO: 32607

Manuscript Type: ORIGINAL ARTICLE

*Retrospective Study*

Eight-week ledipasvir/sofosbuvir in non-cirrhotic, treatment-naïve hepatitis C genotype-1 patients with HCV-RNA < 6 million: Single center, real world effectiveness and safety

Latt NL *et al.* 8-wk LDV/SOF therapy for non-cirrhotic, treatment-naïve hepatitis C genotype-1 patients

Nyan L Latt, Beshoy T Yanny, Derenik Gharibian, Rita Gevorkyan, Amandeep K Sahota

Abstract

AIM

## Match Overview

1	<b>Crossref</b> 17 words Eric Nguyen-Khac. "Noninvasive Diagnosis of Large Esophageal Varices by Fibroscan: Strong Influence of the Ci	<1%
2	<b>Internet</b> 17 words crawled on 06-Mar-2017 <a href="http://repub.eur.nl">repub.eur.nl</a>	<1%
3	<b>Internet</b> 16 words crawled on 16-Sep-2014 <a href="http://www.projectsinknowledge.com">www.projectsinknowledge.com</a>	<1%
4	<b>Crossref</b> 14 words Backus, Lisa I., Pamela S. Belperio, Troy A. Shahoumian, Timothy P. Loomis, and Larry A. Mole. "Real World Eff	<1%
5	<b>Crossref</b> 13 words Wang, Lan S, Lionel S. D'Souza, and Ira M. Jacobson. "Hepatitis C- A clinical review", <i>Journal of Medical Virology</i> ,	<1%
6	<b>Crossref</b> 13 words "Liver", <i>The American Journal of Gastroenterology</i> , 2015.	<1%
7	<b>Internet</b> 13 words crawled on 13-Mar-2014 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	<1%

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 2,970 条结果 (用时 0.84 秒)

## INITIAL TREATMENT OF HCV INFECTION | Recommendations for ...

[www.hcvguidelines.org/full-report/initial-treatment-hcv-infection](http://www.hcvguidelines.org/full-report/initial-treatment-hcv-infection) ▼ [翻译此页](#)

Initial treatment of HCV infection includes **patients** with chronic **hepatitis C** who have ... **Genotype 1a Treatment-Naïve Patients** Without **Cirrhosis** - Recommended ... are **non-black**, HIV-uninfected, and whose **HCV RNA** level is **<6 million IU/mL** ... **8 weeks** of **ledipasvir/sofosbuvir** who had baseline **HCV RNA** levels below **6** ...

您 17-4-25 访问过该网页。

## Hepatitis C Selected Articles - NATAP

[www.natap.org/hcv.htm](http://www.natap.org/hcv.htm) ▼

EASL: Elbasvir/Grazoprevir Plus Sofosbuvir ± Ribavirin in **Treatment-Naïve** And ... Glecaprevir and pibrentasvir for 12 **weeks** for **hepatitis C** virus **genotype 1** infection and prior ... CROI: Efficacy and **Safety** of Direct Antiviral Agents in **Cirrhotic HCV-HIV** ... AASLD: **Real World Effectiveness** of **Ledipasvir/Sofosbuvir** in **Patients** ...

## Core Concepts - Treatment of HCV Genotype 1 ... - Hepatitis C Online

[www.hepatitisc.uw.edu/go/treatment-infection/treatment-genotype-1/.../all](http://www.hepatitisc.uw.edu/go/treatment-infection/treatment-genotype-1/.../all) ▼ [翻译此页](#)作者: DH Spach - [相关文章](#)

2016年7月22日 - All of these regimens are safe and highly **effective**. ... If the **genotype 1** subtype is not known, the **patient** should be **treated** as genotype 1a. ... rate of 94 to 98% for **patients** with either **8** or **12 weeks** of **ledipasvir-sofosbuvir** among **non-cirrhotic patients** with baseline **HCV RNA** levels less than **6 million IU/mL**.

## Role of ledipasvir/sofosbuvir combination for genotype 1 hepatitis C ...



[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 4,630 条结果 (用时 0.46 秒)

## INITIAL TREATMENT OF HCV INFECTION | Recommendations for ...

[www.hcvguidelines.org/full-report/initial-treatment-hcv-infection](http://www.hcvguidelines.org/full-report/initial-treatment-hcv-infection) ▼ [翻译此页](#)

Initial treatment of HCV infection includes **patients** with chronic **hepatitis C** who have ... **Genotype 1a Treatment-Naïve Patients** Without **Cirrhosis** - Recommended ... are **non-black**, **HIV-uninfected**, and whose **HCV RNA** level is **<6 million IU/mL**. .... **8 weeks** of **ledipasvir/sofosbuvir** who had baseline **HCV RNA** levels below **6** ...

## Core Concepts - Treatment of HCV Genotype 1 ... - Hepatitis C Online

[www.hepatitisc.uw.edu/go/treatment-infection/treatment-genotype-1/.../all](http://www.hepatitisc.uw.edu/go/treatment-infection/treatment-genotype-1/.../all) ▼ [翻译此页](#)

作者: DH Spach - [相关文章](#)

2016年7月22日 - All of these regimens are **safe** and highly **effective**. ... If the **genotype 1** subtype is not known, the **patient** should be **treated** as genotype 1a. ... rate of 94 to 98% for **patients** with either **8** or **12 weeks** of **ledipasvir-sofosbuvir** among **non-cirrhotic patients** with baseline **HCV RNA** levels less than **6 million IU/mL**.

## Hepatitis C Selected Articles - NATAP

[www.natap.org/hcv.htm](http://www.natap.org/hcv.htm) ▼

Efficacy of **8 Weeks** of **Sofosbuvir**, **Velpatasvir**, and **Voxilaprevir** in **Patients** With .... **EASL: Real-world effectiveness** and cost per SVR of **Sofosbuvir/Velpatasvir** in ... for **8 weeks** in **treatment-naïve non-cirrhotic patients** with chronic **HCV** genotype ... Chronic **Hepatitis C Virus Genotype 1, 4, or 6** Infection (C-CORAL): A Phase 3 ...

## Summary for AASLD 2016 for Hepatitis C - New HCV two and three ...

[www.natap.org/2016/AASLD/AASLD\\_138.htm](http://www.natap.org/2016/AASLD/AASLD_138.htm) ▼ [翻译此页](#)

At present, besides the Abbvie 3-D regimen, mostly dual **hepatitis C (HCV)** direct acting .... Figure 2: Adherence rates to HCC surveillance in **HCV** infected **cirrhotics** .... **AASLD: Real World Effectiveness of Ledipasvir/Sofosbuvir in Patients** .... In **DAA-naïve patients** with **GT2** infection, **8 week treatment**