

World Journal of *Gastrointestinal Pathophysiology*

World J Gastrointest Pathophysiol 2017 November 15; 8(4): 150-187





REVIEW

- 150** Paneth cells in intestinal physiology and pathophysiology
Gassler N

ORIGINAL ARTICLE

Basic Study

- 161** Pharmacological inhibition of diacylglycerol acyltransferase-1 and insights into postprandial gut peptide secretion
Maciejewski BS, Manion TB, Steppan CM
- 176** Effects of commercially produced almond by-products on chemotherapy-induced mucositis in rats
Whittaker AL, Zhu Y, Howarth GS, Loung CS, Bastian SEP, Wirthensohn MG

Contents

World Journal of Gastrointestinal Pathophysiology
Volume 8 Number 4 November 15, 2017

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Pathophysiology*, Tao-Sheng Chen, PhD, Director, Department of Chemical Biology and Therapeutics, St. Jude Children's Research Hospital, Memphis, TN 38105, United States

AIM AND SCOPE

World Journal of Gastrointestinal Pathophysiology (*World J Gastrointest Pathophysiol*, *WJGP*, online ISSN 2150-5330, DOI: 10.4291), is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJGP is to report rapidly the most recent results in basic and clinical research on gastrointestinal pathophysiology, including all aspects of normal or abnormal function of the gastrointestinal tract, hepatobiliary system, and pancreas. *WJGP* specifically covers growth and development, digestion, secretion, absorption, metabolism and motility relative to the gastrointestinal organs, as well as immune and inflammatory processes, and neural, endocrine and circulatory control mechanisms that affect these organs. This journal will also report new methods and techniques in gastrointestinal pathophysiological research.

We encourage authors to submit their manuscripts to *WJGP*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Gastrointestinal Pathophysiology is now indexed in PubMed, PubMed Central.

FLYLEAF

I-IV Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ya-Jing Lu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Li-Jun Cui*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Gastrointestinal Pathophysiology

ISSN
ISSN 2150-5330 (online)

LAUNCH DATE
April 15, 2010

Frequency
Quarterly

EDITOR-IN-CHIEF
Thomas Y Ma, MD, PhD, Professor, Chief, Division of Gastroenterology and Hepatology, University of New Mexico, MSC10 5550, 1 UNM, Albuquerque, NM 87131, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/2150-5330/editorialboard.htm>

EDITORIAL OFFICE
Xiu-Xia Song, Director
World Journal of Gastrointestinal Pathophysiology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f0publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f0publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
November 15, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f0publishing.com>

Paneth cells in intestinal physiology and pathophysiology

Nikolaus Gassler

Nikolaus Gassler, Institute of Pathology, RWTH Aachen University, Braunschweig 38114, Germany

Author contributions: Gassler N contributed to the study idea, literature research, manuscript writing, and final revision.

Conflict-of-interest statement: None.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Nikolaus Gassler, MA, Professor, Institute of Pathology, RWTH Aachen University, Celler Strasse 38, Braunschweig 38114, Germany. ngassler@ukaachen.de
Telephone: +49-531-5953310
Fax: +49-531-5953449

Received: January 24, 2017

Peer-review started: January 26, 2017

First decision: May 2, 2017

Revised: July 28, 2017

Accepted: August 15, 2017

Article in press: August 16, 2017

Published online: November 15, 2017

Abstract

Small intestinal mucosa is characterised by villus forming connective tissues with highly specialised surface lining epithelial cells essentially contributing to the establishment of the intestinal border. In order to perform these diverse functions, spatially distinct compartments of epithelial differentiation are found along the crypt-villus axis, including Paneth cells as a highly specialised cell type. Paneth cells locate in crypts

and assist undifferentiated columnar cells, called crypt base columnar cells, and rapidly amplifying cells in the regeneration of absorptive and secretory cell types. There is some evidence that Paneth cells are involved in the configuration and function of the stem cell zone as well as intestinal morphogenesis and crypt fission. However, the flow of Paneth cells to crypt bottoms requires strong Wnt signalling guided by EphB3 and partially antagonised by Notch. In addition, mature Paneth cells are essential for the production and secretion of antimicrobial peptides including α -defensins/cryptdins. These antimicrobials are physiologically involved in shaping the composition of the microbiome. The autophagy related 16-like 1 (ATG16L1) is a genetic risk factor and is involved in the exocytosis pathway of Paneth cells as well as a linker molecule to PPAR signalling and lipid metabolism. There is evidence that injuries of Paneth cells are involved in the etiopathogenesis of different intestinal diseases. The review provides an overview of the key points of Paneth cell activities in intestinal physiology and pathophysiology.

Key words: Antimicrobial peptide; Paneth cell; Crohn's disease; Microbiome; Small intestine

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Paneth cells physiologically locate in small intestinal crypts. Wnt signalling promotes their differentiation and movement into crypts, whereas Notch activities antagonise Paneth cell maturation. The cells essentially contribute to crypt morphogenesis and intestinal homeostasis, sharpening the microbiome by secreting antimicrobial peptides, like defensins, and crypt fission. There is increasing molecular evidence that Paneth cells disorders are strongly involved in the pathophysiology of several intestinal diseases including ileal Crohn's disease and necrotising enterocolitis.

Gassler N. Paneth cells in intestinal physiology and pathophysiology. *World J Gastrointest Pathophysiol* 2017; 8(4): 150-160
Available from: URL: <http://www.wjgnet.com/2150-5330/full/v8/i4/150.htm> DOI: <http://dx.doi.org/10.4291/wjgp.v8.i4.150>

INTRODUCTION

The human small intestine is a multi-layered organ with complex cellularity. The intestinal mucosa is characterised by villus forming connective tissues with highly specialized surface lining epithelial cells. The epithelium is highly dynamic and interacts with the underlying mesenchyme and the luminal content. In order for these diverse functions to be performed, spatially distinct compartments of epithelial differentiation are found along the crypt-villus axis (CVA). Stem cells located in the crypts are responsible for the high throughput of surface lining cells within a few days. Histomorphologically, the CVA is a very characteristic mucosal feature and fundamental in explaining the structure and function of the surface lining epithelia. At present, four important cell types are defined, namely absorptive (enterocytes), mucosecreting (goblet cells), enteroendocrine, and Paneth cells^[1]. Some other epithelial and non-epithelial cell types are found at the mucosal surface, including M-cells, brush/tuft/caveolated cells, and several types of intraepithelial lymphocytes.

Enterocytes are most common in the surface epithelium and are responsible for digestion and absorption of nutrients as well as in forming the intestinal border. They are assisted by mucosecreting/goblet cells which produce several mucine types which are chemically different along the intestinal tract. Goblet cells increase in number from oral to aboral as the stool becomes increasingly compacted. Enteroendocrine cells represent the third important cell type and comprise a highly specialised chemosensory system involved in sensing the energy balance and reserve of an individual. A detailed classification of enteroendocrine subtypes is possible by the specific intestinal hormones that are produced and the assistance of Notch signalling in enteroendocrine cell differentiation^[2,3].

Paneth cells, first described by Gustav Albert Schwalbe in 1872, originate directly from intestinal stem cells^[4]. In contrast to other secretory cell types, Paneth cells are physiologically found at the bottom positions of small intestinal crypts (Figure 1). The unique histomorphological feature implicates special functions of Paneth cells in cellular homeostasis as well as in the establishment and configuration of the mucosal barrier as a physical and highly organised immune interface. Recently, some molecular mechanisms underlying secretory disorders of Paneth cells were identified strongly related to intestinal diseases, e.g., ileal Crohn's disease and necrotising enterocolitis. The important etiopathological findings were addressed and summarised with the term "Paneth's disease" and introduced in the literature^[4].

In the following paragraphs, important aspects of Paneth cell physiology and pathophysiology are reviewed. The data clearly demonstrate that Paneth cells are a highly specialized cell type strongly involved in assisting to sharpen and maintain of the microbiome as well as in the establishment of the stem cell niche




Figure 1 Normal small intestinal crypts with basal orientated Paneth cells. A: Paneth cells are characterised by their apical located granules (arrows). Between Paneth cells undifferentiated progenitor cells are found. In the normal *Lamina propria mucosae* a mixed population of immune cells and stroma resident cells is found; B: Occasionally, Paneth cells at the bottom of small intestinal crypts are mixed up with enteroendocrine cells (arrow). They are characterised by basal located granules. In the upper part of the crypt, a mitotic figure is shown.

and promotion of cellular renewal and mucosal morphogenesis. Consequently, Paneth cell disorders are involved in the pathophysiology of intestinal diseases.

PANETH AND STEM CELL NETWORK

The small intestinal epithelium renews within 3-6 d. The extraordinary rate of cell renewal is driven by a vigorous proliferation within crypts and a highly dynamic movement of epithelial columns toward the villus tip. The intestinal epithelia descend from a distinct stem cell zone located in small intestinal crypts. The zone consists of Paneth cells and 4 to 6 independent intestinal adult stem cells adjacent to rapidly cycling progenitors in the upper part of intestinal crypts.

The stem zone model is orientated on the morphological finding of crypt base columnar cells (CBC cells). These undifferentiated cycling cells are intermingled with Paneth cells and are hierarchically followed by Mix cells located directly above the Paneth cells^[5-7]. Mix cells are assumed to be strongly amplifying precursors of the different epithelial cell lines including Paneth cells.

In contrast to the stem cell zone model, a +4 position model has been suggested^[8]. The model was substantiated by the finding that severe radiation sensitivity exists in the +4 position^[9]. In this location, active cell cycling is found and radiation sensitivity indicates sufficient protection of the stem cell compartment from genetic damage. In the proposed model, injured +4 position stem cells are replaced by earlier generations of transit amplifying (TA) cells with a better repair capacity and asymmetric segregation of old and new DNA strands^[10]. Some parallels exist between the both models including definition of a slow and a rapid cycling cell type and an assisting role of Paneth cells in maintaining stem cell behaviour.

Maturing Paneth cells migrate downward into small intestinal crypts, where they reside for 3-6 wk^[11]. Paneth

cells escape from the crypt bottom by cellular fragmentation and phagocytosis from *lamina propria mucosae* infiltrating macrophages. There is experimental evidence that Wnt signalling and the expression of Wnt target genes are essential in the configuration and function of the stem cell zone including establishment of rapidly cycling TA cells^[4,12-14]. In a current model, an increasing gradient of Wnt activity directed into the small intestinal crypt is proposed reflecting the governing action of adjacent mesenchymal cells that release Wnt proteins. At the base of crypts, β -catenin is enriched in the nuclei of progenitor cells implying a strong response to Wnt signalling. The Wnt gradient is crucial for a graded expression of EphB2 and EphB3 acting as cell-sorting receptors along the CVA^[15]. In addition, graded Wnt activity is essential in the differentiation of Paneth cells with accumulation of large granules in the cytoplasm. Terminal differentiation of Paneth cells exists at the crypt bottom, where Wnt activity is maximised and several Wnt target genes are expressed^[16]. It has to be stressed that the expression pattern of Wnt target genes, especially *Lgr5/Gpr49*, is essential in the establishment of the small intestinal crypt stem cell zone including CBC cells interspersed between the maturing Paneth cells^[10]. Recently, intestinal organoids were used to visualize the short-range Wnt gradient in the intestinal stem cell niche^[17].

Paneth cells are important players in the intestinal stem cell niche, but they are identified as nonessential constituents. In a mouse model deficient in *Math1* (*Atoh1*), which totally and permanently lacks Paneth cells, *Lgr5*⁺ CBCs occupied the full crypt base, showed strong proliferation, and generated differentiated progeny over months^[18]. The increased proliferation along the CVA was associated with an increase in crypt height in the duodenum (13%) and ileum (15%). Enhanced or unaffected levels of Wnt target transcripts of genes associated with cell proliferation such as *CD44*, *Myc*, and *Ccnd1* were found. Using a different strategy of intestinal *Math1* deficiency, functional intestinal stem cells were found after Paneth cell ablation^[19]. In this study, *ex vivo* and *in vivo* experiments were performed to clarify the role of mesenchyme in the establishment of the stem cell niche. In the *ex vivo* setting, *Math1*-deficient crypt cells were not able to self-renew or survive indicating for extraepithelial, mesenchymal signals necessary for stem cell activities. Using caspase-8 deleted mesenchyme-free organoids, differentiation of Paneth cells was found^[20].

The Wnt dependency of Paneth cell differentiation is modified by other signalling cascades including the hierarchical Notch. The expression of Notch receptors is found in CBCs, and Paneth cells express the corresponding ligands^[21]. Consequently, Paneth cells are postulated as a source of Notch signalling to maintain CBC functions and crypt activities. It is suggested that some CBC functions unrelated to secretory metaplasia may require a Paneth cell source of Notch signalling.

Inhibition of Notch by pharmacological inactivation of the γ -secretase, an enzyme that is involved in Notch signalling, is already established. In dibenzazepine (DBZ)

- treated wild-type black-6 mice, a significant increase in secretory differentiated cells located throughout small intestinal crypts is found. After DBZ treatment, small intestinal crypts are lined by orthotopic Paneth cells and transdifferentiated cells with morphological features mixed from Paneth cells and mucus-retaining cells (Figure 2). The change in secretory differentiation is called secretory metaplasia.

PANETH CELL METAPLASIA

In distinct intestinal disorders, the physiology of Paneth cells is disturbed. In general, true injuries/diseases of Paneth cells should be separated from such intestinal disorders, where differentiation of Paneth cells is necessary to compensate cellular stress or misdirected mucosal differentiation/tissue homeostasis. The important histomorphological findings include loss, transdifferentiation or metaplasia of Paneth cells.

Metaplasia is defined as the occurrence of differentiated cells in a histomorphological location, where they are physiologically not found. Paneth cell metaplasia is seen throughout the gastro-intestinal tract, but frequently manifests in the stomach and is associated with different intestinal injuries (Figure 3). In addition, Paneth cell metaplasia is found in extra-gastro-intestinal sites, like the lung and tracheobronchial system, pancreatobiliary tract, and the uro-genital tract, but is a rarity in the nasopharyngeal system. It has been postulated that Notch may be involved in forms of pre-cancerous, metaplastic conditions in the stomach, which typically results from a loss of parietal cells and chronic inflammation. The expression of *Math1*, which is physiologically not found in normal gastric mucosa, has been shown to be upregulated in intestinal metaplasia of the stomach^[22,23].

Paneth cell loss is a hallmark of special types of Crohn's disease and is also found in acute inflammation like graft vs host disease (GvHD) grade II and III and ischemic tissue damage^[20,24]. The phenomenon of transdifferentiation, whereby Paneth cells are replaced by lysozyme-producing mucus cells, is described in inflammatory small intestinal disorders and in celiac disease. There is some phenomenological evidence that Paneth cell transdifferentiation could be due to re-programmed stem cells^[25].

At present, it is well established that the molecular framework underlying the process of regular Paneth cell differentiation and maturation strongly depends on Wnt activity^[16]. In contrast to the crypt physiology, the role of Wnt signalling in the development of Paneth cell metaplasia is only marginally elucidated. However, there is some evidence of Wnt activity as a driving force in metaplasia. In an experimental study with the constitutive expression of a β -catenin-Lef1 fusion protein under control of a lung-endoderm-specific promoter from the surfactant protein C gene transgenic lungs included cells expressing marker genes strongly characteristic of intestinal Paneth cells^[26]. The data strongly supports the view that hyperactive Wnt signalling could be crucial




Figure 2 Differentiation of Paneth cells and Notch inhibition. Normal ileal mouse mucosa with normal crypts and Paneth cells: HE staining (A) and alcian-PAS staining (B). Arrows indicate Paneth cells. Ileal mouse tissues after treatment with dibenzazepine show an increase in secretory cells in the crypts with differentiation of Paneth cell-like epithelia (arrows): HE staining (C) and alcian-PAS staining (D).

in stem cell lineage commitment and the generation of intestinal metaplasia. In the lung study, where lysozyme was used as a marker protein of mature Paneth cells, increased Wnt activity was associated with diminished lysozyme expression in metaplastic cells^[26]. This interesting finding is highly suggestive that Wnt activity is an important prerequisite for Paneth cell metaplasia, but additional factors and supportive signalling mechanisms are necessary. The auxiliary components necessary for Paneth cell differentiation and maturation probably differs between tissues. For example, inflammation seems to induce Paneth cell metaplasia/hyperplasia in the colon, but not in the small intestine^[27]. The intestinal microbiome is suggested as an important variable in signalling activities and the expression level of several intestinal pathways and regulates phenomena including inflammation and cellular differentiation.

PANETH CELLS AND ANTIMICROBIAL PEPTIDES

In the intestine, a plethora of microbes is found including autochthonous long-term colonisers and allochthonous transient microorganisms. One important function of commensal microbiota is host defence through the promotion of the mucosal immune system and inhibition of pathological microbes. Another function is the nutrition of the host, because the bacteria have the capacity to ferment components of the diet. The synthesised short-chain fatty acids (SCFAs), vitamins, and amino acids are essentials in organising host physiology and energy balance. SCFAs act as signalling molecules and highly specialised free fatty acid receptors (FFARs) exist in the intestinal epithelium. FFARs contribute to the chemosensory intestinal system and differ in their molecular structure, ligand specificity, and functional properties^[28]. There is evidence that SCFAs are able to induce glucagon-like peptide-1 release from intestinal cells. The molecular link is important to display the integral role of the intestinal microbiome in the regulation host's energy homeostasis. Metabolic disorders and obesity are

associated with changes in the gut microbiota^[29].

In the intestine, a complex molecular system is established to maintain microbiome - host homeostasis and to shape the composition of microbes colonising the intestine. For this purpose, Paneth cells sufficiently express and secrete different types of antimicrobial peptides (AMPs) which are important host-defence substances in the communication between host and microbiome. The secretory capacity of Paneth cells is reflected by the cytoarchitecture with apical clustering of large secretory granules. Ultrastructurally, the endoplasmic reticulum (ER) is hyperplastic and associated with a well-developed Golgi network. Upon prosecretory stimuli like bacterially derived Toll-like receptor ligands the cytosolic calcium content increases *via* KCa3.1, a calcium-activated potassium channel, and leads to granule secretion^[30-33].

Defensins are the major AMP family and contain six cysteines involved in intramolecular disulphide bonds^[34,35]. Two major subfamilies of defensins exist. The α -defensins are strongly synthesised by Paneth cells and neutrophils, whereas β -defensins with constitutive HBD-1 and inducible HBD-2 are found in different types of cells including enterocytes^[36]. In mice, α -defensins are named cryptidins or cryptidins which consist of six major isoforms. Among those isoforms, variant 4 is the most microbicidal peptide^[37]. In addition to α -defensins, Paneth cell granules contain lysozyme as another potent host-defence molecule. The biosynthesis of active α -defensins depends on proteolytic processing. In murine Paneth cells, matrix metalloproteinase 7 (MMP7) is found cleaving defensin precursor molecules to active α -defensins^[38,39]. Disturbed proteolytic activity as found in MMP7 knockout mice is associated with a hampered intestinal microbiome and an increased distribution of bacterial pathogens^[40]. In contrast to mice, trypsin is found in human Paneth cells as an activating protease for prodefensins and the antimicrobial protein Reg3A^[41]. The human α -defensins type 5 (HD5) and 6 (HD6) are essential in host protection from intestinal pathogens. Using a HD5 transgenic mouse model with physiological defensin levels, animals were resistant to enteric *Salmonella* infections due to reduced viability of bacteria, diminished bacterial translocation,




Figure 3 Examples of Paneth cell metaplasia throughout the intestinal tract. A: Paneth cell metaplasia (arrows) in Barrett mucosa. Squamous epithelia of the oesophagus are marked with an arrowhead; B: Chronic atrophic gastritis with Paneth cell metaplasia (arrow); C: Paneth cell metaplasia (arrows) of Brunner's gland. In the upper part, small intestinal mucosa and secretory ducts are shown (arrowheads); D: Colon mucosa in ulcerative colitis with disturbed crypt architecture, increased numbers of stroma infiltrating inflammatory cells, and Paneth cell metaplasia (arrowheads); E: Higher magnification of ulcerative colitis-associated Paneth cell metaplasia (arrows) in colon mucosa as demonstrated in (D); F: Paneth cell metaplasia (arrows) in tubular adenoma of the colon with low-grade dysplasia.

and increased survival after lethal *Salmonella* infections^[42]. In the same model, evidence of a direct role of α -defensins in regulating and shaping the small intestinal microbiome was shown^[40]. In the intestine of HD5 transgenic animals the number of commensals was changed with an increase in *Bacteroides*, a decrease in *Firmicutes*, and the loss of segmented filamentous bacteria (SFB), designated *Candidatus arthromitus*, accompanied by an absence of Th17 cell differentiation. SFB are host specific intestinal symbionts belonging to the *Clostridiaceae* displaying differentiation of filaments that interact intimately with the surface lining intestinal epithelia. Secreted proteins are expressed by SFB including different ADP-ribosyltransferases and a myosin-cross-reactive antigen, all probably involved in modifying the host response and post-natal maturation of the gut immune system^[43,44]. Recently, an SFB-host cell co-culturing system was established producing viable infectious particles, which can colonise intestinal mucosa with the induction of an immune response^[45]. Regional variation in the expression and secretion of Paneth cell AMPs along the intestinal tract are well-balanced by the colonising activity of SFB with the establishment and shaping the intestinal microbiome^[46].

The α -defensin HD6 secreted by human Paneth cells acts in a way that is completely different to HD5. The HD6 is antibacterial by the configuration of self-assembled peptide nanonets and formation of nanofibrils^[47,48].

Targeted bacteria surrounded by the fibrils and nanonets are not able to invade the intestinal mucosa. The HD6 self-assembling function is essentially based on the presence of histidine-27, which forms a salt bridge necessary for multimerising the peptide. The important Paneth cell produced α -defensins HD5 and HD6 can be further classified by their unique functions. The antimicrobial HD5 activity with disruption of bacterial membranes is called "harpoon" activity, whereas HD6 represents so-called "net forming" activities^[49]. In summary, HD5 acts antimicrobially and exhibits lectin-like properties, whereas HD6 entraps microbes and blocks host cell invasion^[50,51].

In addition to defensins, Paneth cells are able to secrete other AMPs including lysozyme, secretory phospholipase A2 (sPLA2), RegIII, angiogenin 4, and cathelicidins^[52-54]. Among the AMPs, RegIII proteins are important in antibacterial defence, and murine RegIII γ shares 65% identity with human RegIII α . RegIII proteins belong to the family of C-type lectin regenerating islet-derived proteins and bind glycan chains of peptidoglycans on the cell wall of gram-positive bacteria. In contrast to other C-type lectin AMPs as mannose-binding lectin, the complement recruitment domains are not constantly expressed in RegIII suggesting a direct anti-bactericidal function^[55].

The AMP LL-37 belongs to the cathelicidin family and acts in a similar way as HD5 by puncturing holes in microbial membranes. LL-37 mediates antimicrobial

activities in addition to immunological functions *via* various cellular receptors^[56]. LL-37 is an inducible AMP, because LL-37 expression increases in the presence of SCFAs. Consequently, the protein is predominantly found in the transit-amplifying zone in colonic crypts, when compared with small intestinal mucosa^[57]. In quantitative terms, the ratio of HD5 to HD6 is about 3:1, whereas HD5 expression levels are higher by a factor of up to 100 than those of lysozyme and sPLA2^[4].

Paneth cells are secretory cells with the excretion of a plethora of molecules including several types of AMPs^[50,52-54]. The secretory function depends on a well-adapted ER, which is controlled and smoothed by autophagy. Disruption of autophagy by the deletion of the unfolded protein response transcription factor X-box binding protein-1 results in ER stress, Paneth cell impairment, and spontaneous inflammation resembling special variants of Crohn's disease^[58]. There is experimental evidence that lipids and lipid metabolising enzymes are involved in the signalling cascade for exocytosis of granules from Paneth cells^[59]. Expression profiling revealed a panel of target genes and related proteins including lipoprotein lipase, apolipoprotein A-IV, stearoyl-CoA desaturase-1, adiponectin, and leptin. Mapped pathways include PPAR signalling, statin pathway, adipocytokine signalling, and polyunsaturated fatty acid biosynthesis. It has been speculated that the lipid-associated exocytosis is an additive mechanism to the chemosensory system of FFARs and enteroendocrine cells to perform synergisms between the intestinal microbiome and host metabolism.

The release of AMPs from Paneth cells into the intestinal mucus and lumen is initiated by stimulation of pattern recognition receptors (PRRs) located on intestinal surfaces. The class of PRRs includes Toll-like receptors (TLRs) activated by lipopolysaccharide and nucleotide-binding oligomerisation domain-containing molecules (NODs), which recognise muramyl dipeptide. Independently of PRRs, bacterial cell wall glycolipids are able to stimulate the release of defensins by Paneth cells. The myeloid differentiation primary response gene (MyD88) acts as an important cytoplasmic TLR limiting bacterial penetration into the intestinal mucosa^[4].

In summary, Paneth cells are essential for the establishment of a mucus layer enriched with AMPs resting on the surface lining epithelia. The composition of AMPs differs along the intestine, implicating diverse functions in communication with the intestinal microbiome.

PANETH CELLS AND INTESTINAL DISEASES

High numbers of *Lamina propria mucosae* infiltrating leukocytes with different types of specialised lymphocytes, plasma cells, mast cells, monocytes, and eosinophiles is a very characteristic histomorphological feature of intestinal mucosa. The balanced correspondence of the immune cells with the intestinal microbiome *via* the surface

epithelium is a prerequisite for intestinal homeostasis. In consequence, an inflammatory response with a further increase in leukocytes including neutrophils is very common in intestinal disorders and is established in the majority of intestinal diseases. There are remarkable differences in the quantity and quality of infiltrating leukocytes between the underlying intestinal disorders. The differences of leukocytes are helpful to classify the basal disease using morphological, immunohistochemical, and functional techniques.

In addition to resting Paneth cells, infiltrating leukocytes are an important transient source for AMPs. They are assisted by metaplastic Paneth cells, found in the large intestine and stomach. In intestinal inflammation, sPLA2, which is physiologically not found in the colon, is expressed by metaplastic Paneth cells^[60,61]. There are experimental data demonstrating a direct role of the microbiome in regulating the defensin production and secretion. Using *Nod2* knockout mice in co-housing experiments, the wild-type microbiome was able to regulate defensin secretion to physiological levels^[62]. The inflammation-related expression profile of AMPs includes some characteristics indicating a defined intestinal disease^[63,64]. For example, β -defensin types 2, 3, and 4 are increased in ulcerative colitis but not in Crohn's disease^[65]. In summary *Nod2* is highly expressed in ileal Paneth cells that essentially contribute to the regulation of ileal microbiota through the secretion of AMPs^[66].

As outlined above, Paneth cells are the main source of AMPs and acts in stabilising the stem cell zone. Consequently, Paneth cells are investigated as a site of origin for intestinal inflammation. This point of view seems of high relevance concerning the pathogenesis of inflammatory bowel diseases (IBD). A fundamental feature of ileal Crohn's disease is a reduced expression of HD5 and HD6. The finding is sometimes paralleled by a reduced Paneth cell number, but may be also a result of an injured microbiome^[4,36,67].

An in-detail analysis of Paneth cells in Crohn's disease revealed subgroups of the disease characterised by unique molecular, morphological, and clinical features. The clinical phenotype of ileal Crohn's disease is especially associated with Paneth cell injuries. There was first evidence from loss-of-function mutations in *NOD2* (SNP 13), a gene that is strongly expressed by Paneth cells^[66,68-70]. Subsequently, the molecular mechanisms underlying several injuries of Paneth cells were described and linked to (ileal) Crohn's disease. One important finding was low Wnt signalling activity in ileal Crohn's disease with diminished Tcf-4 and reduced secretion of defensins^[71,72]. Another milestone was the identification of defective granule exocytosis from Paneth cells with diminished levels of defensins due to abnormal autophagy in homozygosity for the risk allele autophagy related 16-like 1^[59,73]. The molecular characterisation of disturbed autophagy and unresolved ER stress due to genetically and environmentally controlled dysfunction of unfolded protein response (UPR) was the breakthrough in understanding the underlying mechanisms for

the threshold for the development of (ileal) Crohn's disease^[58,59,74].

In addition to Paneth cells, enterocytes are an additional source to produce AMPs including Reg proteins, defensins, and cathelicidins^[50,52-54]. This fact is of high relevance to physiologically compensate the perinatal period prior to the establishment of Paneth cells and to pathologically balance Paneth cell disorders^[75]. Paneth cells are found in the prenatal small intestine, but the number of cells is highly variable. In view of this observation, insufficient Paneth cell maturation is in discussion as an important variable in the pathogenesis of necrotising enterocolitis. The multifactorial disease is preferentially found in premature infants. Clinically, a sudden onset and high mortality in babies born after 35 weeks' gestation are characteristic.

At present, two different pathophysiological models for the development of necrotising enterocolitis are in discussion; the top-down and the bottom-up hypothesis^[76]. In the top-down scenario, bacterial invasion is found in the villus that triggers thrombosis of small arteries via activation of platelet-activating factor (PAF). In contrast to the top-down scenario, bacterial translocation at the crypt basis with consecutive activation of Paneth cells and arterial thrombosis are the key points in the bottom-up model. In very preterm neonates, the time period between birth and onset of the necrotising enterocolitis is frequently long. In this time maturation of the intestinal immune system is found, a pre-requisite for the initiation and development of enterocolitis^[77]. In animal models, toxic Paneth cell ablation and gastral infection with *Klebsiella pneumoniae* is able to induce necrotising enterocolitis in a bottom-up scenario^[78]. The variable number of Paneth cells found in human necrotising enterocolitis is discussed as enterocolitis associated necrobiosis and does not contradict the bottom-up model^[79,80]. It has to be stressed that the immense damage to intestinal tissues in necrotising enterocolitis is not plausible from the thrombosis of small arteries. Therefore, a two-hit model has been proposed including at first ischemic damage of the intestinal mucosa. In a second step, reperfusion is found with uncontrolled activation of signalling cascades and enzymes aggravating the tissue damage^[81].

Barrett mucosa is frequently found in the oesophageal junction and is due to chronic gastroesophageal reflux. Histomorphologically, the epithelial cells display intestinal differentiation with goblet cells and aberrant mucus retention^[82]. In contrast to complete intestinal metaplasia, brush border and Paneth cells are not commonly found in Barrett mucosa. However, there is evidence of Paneth cell metaplasia in Barrett oesophagus (Figure 3A). The phenomenon is found in about 30% of the cases^[83]. The frequency of Paneth cells in Barrett mucosa is inversely correlated to the development of epithelial dysplasia. In long-segment Barrett oesophagus, the metaplasia is more frequent than in short-segment lesions. The persistence of Barrett mucosa when Paneth cell metaplasia is found could be due to a dys-balanced Notch and Wnt signalling^[84,85]. There is no evidence that Paneth cell metaplasia is

associated with the progression of Barrett mucosa to dysplasia and adenocarcinoma^[83].

In GvHD the intestinal tract is frequently affected and Paneth cell loss is a hallmark of the acute inflammation associated with changes in the composition of the microbiome^[86,87]. Due to the loss of Paneth cells a striking disease of α -defensins is found favouring a shift from commensal to pathologic microbiota^[88]. The dysbiotic shift is assumed to be crucial for severe septic complications in GvHD. In particular, the diminished number of Paneth cells correlates with higher disease severity and poor treatment response in patients suffering from GvHD^[89]. Enteroendocrine cells have been recently discovered as sensors of the intestinal microbiome. The cells are involved in the pathogenesis of intestinal disorders and express specific receptors which can respond to bacterial products^[90]. In germ-free mice, the number of enteroendocrine cells is drastically reduced^[91], whereas an increase in microbiome metabolites such as in GvHD is inductive for secretion of immunomodulatory enteroendocrine hormone peptides^[92]. The inductive mechanisms could be a pathophysiological link for hyperplasia of enteroendocrine cells in GvHD.

PANETH CELLS AND MORPHOGENESIS

Crypt fission, the division of a single crypt into two daughters, is fundamental in intestinal tissue expansion and morphogenesis, but is also found in tumourigenesis driving the clonal expansion of mutant adenomatous crypts^[93-96]. A specific cellular arrangement in the intestinal stem cell niche is observed that controls crypt fission. In a recent model, Paneth cells and CBCs (Lgr5+) are both importantly involved in crypt morphogenesis^[97]. The findings summarise the data from intestinal organoids, where Paneth cells are essentially involved in crypt budding^[98,99], and as mice lacking intestinal Paneth cells, which are able to sufficiently repair crypt injuries^[21,100]. The morphogenic activities of Paneth cells depend on Wnt signalling and redundant sources contribute^[101,102]. Important for the crypt fission model is the observation that Paneth cells adhere to their substrate more strongly than other crypt cells. In view with this strong evidence is given that so-called "mislocalised" Paneth cells change the symmetry of fission in determining the site of fission^[97].

CONCLUSION

Paneth cells are important secretory cells in the small intestinal mucosa. They are found as metaplastic cells in several other locations. Important secretory products of Paneth cells belong to the AMPs. They are essential for the dialogue between the microbiome and the host. In addition to Paneth cell functions in controlling the microbiome, the cells are functionally and structurally involved in forming the stem cell zone of small intestinal crypts and involved in morphogenesis of the CVA. In the last decade, molecular mechanisms of different Paneth cell injuries have been identified and correlated with intestinal

diseases. There is evidence from these experiments that ileal Crohn's disease as well as necrotising enterocolitis are strongly associated with diseased Paneth cells. The important role of Paneth cells in intestinal physiology and pathophysiology strongly supports the view that these cells are the guardian of small intestinal crypts.

REFERENCES

- 1 **Sancho E**, Batlle E, Clevers H. Live and let die in the intestinal epithelium. *Curr Opin Cell Biol* 2003; **15**: 763-770 [PMID: 14644203 DOI: 10.1016/j.cceb.2003.10.012]
- 2 **Schonhoff SE**, Giel-Moloney M, Leiter AB. Minireview: Development and differentiation of gut endocrine cells. *Endocrinology* 2004; **145**: 2639-2644 [PMID: 15044355 DOI: 10.1210/en.2004-0051]
- 3 **Beehler-Evans R**, Micchelli CA. Generation of enteroendocrine cell diversity in midgut stem cell lineages. *Development* 2015; **142**: 654-664 [PMID: 25670792 DOI: 10.1242/dev.114959]
- 4 **Wehkamp J**, Stange EF. Paneth's disease. *J Crohns Colitis* 2010; **4**: 523-531 [PMID: 21122555 DOI: 10.1016/j.crohns.2010.05.010]
- 5 **Bjerknes M**, Cheng H. Clonal analysis of mouse intestinal epithelial progenitors. *Gastroenterology* 1999; **116**: 7-14 [PMID: 9869596 DOI: 10.1016/S0016-5085(99)70222-2]
- 6 **Bjerknes M**, Cheng H. Intestinal epithelial stem cells and progenitors. *Methods Enzymol* 2006; **419**: 337-383 [PMID: 17141062 DOI: 10.1016/S0076-6879(06)19014-X]
- 7 **Bjerknes M**, Cheng H. Neurogenin 3 and the enteroendocrine cell lineage in the adult mouse small intestinal epithelium. *Dev Biol* 2006; **300**: 722-735 [PMID: 17007831 DOI: 10.1016/j.ydbio.2006.07.040]
- 8 **Potten CS**. Extreme sensitivity of some intestinal crypt cells to X and gamma irradiation. *Nature* 1977; **269**: 518-521 [PMID: 909602]
- 9 **Potten CS**, Kovacs L, Hamilton E. Continuous labelling studies on mouse skin and intestine. *Cell Tissue Kinet* 1974; **7**: 271-283 [PMID: 4837676]
- 10 **Barker N**, van de Wetering M, Clevers H. The intestinal stem cell. *Genes Dev* 2008; **22**: 1856-1864 [PMID: 18628392 DOI: 10.1101/gad.1674008]
- 11 **Barker N**. The canonical Wnt/beta-catenin signalling pathway. *Methods Mol Biol* 2008; **468**: 5-15 [PMID: 19099242 DOI: 10.1007/978-1-59745-249-6_1]
- 12 **van de Wetering M**, Sancho E, Verweij C, de Lau W, Oving I, Hurlstone A, van der Horn K, Batlle E, Coudreuse D, Haramis AP, Tjon-Pon-Fong M, Moerer P, van den Born M, Soete G, Pals S, Eilers M, Medema R, Clevers H. The beta-catenin/TCF-4 complex imposes a crypt progenitor phenotype on colorectal cancer cells. *Cell* 2002; **111**: 241-250 [PMID: 12408868 DOI: 10.1016/S0092-8674(02)01014-0]
- 13 **Kuhnert F**, Davis CR, Wang HT, Chu P, Lee M, Yuan J, Nusse R, Kuo CJ. Essential requirement for Wnt signaling in proliferation of adult small intestine and colon revealed by adenoviral expression of Dickkopf-1. *Proc Natl Acad Sci USA* 2004; **101**: 266-271 [PMID: 14695885 DOI: 10.1073/pnas.2536800100]
- 14 **Gregorieff A**, Pinto D, Begthel H, Destree O, Kielman M, Clevers H. Expression pattern of Wnt signaling components in the adult intestine. *Gastroenterology* 2005; **129**: 626-638 [PMID: 16083717 DOI: 10.1053/j.gastro.2005.06.007]
- 15 **Battle E**, Henderson JT, Begthel H, van den Born MM, Sancho E, Huls G, Meeldijk J, Robertson J, van de Wetering M, Pawson T, Clevers H. Beta-catenin and TCF mediate cell positioning in the intestinal epithelium by controlling the expression of EphB/ephrinB. *Cell* 2002; **111**: 251-263 [PMID: 12408869 DOI: 10.1016/S0092-8674(02)01015-2]
- 16 **van Es JH**, Jay P, Gregorieff A, van Gijn ME, Jonkheer S, Hatzis P, Thiele A, van den Born M, Begthel H, Brabletz T, Taketo MM, Clevers H. Wnt signalling induces maturation of Paneth cells in intestinal crypts. *Nat Cell Biol* 2005; **7**: 381-386 [PMID: 15778706 DOI: 10.1038/ncb1240]
- 17 **Farin HF**, Jordens I, Mosa MH, Basak O, Korving J, Tauriello DV, de Punder K, Angers S, Peters PJ, Maurice MM, Clevers H. Visualization of a short-range Wnt gradient in the intestinal stem-cell niche. *Nature* 2016; **530**: 340-343 [PMID: 26863187 DOI: 10.1038/nature16937]
- 18 **Kim TH**, Escudero S, Shivdasani RA. Intact function of Lgr5 receptor-expressing intestinal stem cells in the absence of Paneth cells. *Proc Natl Acad Sci USA* 2012; **109**: 3932-3937 [PMID: 22355124 DOI: 10.1073/pnas.1113890109]
- 19 **Sato T**, van Es JH, Snippert HJ, Stange DE, Vries RG, van den Born M, Barker N, Shroyer NF, van de Wetering M, Clevers H. Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature* 2011; **469**: 415-418 [PMID: 21113151 DOI: 10.1038/nature09637]
- 20 **Günther C**, Martini E, Wittkopf N, Amann K, Weigmann B, Neumann H, Waldner MJ, Hedrick SM, Tenzer S, Neurath MF, Becker C. Caspase-8 regulates TNF- α -induced epithelial necroptosis and terminal ileitis. *Nature* 2011; **477**: 335-339 [PMID: 21921917 DOI: 10.1038/nature10400]
- 21 **Durand A**, Donahue B, Peignon G, Letourneur F, Cagnard N, Slomianny C, Perret C, Shroyer NF, Romagnolo B. Functional intestinal stem cells after Paneth cell ablation induced by the loss of transcription factor Math1 (Atoh1). *Proc Natl Acad Sci USA* 2012; **109**: 8965-8970 [PMID: 22586121 DOI: 10.1073/pnas.1201652109]
- 22 **Mutoh H**, Sakamoto H, Hayakawa H, Arai Y, Satoh K, Nokubi M, Sugano K. The intestine-specific homeobox gene Cdx2 induces expression of the basic helix-loop-helix transcription factor Math1. *Differentiation* 2006; **74**: 313-321 [PMID: 16831200 DOI: 10.1111/j.1432-0436.2006.00074.x]
- 23 **Demitrack ES**, Samuelson LC. Notch regulation of gastrointestinal stem cells. *J Physiol* 2016; **594**: 4791-4803 [PMID: 26848053 DOI: 10.1111/JP271667]
- 24 **Grootjans J**, Hodin CM, de Haan JJ, Derikx JP, Rouschop KM, Verheyen FK, van Dam RM, Dejong CH, Buurman WA, Lenaerts K. Level of activation of the unfolded protein response correlates with Paneth cell apoptosis in human small intestine exposed to ischemia/reperfusion. *Gastroenterology* 2011; **140**: 529-539.e3 [PMID: 20965186 DOI: 10.1053/j.gastro.2010.10.040]
- 25 **Rubio CA**. Lysozyme-rich mucus metaplasia in duodenal crypts supersedes Paneth cells in celiac disease. *Virchows Arch* 2011; **459**: 339-346 [PMID: 21769618 DOI: 10.1007/s00428-011-1129-3]
- 26 **Okubo T**, Hogan BL. Hyperactive Wnt signaling changes the developmental potential of embryonic lung endoderm. *J Biol* 2004; **3**: 11 [PMID: 15186480 DOI: 10.1186/jbiol3]
- 27 **Perminow G**, Beisner J, Koslowski M, Lyckander LG, Stange E, Vatn MH, Wehkamp J. Defective paneth cell-mediated host defense in pediatric ileal Crohn's disease. *Am J Gastroenterol* 2010; **105**: 452-459 [PMID: 19904243 DOI: 10.1038/ajg.2009.643]
- 28 **Kaemmerer E**, Plum P, Klaus C, Weiskirchen R, Liedtke C, Adolf M, Schippers A, Wagner N, Reinartz A, Gassler N. Fatty acid binding receptors in intestinal physiology and pathophysiology. *World J Gastrointest Pathophysiol* 2010; **1**: 147-153 [PMID: 21607156 DOI: 10.4291/wjgp.v1.i5.147]
- 29 **Kuwahara A**. Contributions of colonic short-chain Fatty Acid receptors in energy homeostasis. *Front Endocrinol (Lausanne)* 2014; **5**: 144 [PMID: 25228897 DOI: 10.3389/fendo.2014.00144]
- 30 **Satoh Y**, Habara Y, Ono K, Kanno T. Carbamylcholine- and catecholamine-induced intracellular calcium dynamics of epithelial cells in mouse ileal crypts. *Gastroenterology* 1995; **108**: 1345-1356 [PMID: 7729625]
- 31 **Ayabe T**, Satchell DP, Wilson CL, Parks WC, Selsted ME, Ouellette AJ. Secretion of microbicidal alpha-defensins by intestinal Paneth cells in response to bacteria. *Nat Immunol* 2000; **1**: 113-118 [PMID: 11248802]
- 32 **Ayabe T**, Wulff H, Darmoul D, Cahalan MD, Chandy KG, Ouellette AJ. Modulation of mouse Paneth cell alpha-defensin secretion by mIKCa1, a Ca²⁺-activated, intermediate conductance potassium channel. *J Biol Chem* 2002; **277**: 3793-3800 [PMID: 11724775 DOI: 10.1074/jbc.M107507200]
- 33 **Ouellette AJ**. Paneth cell alpha-defensin synthesis and function. *Curr Top Microbiol Immunol* 2006; **306**: 1-25 [PMID: 16909916]
- 34 **Ganz T**. Defensins: antimicrobial peptides of innate immunity. *Nat Rev Immunol* 2003; **3**: 710-720 [PMID: 12949495 DOI: 10.1038/nri1180]

- 35 **Wong JH**, Xia L, Ng TB. A review of defensins of diverse origins. *Curr Protein Pept Sci* 2007; **8**: 446-459 [PMID: 17979760]
- 36 **Wehkamp J**, Fellermann K, Herrlinger KR, Bevins CL, Stange EF. Mechanisms of disease: defensins in gastrointestinal diseases. *Nat Clin Pract Gastroenterol Hepatol* 2005; **2**: 406-415 [PMID: 16265431]
- 37 **Masuda K**, Nakamura K, Yoshioka S, Fukaya R, Sakai N, Ayabe T. Regulation of microbiota by antimicrobial peptides in the gut. *Adv Otorhinolaryngol* 2011; **72**: 97-99 [PMID: 21865701 DOI: 10.1159/000324625]
- 38 **Shirafuji Y**, Tanabe H, Satchell DP, Henschen-Edman A, Wilson CL, Ouellette AJ. Structural determinants of pro-cryptdin recognition and cleavage by matrix metalloproteinase-7. *J Biol Chem* 2003; **278**: 7910-7919 [PMID: 12482850 DOI: 10.1074/jbc.M210600200]
- 39 **Weeks CS**, Tanabe H, Cummings JE, Crampton SP, Sheynis T, Jelinek R, Vanderlick TK, Cocco MJ, Ouellette AJ. Matrix metalloproteinase-7 activation of mouse paneth cell pro-alpha-defensins: SER43 down arrow ILE44 proteolysis enables membrane-disruptive activity. *J Biol Chem* 2006; **281**: 28932-28942 [PMID: 16822871 DOI: 10.1074/jbc.M602041200]
- 40 **Salzman NH**, Hung K, Haribhai D, Chu H, Karlsson-Sjöberg J, Amir E, Tegatz P, Barman M, Hayward M, Eastwood D, Stoel M, Zhou Y, Sodergren E, Weinstock GM, Bevins CL, Williams CB, Bos NA. Enteric defensins are essential regulators of intestinal microbial ecology. *Nat Immunol* 2010; **11**: 76-83 [PMID: 19855381 DOI: 10.1038/ni.1825]
- 41 **Ghosh D**, Porter E, Shen B, Lee SK, Wilk D, Drazba J, Yadav SP, Crabb JW, Ganz T, Bevins CL. Paneth cell trypsin is the processing enzyme for human defensin-5. *Nat Immunol* 2002; **3**: 583-590 [PMID: 12021776 DOI: 10.1038/ni797]
- 42 **Salzman NH**, Ghosh D, Huttner KM, Paterson Y, Bevins CL. Protection against enteric salmonellosis in transgenic mice expressing a human intestinal defensin. *Nature* 2003; **422**: 522-526 [PMID: 12660734 DOI: 10.1038/nature01520]
- 43 **Pamp SJ**, Harrington ED, Quake SR, Relman DA, Blainey PC. Single-cell sequencing provides clues about the host interactions of segmented filamentous bacteria (SFB). *Genome Res* 2012; **22**: 1107-1119 [PMID: 22434425 DOI: 10.1101/gr.131482.111]
- 44 **Schnupf P**, Gaboriau-Routhiau V, Cerf-Bensussan N. Host interactions with Segmented Filamentous Bacteria: an unusual trade-off that drives the post-natal maturation of the gut immune system. *Semin Immunol* 2013; **25**: 342-351 [PMID: 24184014 DOI: 10.1016/j.smim.2013.09.001]
- 45 **Schnupf P**, Gaboriau-Routhiau V, Gros M, Friedman R, Moya-Nilges M, Nigro G, Cerf-Bensussan N, Sansonetti PJ. Growth and host interaction of mouse segmented filamentous bacteria in vitro. *Nature* 2015; **520**: 99-103 [PMID: 25600271 DOI: 10.1038/nature14027]
- 46 **Karlsson J**, Pütsep K, Chu H, Kays RJ, Bevins CL, Andersson M. Regional variations in Paneth cell antimicrobial peptide expression along the mouse intestinal tract. *BMC Immunol* 2008; **9**: 37 [PMID: 18637162 DOI: 10.1186/1471-2172-9-37]
- 47 **Chu H**, Pazgier M, Jung G, Nuccio SP, Castillo PA, de Jong MF, Winter MG, Winter SE, Wehkamp J, Shen B, Salzman NH, Underwood MA, Tsois RM, Young GM, Lu W, Lehrer RI, Bäuml AJ, Bevins CL. Human α -defensin 6 promotes mucosal innate immunity through self-assembled peptide nanonets. *Science* 2012; **337**: 477-481 [PMID: 22722251 DOI: 10.1126/science.1218831]
- 48 **Ouellette AJ**, Selsted ME. Immunology. HD6 defensin nanonets. *Science* 2012; **337**: 420-421 [PMID: 22837514 DOI: 10.1126/science.1225906]
- 49 **Bevins CL**. Innate immune functions of α -defensins in the small intestine. *Dig Dis* 2013; **31**: 299-304 [PMID: 24246978 DOI: 10.1159/000354681]
- 50 **Chairatana P**, Nolan EM. Defensins, lectins, mucins, and secretory immunoglobulin A: microbe-binding biomolecules that contribute to mucosal immunity in the human gut. *Crit Rev Biochem Mol Biol* 2017; **52**: 45-56 [PMID: 27841019 DOI: 10.1080/10409238.2016.124365]
- 51 **Nakamura K**, Sakuragi N, Takakuwa A, Ayabe T. Paneth cell α -defensins and enteric microbiota in health and disease. *Biosci Microbiota Food Health* 2016; **35**: 57-67 [PMID: 27200259 DOI: 10.12938/bmfh.2015-019]
- 52 **Peterson LW**, Artis D. Intestinal epithelial cells: regulators of barrier function and immune homeostasis. *Nat Rev Immunol* 2014; **14**: 141-153 [PMID: 24566914 DOI: 10.1038/nri3608]
- 53 **Clevers HC**, Bevins CL. Paneth cells: maestros of the small intestinal crypts. *Annu Rev Physiol* 2013; **75**: 289-311 [PMID: 23398152 DOI: 10.1146/annurev-physiol-030212-183744]
- 54 **Mowat AM**, Agace WW. Regional specialization within the intestinal immune system. *Nat Rev Immunol* 2014; **14**: 667-685 [PMID: 25234148 DOI: 10.1038/nri3738]
- 55 **Cash HL**, Whitham CV, Behrendt CL, Hooper LV. Symbiotic bacteria direct expression of an intestinal bactericidal lectin. *Science* 2006; **313**: 1126-1130 [PMID: 16931762 DOI: 10.1126/science.1127119]
- 56 **Vandamme D**, Landuyt B, Luyten W, Schoofs L. A comprehensive summary of LL-37, the factotum human cathelicidin peptide. *Cell Immunol* 2012; **280**: 22-35 [PMID: 23246832 DOI: 10.1016/j.cellimm.2012.11.009]
- 57 **Kopp ZA**, Jain U, Van Limbergen J, Stadnyk AW. Do antimicrobial peptides and complement collaborate in the intestinal mucosa? *Front Immunol* 2015; **6**: 17 [PMID: 25688244 DOI: 10.3389/fimmu.2015.00017]
- 58 **Adolph TE**, Tomczak MF, Niederreiter L, Ko HJ, Böck J, Martinez-Naves E, Glickman JN, Tschurtschenthaler M, Hartwig J, Hosomi S, Flak MB, Cusick JL, Kohno K, Iwakaki T, Billmann-Born S, Raine T, Bharti R, Lucius R, Kweon MN, Marciniak SJ, Choi A, Hagen SJ, Schreiber S, Rosenstiel P, Kaser A, Blumberg RS. Paneth cells as a site of origin for intestinal inflammation. *Nature* 2013; **503**: 272-276 [PMID: 24089213 DOI: 10.1038/nature12599]
- 59 **Cadwell K**, Liu JY, Brown SL, Miyoshi H, Loh J, Lennerz JK, Kishi C, Kc W, Carrero JA, Hunt S, Stone CD, Brunt EM, Xavier RJ, Sleckman BP, Li E, Mizushima N, Stappenbeck TS, Virgin HW 4th. A key role for autophagy and the autophagy gene Atg16l1 in mouse and human intestinal Paneth cells. *Nature* 2008; **456**: 259-263 [PMID: 18849966 DOI: 10.1038/nature07416]
- 60 **Haapamäki MM**, Grönroos JM, Nurmi H, Alanen K, Nevalainen TJ. Gene expression of group II phospholipase A2 in intestine in Crohn's disease. *Am J Gastroenterol* 1999; **94**: 713-720 [PMID: 10086656 DOI: 10.1111/j.1572-0241.1999.00941.x]
- 61 **Cunliffe RN**, Mahida YR. Expression and regulation of antimicrobial peptides in the gastrointestinal tract. *J Leukoc Biol* 2004; **75**: 49-58 [PMID: 14525966 DOI: 10.1189/jlb.0503249]
- 62 **Shanahan MT**, Carroll IM, Grossniklaus E, White A, von Furstenberg RJ, Barner R, Fodor AA, Henning SJ, Sartor RB, Gulati AS. Mouse Paneth cell antimicrobial function is independent of Nod2. *Gut* 2014; **63**: 903-910 [PMID: 23512834 DOI: 10.1136/gutjnl-2012-304190]
- 63 **Machado LR**, Ottolini B. An evolutionary history of defensins: a role for copy number variation in maximizing host innate and adaptive immune responses. *Front Immunol* 2015; **6**: 115 [PMID: 25852686 DOI: 10.3389/fimmu.2015.00115]
- 64 **Cobo ER**, Chadee K. Antimicrobial Human β -Defensins in the Colon and Their Role in Infectious and Non-Infectious Diseases. *Pathogens* 2013; **2**: 177-192 [PMID: 25436887 DOI: 10.3390/pathogens2010177]
- 65 **Fahlgren A**, Hammarstrom S, Danielsson A, Hammarstrom ML. beta-Defensin-3 and -4 in intestinal epithelial cells display increased mRNA expression in ulcerative colitis. *Clin Exp Immunol* 2004; **137**: 379-385 [PMID: 15270856 DOI: 10.1111/j.1365-2249.2004.02543.x]
- 66 **Sidiq T**, Yoshihama S, Downs I, Kobayashi KS. Nod2: A Critical Regulator of Ileal Microbiota and Crohn's Disease. *Front Immunol* 2016; **7**: 367 [PMID: 27703457 DOI: 10.3389/fimmu.2016.00367]
- 67 **Wehkamp J**, Salzman NH, Porter E, Nuding S, Weichenthal M, Petras RE, Shen B, Schaeffeler E, Schwab M, Linzmeier R, Feathers RW, Chu H, Lima H Jr, Fellermann K, Ganz T, Stange EF, Bevins CL. Reduced Paneth cell alpha-defensins in ileal Crohn's disease. *Proc Natl Acad Sci USA* 2005; **102**: 18129-18134 [PMID: 16330776 DOI: 10.1073/pnas.0505256102]
- 68 **Hampe J**, Cuthbert A, Croucher PJ, Mirza MM, Mascheretti S, Fisher S, Frenzel H, King K, Hasselmeier A, MacPherson AJ, Bridger S, van Deventer S, Forbes A, Nikolaus S, Lennard-Jones JE, Foelsch UR, Krawczak M, Lewis C, Schreiber S, Mathew CG. Association between insertion mutation in NOD2 gene and Crohn's disease in

- German and British populations. *Lancet* 2001; **357**: 1925-1928 [PMID: 11425413 DOI: 10.1016/S0140-6736(00)05063-7]
- 69 **Lala S**, Ogura Y, Osborne C, Hor SY, Bromfield A, Davies S, Ogunbiyi O, Nuñez G, Keshav S. Crohn's disease and the NOD2 gene: a role for paneth cells. *Gastroenterology* 2003; **125**: 47-57 [PMID: 12851870]
 - 70 **Rosenstiel P**, Fantini M, Bräutigam K, Kühbacher T, Waetzig GH, Seegert D, Schreiber S. TNF-alpha and IFN-gamma regulate the expression of the NOD2 (CARD15) gene in human intestinal epithelial cells. *Gastroenterology* 2003; **124**: 1001-1009 [PMID: 12671897 DOI: 10.1053/gast.2003.50157]
 - 71 **Wehkamp J**, Wang G, Kübler I, Nuding S, Gregorieff A, Schnabel A, Kays RJ, Fellermann K, Burk O, Schwab M, Clevers H, Bevins CL, Stange EF. The Paneth cell alpha-defensin deficiency of ileal Crohn's disease is linked to Wnt/Tcf-4. *J Immunol* 2007; **179**: 3109-3118 [PMID: 17709525]
 - 72 **Koslowski MJ**, Kübler I, Chamaillard M, Schaeffeler E, Reinisch W, Wang G, Beisner J, Teml A, Peyrin-Biroulet L, Winter S, Herrlinger KR, Rutgeerts P, Vermeire S, Cooney R, Fellermann K, Jewell D, Bevins CL, Schwab M, Stange EF, Wehkamp J. Genetic variants of Wnt transcription factor TCF-4 (TCF7L2) putative promoter region are associated with small intestinal Crohn's disease. *PLoS One* 2009; **4**: e4496 [PMID: 19221600 DOI: 10.1371/journal.pone.0004496]
 - 73 **Hampe J**, Franke A, Rosenstiel P, Till A, Teuber M, Huse K, Albrecht M, Mayr G, De La Vega FM, Briggs J, Günther S, Prescott NJ, Onnie CM, Häslér R, Sipos B, Fölsch UR, Lengauer T, Platzer M, Mathew CG, Krawczak M, Schreiber S. A genome-wide association scan of nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in ATG16L1. *Nat Genet* 2007; **39**: 207-211 [PMID: 17200669 DOI: 10.1038/ng1954]
 - 74 **Kaser A**, Lee AH, Franke A, Glickman JN, Zeissig S, Tilg H, Nieuwenhuis EE, Higgins DE, Schreiber S, Glimcher LH, Blumberg RS. XBP1 links ER stress to intestinal inflammation and confers genetic risk for human inflammatory bowel disease. *Cell* 2008; **134**: 743-756 [PMID: 18775308 DOI: 10.1016/j.cell.2008.07.021]
 - 75 **Ménard S**, Förster V, Lotz M, Gütle D, Duerr CU, Gallo RL, Henriques-Normark B, Pütsep K, Andersson M, Glocker EO, Hornfeld MW. Developmental switch of intestinal antimicrobial peptide expression. *J Exp Med* 2008; **205**: 183-193 [PMID: 18180308 DOI: 10.1084/jem.20071022]
 - 76 **McElroy SJ**, Underwood MA, Sherman MP. Paneth cells and necrotizing enterocolitis: a novel hypothesis for disease pathogenesis. *Neonatology* 2013; **103**: 10-20 [PMID: 23006982 DOI: 10.1159/000342340]
 - 77 **Yee WH**, Soraisham AS, Shah VS, Aziz K, Yoon W, Lee SK; Canadian Neonatal Network. Incidence and timing of presentation of necrotizing enterocolitis in preterm infants. *Pediatrics* 2012; **129**: e298-e304 [PMID: 22271701 DOI: 10.1542/peds.2011-2022]
 - 78 **Zhang C**, Sherman MP, Prince LS, Bader D, Weitkamp JH, Slaughter JC, McElroy SJ. Paneth cell ablation in the presence of Klebsiella pneumoniae induces necrotizing enterocolitis (NEC)-like injury in the small intestine of immature mice. *Dis Model Mech* 2012; **5**: 522-532 [PMID: 22328592 DOI: 10.1242/dmm.009001]
 - 79 **Coutinho HB**, da Mota HC, Coutinho VB, Robalinho TI, Furtado AF, Walker E, King G, Mahida YR, Sewell HF, Wakelin D. Absence of lysozyme (muramidase) in the intestinal Paneth cells of newborn infants with necrotizing enterocolitis. *J Clin Pathol* 1998; **51**: 512-514 [PMID: 9797727]
 - 80 **Schaart MW**, de Bruijn AC, Bouwman DM, de Krijger RR, van Goudoever JB, Tibboel D, Renes IB. Epithelial functions of the residual bowel after surgery for necrotizing enterocolitis in human infants. *J Pediatr Gastroenterol Nutr* 2009; **49**: 31-41 [PMID: 19458550 DOI: 10.1097/MPG.0b013e318186d341]
 - 81 **Neu J**, Mshvildadze M, Mai V. A roadmap for understanding and preventing necrotizing enterocolitis. *Curr Gastroenterol Rep* 2008; **10**: 450-457 [PMID: 18799119]
 - 82 **Dunbar KB**, Spechler SJ. Controversies in Barrett esophagus. *Mayo Clin Proc* 2014; **89**: 973-984 [PMID: 24867396 DOI: 10.1016/j.mayocp.2014.01.022]
 - 83 **Chen W**, Frankel WL, Cronley KM, Yu L, Zhou X, Yearsley MM. Significance of paneth cell metaplasia in Barrett esophagus: a morphologic and clinicopathologic study. *Am J Clin Pathol* 2015; **143**: 665-671 [PMID: 25873500 DOI: 10.1309/AJCPVUJMCVBC9PKM]
 - 84 **Tamagawa Y**, Ishimura N, Uno G, Yuki T, Kazumori H, Ishihara S, Amano Y, Kinoshita Y. Notch signaling pathway and Cdx2 expression in the development of Barrett's esophagus. *Lab Invest* 2012; **92**: 896-909 [PMID: 22449796 DOI: 10.1038/labinvest.2012.56]
 - 85 **Krishnadath KK**, Wang KK. Molecular pathogenesis of Barrett esophagus: current evidence. *Gastroenterol Clin North Am* 2015; **44**: 233-247 [PMID: 26021192 DOI: 10.1016/j.gtc.2015.02.002]
 - 86 **Kroemer A**, Elsabbagh AM, Matsumoto CS, Zasloff M, Fishbein TM. The microbiome and its implications in intestinal transplantation. *Curr Opin Organ Transplant* 2016; **21**: 135-139 [PMID: 26761420 DOI: 10.1097/MOT.0000000000000278]
 - 87 **Fishbein T**, Novitskiy G, Mishra L, Matsumoto C, Kaufman S, Goyal S, Shetty K, Johnson L, Lu A, Wang A, Hu F, Kallakury B, Lough D, Zasloff M. NOD2-expressing bone marrow-derived cells appear to regulate epithelial innate immunity of the transplanted human small intestine. *Gut* 2008; **57**: 323-330 [PMID: 17965060 DOI: 10.1136/gut.2007.133322]
 - 88 **Eriguchi Y**, Takashima S, Oka H, Shimoji S, Nakamura K, Uryu H, Shimoda S, Iwasaki H, Shimono N, Ayabe T, Akashi K, Teshima T. Graft-versus-host disease disrupts intestinal microbial ecology by inhibiting Paneth cell production of α -defensins. *Blood* 2012; **120**: 223-231 [PMID: 22535662 DOI: 10.1182/blood-2011-12-401166]
 - 89 **Levine JE**, Huber E, Hammer ST, Harris AC, Greenon JK, Braun TM, Ferrara JL, Holler E. Low Paneth cell numbers at onset of gastrointestinal graft-versus-host disease identify patients at high risk for nonrelapse mortality. *Blood* 2013; **122**: 1505-1509 [PMID: 23760615 DOI: 10.1182/blood-2013-02-485813]
 - 90 **Worthington JJ**. The intestinal immunoendocrine axis: novel cross-talk between enteroendocrine cells and the immune system during infection and inflammatory disease. *Biochem Soc Trans* 2015; **43**: 727-733 [PMID: 26551720 DOI: 10.1042/BST20150090]
 - 91 **Nogueira AM**, Barbosa AJ. Immunocytochemical study of intestinal endocrine cells in germ-free mice. *Eur J Histochem* 1994; **38**: 213-218 [PMID: 7530514]
 - 92 **Canl PD**, Everard A, Duparc T. Gut microbiota, enteroendocrine functions and metabolism. *Curr Opin Pharmacol* 2013; **13**: 935-940 [PMID: 24075718 DOI: 10.1016/j.coph.2013.09.008]
 - 93 **Park HS**, Goodlad RA, Wright NA. Crypt fission in the small intestine and colon. A mechanism for the emergence of G6PD locus-mutated crypts after treatment with mutagens. *Am J Pathol* 1995; **147**: 1416-1427 [PMID: 7485404]
 - 94 **Graham TA**, Humphries A, Sanders T, Rodriguez-Justo M, Tadrous PJ, Preston SL, Novelli MR, Leedham SJ, McDonald SA, Wright NA. Use of methylation patterns to determine expansion of stem cell clones in human colon tissue. *Gastroenterology* 2011; **140**: 1241-1250.e1-9 [PMID: 21192938 DOI: 10.1053/j.gastro.2010.12.036]
 - 95 **Humphries A**, Cereser B, Gay LJ, Miller DS, Das B, Gutteridge A, Elia G, Nye E, Jeffery R, Poulson R, Novelli MR, Rodriguez-Justo M, McDonald SA, Wright NA, Graham TA. Lineage tracing reveals multipotent stem cells maintain human adenomas and the pattern of clonal expansion in tumor evolution. *Proc Natl Acad Sci USA* 2013; **110**: E2490-E2499 [PMID: 23766371 DOI: 10.1073/pnas.1220353110]
 - 96 **Fujimitsu Y**, Nakanishi H, Inada K, Yamachika T, Ichinose M, Fukami H, Tatematsu M. Development of aberrant crypt foci involves a fission mechanism as revealed by isolation of aberrant crypts. *Jpn J Cancer Res* 1996; **87**: 1199-1203 [PMID: 9045953]
 - 97 **Langlands AJ**, Almet AA, Appleton PL, Newton IP, Osborne JM, Näthke IS. Paneth Cell-Rich Regions Separated by a Cluster of Lgr5+ Cells Initiate Crypt Fission in the Intestinal Stem Cell Niche. *PLoS Biol* 2016; **14**: e1002491 [PMID: 27348469 DOI: 10.1371/journal.pbio.1002491]
 - 98 **Pin C**, Parker A, Gunning AP, Ohta Y, Johnson IT, Carding SR, Sato T. An individual based computational model of intestinal crypt fission and its application to predicting unrestricted growth of the intestinal epithelium. *Integr Biol (Camb)* 2015; **7**: 213-228 [PMID: 25537618 DOI: 10.1039/c4ib00236a]
 - 99 **Sato T**, Vries RG, Snippert HJ, van de Wetering M, Barker N, Stange

- DE, van Es JH, Abo A, Kujala P, Peters PJ, Clevers H. Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* 2009; **459**: 262-265 [PMID: 19329995 DOI: 10.1038/nature07935]
- 100 **Metcalf C**, Kljavin NM, Ybarra R, de Sauvage FJ. Lgr5+ stem cells are indispensable for radiation-induced intestinal regeneration. *Cell Stem Cell* 2014; **14**: 149-159 [PMID: 24332836 DOI: 10.1016/j.stem.2013.11008]
- 101 **Joo M**, Shahsafaie A, Odze RD. Paneth cell differentiation in colonic epithelial neoplasms: evidence for the role of the Apc/beta-catenin/Tcf pathway. *Hum Pathol* 2009; **40**: 872-880 [PMID: 19269007 DOI: 10.1016/j.humpath.2008.12.003]
- 102 **Farin HF**, Van Es JH, Clevers H. Redundant sources of Wnt regulate intestinal stem cells and promote formation of Paneth cells. *Gastroenterology* 2012; **143**: 1518-1529.e7 [PMID: 22922422 DOI: 10.1053/j.gastro.2012.08.031]

P- Reviewer: Hokama A, Wejman J, Xu WX **S- Editor:** Kong JX
L- Editor: A **E- Editor:** Lu YJ





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

