

[全部](#) [图片](#) [视频](#) [新闻](#) [地图](#) [图书](#)

找到约 7,030 条结果

**时间不限**

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

**所有结果**

精确匹配

[Incidence and management of gastrointestinal bleeding with ... - NCBI](#)<https://www.ncbi.nlm.nih.gov/pubmed/22541185>

BACKGROUND: Continuous flow left ventricular assist devices (CF-LVADs) have ... of care for patients in advanced heart failure (HF) requiring long-term mechanical ... CI, 0.98 -0.99; p=0.001) were independent predictors of GI hemorrhage.

[Gastrointestinal bleeding in a patient with a continuous-flow ... - NCBI](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4840201/>

28 Apr 2016 ... A 66 years old male with a history of congestive heart failure .... As Crow observed, the rate of GI bleed in CF-LVAD patients was far ... in CF-LVAD patients vs 4.6 events/100 patient years in mechanical valve recipients][5].

[Gastrointestinal Bleeding in Recipients of the HeartWare Ventricular ...](#)[www.heartfailure.onlinejacc.org/article.aspx?articleID=2194997](http://www.heartfailure.onlinejacc.org/article.aspx?articleID=2194997)

Mini-Focus Issue: Mechanical Support and Bleeding: Clinical Research | April 2015 ... Objectives This study evaluated gastrointestinal bleeding (GIB) in patients ... for recipients of continuous flow device left ventricular assist devices ( CF-LVAD). ... Mean time to first bleed was 273.1 days and 86.1% of events occurred ...

[Gastrointestinal bleed after left ventricular assist device implantation ...](#)[www.annalscts.com/article/view/4755/5645](http://www.annalscts.com/article/view/4755/5645)

8 Jun 2014 ... Gastrointestinal bleed after left ventricular assist device ... Results: The incidence of GIB after CF-LVAD varies between 18-40% in numerous studies. ... The New York Heart Association (NYHA) classifies patients with HF by their ... may require mechanical support for their failing hearts with an LVAD.

[\[PDF\] Left Ventricular Assist Devices and Gastrointestinal Bleeding: A ...](#)[onlinelibrary.wiley.com/doi/10.1002/clc.22096/pdf](http://onlinelibrary.wiley.com/doi/10.1002/clc.22096/pdf)

Hypothesis: GI bleeding occurs frequently in patients with LVADs, especially with devices with nonpulsatile .... GI hemorrhage, AVM, and heart-assist devices.

[\[PDF\] Gastrointestinal Bleeding with Continuous Flow Left - InTech](#)

Name of Journal: World Journal of Gastroenterology

Manuscript NO: 33234

Manuscript Type: EDITORIAL

## Bleeding with the artificial heart: Gastrointestinal hemorrhage in CF-LVAD patients

Grigoriy E Gurvits, Elena Fradkov

### Abstract

Continuous-flow left ventricular assist devices (CF-LVADs) have significantly improved outcomes for patients with end-stage heart failure when used as a bridge to cardiac transplantation or, more recently, as destination therapy. However, its implantations carries a risk of complications including infection, device malfunction, embolism, right ventricular failure, thromboembolic disease, reoperative and

### Match Overview

1	<b>Internet</b> 36 words crawled on 19-Aug-2016 <a href="http://f1000.com">f1000.com</a>	1%
2	<b>Crossref</b> 30 words Leslie W. Miller. "Use of a Continuous-Flow Device in Patients Awaiting Heart Transplantation", <i>New England Journal</i>	1%
3	<b>Crossref</b> 10 words Marsano, Joseph, Jay Desai, Shannon Chang, Michelle Chau, Mark Pochapin, and Grigoriy E. Gurvits. "Characteristics	<1%

Bleeding with the artificial heart: Gastrointestinal hemorrhage in CF-LVAD patie

全部 图片 视频 新闻 地图 图书

找到约 7,040 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Incidence and management of gastrointestinal bleeding with ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pubmed/22541185>

BACKGROUND: Continuous flow left ventricular assist devices (CF-LVADs) have ... of care for patients in advanced heart failure (HF) requiring long-term mechanical ... CI, 0.98 -0.99; p=0.001) were independent predictors of GI hemorrhage.

[Gastrointestinal bleeding in a patient with a continuous-flow ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4840201/>

28 Apr 2016 ... A 66 years old male with a history of congestive heart failure ... As Crow observed, the rate of GI bleed in CF-LVAD patients was far ... in CF-LVAD patients vs 4.6 events/100 patient years in mechanical valve recipients[5].

[Gastrointestinal Bleeding in Recipients of the HeartWare Ventricular ...](#)

[www.heartfailure.onlinejacc.org/article.aspx?articleID=2194997](http://www.heartfailure.onlinejacc.org/article.aspx?articleID=2194997)

Mini-Focus Issue: Mechanical Support and Bleeding: Clinical Research | April 2015 ... Objectives This study evaluated gastrointestinal bleeding (GIB) in patients ... for recipients of continuous flow device left ventricular assist devices ( CF-LVAD). ... Mean time to first bleed was 273.1 days and 86.1% of events occurred ...

[Gastrointestinal bleed after left ventricular assist device implantation ...](#)

[www.annalscts.com/article/view/4755/5645](http://www.annalscts.com/article/view/4755/5645)

8 Jun 2014 ... Gastrointestinal bleed after left ventricular assist device ... Results: The incidence of GIB after CF-LVAD varies between 18-40% in numerous studies. ... The New York Heart Association (NYHA) classifies patients with HF by their ... may require mechanical support for their failing hearts with an LVAD.

[\[PDF\] Left Ventricular Assist Devices and Gastrointestinal Bleeding: A ...](#)

[onlinelibrary.wiley.com/doi/10.1002/clc.22096/pdf](http://onlinelibrary.wiley.com/doi/10.1002/clc.22096/pdf)

Hypothesis: GI bleeding occurs frequently in patients with LVADs, especially with devices with nonpulsatile .... GI hemorrhage, AVM, and heart-assist devices.

[\[PDF\] Gastrointestinal Bleeding with Continuous Flow Left - InTech](#)

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 7,060 条结果 (用时 0.65 秒)

### Incidence and management of gastrointestinal bleeding with ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/22541185> - 翻译此页

作者: A Aggarwal - 2012 - 被引用次数: 80 - 相关文章

BACKGROUND: Continuous flow left ventricular assist devices (CF-LVADs) have ... of care for patients in advanced heart failure (HF) requiring long-term mechanical ... CI, 0.98 -0.99; p=0.001) were independent predictors of GI hemorrhage.

### Gastrointestinal bleeding in a patient with a continuous-flow ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: RV Mirasol - 2016 - 相关文章

2016年4月28日 - A 66 years old male with a history of congestive heart failure .... As Crow observed, the rate of GI bleed in CF-LVAD patients was far ... in CF-LVAD patients vs 4.6 events/100 patient years in mechanical valve recipients)[5].

### Gastrointestinal Bleeding in Recipients of the ... - JACC: Heart Failure

[www.heartfailure.onlinejacc.org/article.aspx?articleID=2194997](http://www.heartfailure.onlinejacc.org/article.aspx?articleID=2194997) - 翻译此页

Mini-Focus Issue: Mechanical Support and Bleeding: Clinical Research | April 2015 ... Objectives This study evaluated gastrointestinal bleeding (GIB) in patients ... for recipients of continuous flow device left ventricular assist devices (CF-LVAD). ... Mean time to first bleed was 273.1 days and 86.1% of events occurred ...

### Gastrointestinal bleed after left ventricular assist device implantation ...

[www.annalscts.com](http://www.annalscts.com) > Home > Vol 3, No 5 (September 2014) > 翻译此页

作者: L Harvey - 2014 - 被引用次数: 31 - 相关文章

2014年6月8日 - Gastrointestinal bleed after left ventricular assist device ... Results: The incidence of GIB after CF-LVAD varies between 18-40% in numerous studies. ... The New York Heart Association (NYHA) classifies patients with HF by their ... may require mechanical support for their failing hearts with an LVAD.

### Left Ventricular Assist Devices and Gastrointestinal Bleeding: A ...