

1-Aug-2017 06:04PM

3402 words • 7 matches • 5 sources

FAQ



33392-Review

BY KAMAL ABULEBDA

Quotes Excluded  
Bibliography Excluded4%  
SIMILARName of Journal: *World Journal of Critical Care*

Manuscript NO: 33392

Manuscript Type: ORIGINAL ARTICLE

*Retrospective Study*Intensivist-based deep sedation using propofol for pediatric outpatient flexible  
bronchoscopyKamal Abulebda, Samer Abu-Sultaneh, Sheikh Sohail Ahmed, Elizabeth A.S. Moser,  
Renee C McKinney, Riad Lutfi

## Match Overview

- | Match Number | Source   | Words    | Similarity |
|--------------|----------|----------|------------|
| 1            | Crossref | 41 words | 2%         |
| 2            | Crossref | 22 words | 1%         |
| 3            | Crossref | 17 words | 1%         |
| 4            | Crossref | 16 words | 1%         |
| 5            | Internet | 14 words | 1%         |
- 1 Sokol, Sarah, Ishaq Lat, and John Kress. "890 : Implications of the first 48 hours of sedation on duration of mechanical ventilation." *Journal of Intensive Care Medicine* 28(10): 890-895, 2013.
- 2 Tuffi, Antonio, Antonino, Alessandra Casuccio, Carmelo Buttà, Rosaria Pecoraro, Domenico Di Raimondo, Vittoriano, et al. "Propofol sedation in pediatric patients undergoing flexible bronchoscopy: a retrospective study." *Journal of Intensive Care Medicine* 28(10): 890-895, 2013.
- 3 Strayer, R.J. "Adverse events associated with ketamine for procedural sedation in adults." *American Journal of Emergency Medicine* 28(10): 890-895, 2013.
- 4 Kamal, Pradip P., Courtney E. McCracken, Scott E. Gillespie, James D. Fortenberry, Jana A. Stockwell, Joseph P. Cravens, et al. "Propofol sedation in pediatric patients undergoing flexible bronchoscopy: a retrospective study." *Journal of Intensive Care Medicine* 28(10): 890-895, 2013.
- 5 crawled on 13-Jun-2017  
[www.wjgnet.com](http://www.wjgnet.com)

891: [Propofol-based sedation: Intensivist-run procedural sedation for pediatric outpatient bronchoscopy.](#)

R Lutfi, S Hicks, R McKinney, M Nitu... - Critical Care ..., 2013 - journals.lww.com

... 3, -4 or-5 within the first 48 hours were classified as [deep sedation](#) (DS). .... Our [intensivist-based procedural sedation](#) team has assisted pulmonologists to sedate children [with flexible](#) ... can be safely [sedated](#) for [flexible fiberoptic bronchoscopy](#) via an intensivist-run [sedation](#) team in ...

[所有 2 个版本](#) [引用](#) [保存](#)

893: [A multicenter evaluation of the incidence of severe, refractory alcohol withdrawal.](#)

N Benedict, E Forsberg, B Lohr... - Critical Care ..., 2013 - journals.lww.com

... 3, -4 or-5 within the first 48 hours were classified as [deep sedation](#) (DS). .... Our [intensivist-based procedural sedation](#) team has assisted pulmonologists to sedate children [with flexible](#) ... can be safely [sedated](#) for [flexible fiberoptic bronchoscopy](#) via an intensivist-run [sedation](#) team in ...

[所有 2 个版本](#) [引用](#) [保存](#)

890: [Implications of the first 48 hours of sedation on duration of mechanical ventilation.](#)

S Sokol, I Lat, J Kress - Critical Care Medicine, 2013 - journals.lww.com

... 3, -4 or-5 within the first 48 hours were classified as [deep sedation](#) (DS). .... Our [intensivist-based procedural sedation](#) team has assisted pulmonologists to sedate children [with flexible](#) ... can be safely [sedated](#) for [flexible fiberoptic bronchoscopy](#) via an intensivist-run [sedation](#) team in ...

[所有 2 个版本](#) [引用](#) [保存](#)

892: [Correlation between bi-spectral index and sedation scales in mechanically ventilated patients.](#)

D Pradhan, P Bhattacharyya, J Mitra - Critical Care Medicine, 2013 - journals.lww.com

... 3, -4 or-5 within the first 48 hours were classified as [deep sedation](#) (DS). .... Our [intensivist-based procedural sedation](#) team has assisted pulmonologists to sedate children [with flexible](#) ... can be safely [sedated](#) for [flexible fiberoptic bronchoscopy](#) via an intensivist-run [sedation](#) team in ...

[所有 2 个版本](#) [引用](#) [保存](#)

[全部](#)[地图](#)[图片](#)[购物](#)[新闻](#)[更多](#)[设置](#)[工具](#)

找到约 5,890 条结果 (用时 0.85 秒)

[\[PDF\] Propofol-based sedation by intensivists provides an effective and sa...](#)  
[www2.pedsedation.org/.../Abulebda\\_Propofol%20based%20sedation%20by...](#) • [翻译此页](#)  
sedation for **pediatric outpatient flexible bronchoscopy**. Introduction: ... sedation team performs **deep sedation** to assist pulmonologist sedating these patients ... A total of 458 patients were sedated **using propofol only or propofol and ketamine** ...

[2016 Society for Pediatric Sedation Conference - Abstracts](#)  
[www2.pedsedation.org/sections/meetings/2016conference/.../abstracts.html](#) • [翻译此页](#)  
**Propofol-sparing deep sedation** following induction **with** dexmedetomidine: A case report. SPS - 02  
... **Propofol-based sedation by intensivists** provides an effective and safe sedation for **pediatric outpatient flexible bronchoscopy**. SPS - 13 ...

[Download PDF - Pediatrics](#)  
[https://pediatrics.aappublications.org/content/.../peds.2016-1212.full-text.pdf](#) • [翻译此页](#)  
filed conflict of interest statements **with** the American Academy ... Patients Before, During, and After Sedation for Diagnostic ... a time-**based** record that includes the name ... Complications of **flexible bronchoscopy** ... **propofol** for **deep sedation** and ... for **outpatient** dental procedures: ... S. **Propofol** sedation: **intensivists'**

[Pediatric Sedation: A Global Challenge - NCBI - NIH](#)  
[https://www.ncbi.nlm.nih.gov/](#) • [NCBI](#) • [Literature](#) • [PubMed Central \(PMC\)](#) • [翻译此页](#)

作者: D Gozal - 2010 - 被引用次数: 17 - [相关文章](#)

2010年10月19日 - **Pediatric** sedation is a challenge which spans all continents and has grown ... an oxygen mask before sedation) or for **flexible bronchoscopy** (97 children), ... The hospitalists deliver over 2,000 sedation per year, mostly ketamine **based** ... **With deep sedation**, supplemental oxygen was determined to mask ...

[Challenges in paediatric procedural sedation: political, economic, and ...](#)  
[bj.a.oxfordjournals.org/content/113/suppl\\_2/i48.full.pdf](#) • [翻译此页](#)



全部

地图

图片

新闻

视频

更多

设置

工具

找到约 8,850 条结果 (用时 0.65 秒)

**[PDF] Review of Pediatric Sedation - CiteSeerX**[citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.457.8628&rep...](https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.457.8628&rep...) ▼ 翻译此页

作者: JP Cravero - 2004 - 被引用次数: 140 - 相关文章

current guidelines **use** the terminology of minimal sedation, moderate sedation, **deep sedation**, and anes- thesia. This language is consistent **with** that used by.

**[PDF] Kamal Abulebda, Renee McKinney, Riad Lutfi Pediatric Critical Care ...**[www2.pedsedation.org/.../Abulebda\\_Propofol%20based%20sedation%20by...](http://www2.pedsedation.org/.../Abulebda_Propofol%20based%20sedation%20by...) ▼ 翻译此页

**Propofol-based** sedation by **intensivists** provides an effective and safe sedation for **pediatric outpatient flexible bronchoscopy**. Introduction: ... sedation team performs **deep sedation** to assist pulmonologist sedating these patients **using** ... performed. A total of 458 patients were sedated **using propofol** only or **propofol** and.

**The safety of propofol sedation for elective nonintubated ...**<https://www.semanticscholar.org/...propofol-sedation.../eda65e385b97dd76e...> ▼ 翻译此页

**Propofol** sedation proved safe **with** a 2.1% prevalence of minor adverse events ... safety of **deep sedation** provided by **pediatric intensivists** for elective nonintubated ... was done on all **outpatient** elective **pediatric** esophagogastroduodenoscopy ... of adverse events associated **with propofol-based** sedation/anesthesia care.

**Deep Sedation With Inhaled Sevoflurane for Pediatric Outpati ...**[mobile.journals.lww.com/.../Deep\\_Sedation\\_With\\_Inhaled\\_Sevoflurane\\_for...](http://mobile.journals.lww.com/.../Deep_Sedation_With_Inhaled_Sevoflurane_for...) - 翻译此页

**Deep Sedation With Inhaled Sevoflurane for Pediatric Outpatient** ... The intravenous drugs were administered by **pediatric intensivists** in the intensive care unit. .... Patient charges were calculated **based** on time units for room **use** in the ICU ... Patients sedated **with propofol** woke up significantly sooner than patients given ...