

Benefit of neoadjuvant concurrent chemoradiotherapy for locally advanced per

全部 图片 视频 新闻 地图 图书

找到约 3,090 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Benefit of neoadjuvant concurrent chemoradiotherapy for locally advanced perihilar cholangiocarcinoma](#)

[... neoadjuvant chemoradiotherapy for locally advanced ...](#) - Ryan - 被引用次数: 226

[Concurrent chemoradiotherapy in resected ...](#) - Nelson - 被引用次数: 87

[Chemotherapy for cholangiocarcinoma: An update](#) - Ramírez-Merino - 被引用次数: 50

[Chemotherapy for cholangiocarcinoma: An update](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3731530/>

15 Jul 2013 ... The **locally advanced cholangiocarcinoma** setting includes a ... arising from the epithelial cells of the intrahepatic, **perihilar** and extrahepatic bile ducts. The **benefit of neoadjuvant chemoradiotherapy** was also suggested in a ... prefer the use of gemcitabine plus oxaliplatin **concurrent** with radiotherapy.

[Outcomes of concurrent chemoradiotherapy versus chemotherapy ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3880040/>

21 Dec 2013 ... Although **neoadjuvant concurrent chemoradiotherapy** (CCRT) and liver ... **perihilar cholangiocarcinoma** [13,14] and **locally advanced IHCC** [15,16]. the **benefits** of CCRT in patients with advanced-stage unresectable biliary ...

[Neoadjuvant therapy for downstaging of locally advanced hilar ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3967880/>

26 Aug 2013 ... The **benefit** of an aggressive approach if a R0 resection is achieved needs to be ... The only exception is the use of **neoadjuvant chemoradiation** prior to the ... certain number of patients with **locally advanced** hilar cholangiocarcinoma, Liver transplantation for unresectable **perihilar cholangiocarcinoma**.

[Neoadjuvant treatment of biliary tract cancer: state-of-the-art and ...](#)

www.futuremedicine.com/doi/full/10.2217/hep.15.43 ▼

A proper selection of BTC patients who may **benefit** from a **neoadjuvant** strategy is ... **Neoadjuvant** therapy in **locally advanced** GBC has been mainly explored by Nine (five **perihilar** and four distal **cholangiocarcinoma**) of 51 patients with ... or gemcitabine-based **concurrent chemoradiotherapy for locally advanced** BTC ...

Match Overview

1	Internet 75 words crawled on 06-Sep-2010 radonc.duke.edu	2%
2	Internet 39 words crawled on 27-Dec-2016 www.beta.esmo.org	1%
3	Crossref 35 words Grendar, Jan, Petra Grendarova, Richie Sinha, and Eliza h Dixon. "Neoadjuvant therapy for downstaging of loca ...	1%
4	Internet 31 words crawled on 22-Jan-2016 www.wjgnet.com	1%
5	Crossref 30 words Hilar Cholangiocarcinoma, 2013.	1%
6	Internet 25 words crawled on 15-Mar-2017 www.jove.com	1%
7	Crossref 12 words &NA;.. "15th World Conference on Lung Cancer:", Journ al of Thoracic Oncology, 2013.	<1%
8	Crossref 10 words Moon, Do Chang, Hee Seung Lee, Yong Il Lee, Moon Ja e Chung, Jeong Youp Park, Seung Woo Park, Si Young	<1%
9	Internet 10 words crawled on 25-Dec-2016	<1%

⁹
Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 33402

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Benefit of neoadjuvant concurrent chemoradiotherapy for locally advanced perihi
lar cholangiocarcinoma

Jang Han Jung, Hyun Jik Lee, Hee Seung Lee, Jung Hyun Jo, In Rae Cho, ⁶Moon Jae
Chung, Jeong Youp Park, Seung Woo Park, Si Young Song, Seungmin Bang

Abstract

Benefit of neoadjuvant concurrent chemoradiotherapy for locally advanced per

全部

图片

视频

新闻

地图

图书

找到约 4,100 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Google 学术: Benefit of neoadjuvant concurrent chemoradiotherapy for locally advanced perihilar cholangiocarcinoma](#)

[... neoadjuvant chemoradiotherapy for locally advanced ...](#) - Ryan - 被引用次数: 231

[Concurrent chemoradiotherapy in resected ...](#) - Nelson - 被引用次数: 90

[Chemotherapy for cholangiocarcinoma: An update](#) - Ramirez-Merino - 被引用次数: 54

[Neoadjuvant treatment of biliary tract cancer: state-of-the-art and ...](#)

[www.futuremedicine.com/doi/full/10.2217/hep.15.43](#) ▼

A proper selection of BTC patients who may benefit from a neoadjuvant strategy is ... Neoadjuvant therapy in locally advanced GBC has been mainly explored by Nine (five perihilar and four distal cholangiocarcinoma) of 51 patients with ... or gemcitabine-based concurrent chemoradiotherapy for locally advanced BTC ...

[Outcomes of concurrent chemoradiotherapy versus chemotherapy ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3880040/>

21 Dec 2013 ... Although neoadjuvant concurrent chemoradiotherapy (CCRT) and liver ... perihilar cholangiocarcinoma [13,14] and locally advanced IHCC [15,16], ... the benefits of CCRT in patients with advanced-stage unresectable biliary ...

[Concurrent chemoradiotherapy for locally advanced breast cancer ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4324341/>

Concurrent chemoradiotherapy for locally advanced breast cancer—time for a new ... More recently, neoadjuvant therapy, particularly chemotherapy, is being ... Sadly, however, such an advantage has not been seen in most studies”, but its ...

[Chemotherapy for cholangiocarcinoma: An update - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3731530/>

15 Jul 2013 ... The locally advanced cholangiocarcinoma setting includes a ... arising from the epithelial cells of the intrahepatic, perihilar and extrahepatic bile ducts. The benefit of neoadjuvant chemoradiotherapy was also suggested in a ... prefer the use of gemcitabine plus oxaliplatin concurrent with radiotherapy.

找到约 4,350 条结果 (用时 1.00 秒)

Google 学术 : Benefit of neoadjuvant concurrent chemoradiotherapy for locally advanced perihilar cholangiocarcinoma

... neoadjuvant chemoradiotherapy for locally advanced ... - Ryan - 被引用次数 : 237

Concurrent chemoradiotherapy in resected ... - Nelson - 被引用次数 : 90

Chemotherapy for cholangiocarcinoma: An update - Ramírez-Merino - 被引用次数 : 55

Neoadjuvant treatment of biliary tract cancer: state-of-the-art and new ...

www.futuremedicine.com > Journal home > TOC > 翻译此页

作者 : M Imperatori - 2016 - 被引用次数 : 1 - 相关文章

Gall bladder cancer (GBC), and intrahepatic and extrahepatic (perihilar or distal bile ... A proper selection of BTC patients who may benefit from a neoadjuvant ... Neoadjuvant therapy in locally advanced GBC has been mainly explored by or gemcitabine-based concurrent chemoradiotherapy for locally advanced BTC ...

Chemotherapy for cholangiocarcinoma: An update - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : N Ramírez-Merino - 2013 - 被引用次数 : 52 - 相关文章

2013年7月15日 - The locally advanced cholangiocarcinoma setting includes a ... arising from the epithelial cells of the intrahepatic, perihilar and extrahepatic bile ducts. The benefit of neoadjuvant chemoradiotherapy was also suggested in a ... prefer the use of gemcitabine plus oxaliplatin concurrent with radiotherapy.

Neoadjuvant therapy for downstaging of locally advanced hilar ... - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3967880/> - 翻译此页

作者 : J Grendar - 2014 - 被引用次数 : 15 - 相关文章

2013年8月26日 - The benefit of an aggressive approach if a R0 resection is achieved needs to be ...