


World Journal of *Transplantation*

World J Transplant 2017 October 24; 7(5): 250-275





REVIEW

- 250 Peripheral blood stem cell mobilization in multiple myeloma: Growth factors or chemotherapy?
Wallis WD, Qazilbash MH

ORIGINAL ARTICLE

Retrospective Cohort Study

- 260 Prediction of delayed graft function using different scoring algorithms: A single-center experience
Michalak M, Wouters K, Fransen E, Hellemans R, Van Craenenbroeck AH, Couttenye MM, Bracke B, Ysebaert DK, Hartman V, De Greef K, Chapelle T, Roeyen G, Van Beeumen G, Emonds MP, Abramowicz D, Bosmans JL

Observational Study

- 269 Graft loss among renal-transplant recipients with early reduction of immunosuppression for BK viremia
Azar MM, Assi R, Valika AK, Banach DB, Hall IE, Landry ML, Malinis MF

Contents

World Journal of Transplantation
Volume 7 Number 5 October 24, 2017

ABOUT COVER

Editorial Board Member of *World Journal of Transplantation*, Daniel KL Cheuk, MD, Doctor, Department of Pediat and Adolescent Medical, University of Hong Kong, Hong Kong, China

AIM AND SCOPE

World Journal of Transplantation (*World J Transplant*, *WJT*, online ISSN 2220-3230, DOI: 10.5500) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJT covers topics concerning organ and tissue donation and preservation; tissue injury, repair, inflammation, and aging; immune recognition, regulation, effector mechanisms, and opportunities for induction of tolerance, thoracic transplantation (heart, lung), abdominal transplantation (kidney, liver, pancreas, islets), transplantation of tissues, cell therapy and islet transplantation, clinical transplantation, experimental transplantation, immunobiology and genomics, and xenotransplantation. The current columns of *WJT* include editorial, frontier, diagnostic advances, therapeutics advances, field of vision, mini-reviews, review, topic highlight, medical ethics, original articles, case report, clinical case conference (Clinicopathological conference), and autobiography.

AIM AND SCOPE

World Journal of Transplantation is now indexed in PubMed, PubMed Central.

FLYLEAF

I-IV Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ya-Jing Lu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Li-Jun Cui*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Transplantation

ISSN
ISSN 2220-3230 (online)

LAUNCH DATE
December 24, 2011

FREQUENCY
Bimonthly

EDITOR-IN-CHIEF
Maurizio Salvadori, MD, Professor, Renal Unit, Careggi University Hospital, Florence 50139, Italy

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/2220-3230/editorialboard.htm>

EDITORIAL OFFICE
Xiu-Xia Song, Director

World Journal of Transplantation
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
October 24, 2017

COPYRIGHT

© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS

<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION

<http://www.f6publishing.com>

Observational Study

Graft loss among renal-transplant recipients with early reduction of immunosuppression for BK viremia

Marwan M Azar, Roland Assi, Aziz K Valika, David B Banach, Isaac E Hall, Marie-Louise Landry, Maricar F Malinis

Marwan M Azar, Department of Pathology, Section of Microbiology, Massachusetts General Hospital, Boston, MA 02145, United States

Marwan M Azar, Maricar F Malinis, Section of Infectious Diseases, Yale School of Medicine, New Haven, CT 06510, United States

Roland Assi, Maricar F Malinis, Department of Surgery, Yale University School of Medicine, New Haven, CT 06510, United States

Aziz K Valika, Adventist Health Partners, Chicago, IL 60521, United States

David B Banach, Division of Infectious Diseases, University of Connecticut School of Medicine, Farmington, CT 06032, United States

Isaac E Hall, Division of Hypertension and Nephrology, University of Utah School of Medicine, Dalt Lake City, UT 84132, United States

Marie-Louise Landry, Department of Laboratory Medicine, Yale University School of Medicine, New Haven, CT 06510, United States

Author contributions: Azar MM and Malinis MF analyzed data; Azar MM drafted manuscript; Assi R, Valika AK, Banach DB, Hall IE, Landry ML and Malinis MF revised and edited manuscript; all authors contributed significantly to this work.

Institutional review board statement: The Yale University Institutional Review Board approved this study and all procedures conducted were in accordance with the Helsinki Declaration of 1975.

Conflict-of-interest statement: The authors have no conflicts of interest.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external

reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Marwan M Azar, MD, Microbiology Fellow, Department of Pathology, Section of Microbiology, Massachusetts General Hospital, 55 Fruit Street, Boston, MA 02145, United States. mmazar@mgh.harvard.edu
Telephone: +1-617-6434393
Fax: +1-888-3244639

Received: February 12, 2017

Peer-review started: February 15, 2017

First decision: April 17, 2017

Revised: June 28, 2017

Accepted: August 2, 2017

Article in press: August 2, 2017

Published online: October 24, 2017

Abstract

AIM

To review the incidence of graft loss and acute rejection among renal transplant recipients with early reduction of immunosuppression for BK viremia.

METHODS

We performed a retrospective analysis of consecutive *de-novo* kidney-only transplants from January 2009 to December 2012 to evaluate the incidence of Polyomavirus associated nephropathy (PyVAN). Recipient plasma was screened for BKV DNA *via* quantitative polymerase chain reaction (PCR) at months 1, 3, 6, 9 and 12 post-transplant and on worsening graft function.

Immunosuppression was reduced at ≥ 3 -log copies/mL. Those with viremia of ≥ 4 -log copies/mL (presumptive PyVAN) underwent renal transplant biopsy. Presumptive PyVAN (PP) and definitive PyVAN (DP; biopsy-proven) were treated by immunosuppression reduction (IR) only.

RESULTS

Among 319 kidney transplant recipients, the median age was 53 years (range 19-83), 65.8% were male, and 58.9% were white. Biopsy-proven acute rejection was found in 18.5% within 0-168 wk. Death-censored graft loss occurred in 5.3% ($n = 17$) and graft loss attributable to PyVAN was 0.6% ($n = 2$). Forty-seven patients were diagnosed with PP (14.7%) and 18 (5.6%) with DP. Graft loss among participants with PyVAN (8.5%) and those without (4.8%) was not significantly different. Deceased donor kidney transplantation (OR = 2.3, 95%CI = 1.1-4.6) and AR (OR = 2.3, 95%CI = 1.2-4.7) were associated with PyVAN in the multivariate analysis. BK viremia between 3 and 4-log copies/mL occurred in 27 patients, all of whom underwent IR. Of these, 16 (59%) never developed PyVAN while 11 (41%) developed PyVAN (4 DP, 7 PP) within a range of 11-39 wk.

CONCLUSION

Instituting an early reduction of immunosuppression, in the absence of adjunctive antivirals, is effective at preventing PyVAN and may be associated with a lower incidence of graft-loss without a reciprocal increase in the incidence of acute rejection.

Key words: BK virus; Renal transplant; Screening; PyVAN; Prevention; Graft loss

© The Author(s) 2017. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The authors describe results of a retrospective study of 319 renal transplant recipients who underwent reduction of immunosuppression for BK viremia at a BK viral of ≥ 3 -log copies/mL. Instituting early reduction of immunosuppression in the absence of adjunctive antivirals was effective in reducing the incidence of graft loss due to Polyoma-virus associated nephropathy (PyVAN) without a reciprocal increase in acute rejection. Our study also emphasizes that efforts to implement universal BK virus polymerase chain reaction assay standards recently developed by the World Health Organization are key in establishing a preventative strategy for PyVAN that is widely applicable and highly effective.

Azar MM, Assi R, Valika AK, Banach DB, Hall IE, Landry ML, Malinis MF. Graft loss among renal-transplant recipients with early reduction of immunosuppression for BK viremia. *World J Transplant* 2017; 7(5): 269-275 Available from: URL: <http://www.wjgnet.com/2220-3230/full/v7/i5/269.htm> DOI: <http://dx.doi.org/10.5500/wjt.v7.i5.269>

INTRODUCTION

BK virus (BKV) is a polyomavirus that causes widespread sub-clinical infection at a young age and subsequently establishes long-term latency in cells of the renal and urinary systems. In recipients of renal allograft transplants and allogeneic hematopoietic stem cell transplants, high-level BKV replication may lead to overt clinical disease. BK-polyoma virus associated nephropathy (PyVAN) is a major complication of kidney transplantation, occurring in 1%-10% of renal transplant recipients^[1,2]. PyVAN is directly associated with graft failure^[3,4] due to progressive interstitial nephritis and indirectly linked to allograft rejection due to immunosuppression reduction (IR), which is the cornerstone of PyVAN treatment^[5]. Guidelines currently recommend prospective screening for BKV reactivation post-transplantation, by using urine cytology to detect decoy cells or testing for high-level BK viruria and/or viremia. In the event of a sustained BK viremia of ≥ 4 -log copies/mL for more than 3 wk, a renal biopsy is recommended to confirm the diagnosis of PyVAN by demonstrating polyomavirus cytopathic changes with confirmation by immunohistochemical staining^[6]. In addition, a prompt reduction of immunosuppression is critical in an attempt to abrogate the full-fledged manifestations of the disease. Although agents with anti-BK activity such as leflunomide^[7], cidofovir^[8] and quinolones^[9] have been used^[10], randomized controlled trials proving their efficacy are lacking.

The positive predictive value of BK viremia at a cut-off of 7×10^3 copies/mL (approximately 4 log copies/mL) has been estimated at 50% to 60% for detecting proven PyVAN within 2 to 6 wk but rises to 90% when a threshold of 6-log copies/mL is implemented^[1,11,12]. A lower threshold of 3-log copies/mL may increase the sensitivity, leading to the identification of more cases, and earlier in the natural course when intervention may be more effective and graft loss more likely to be averted. In this paper, we aim to assess the incidence of PyVAN and graft loss in a single transplant center while implementing a reduction of immunosuppression at BKV loads of ≥ 3 -log BKV copies/mL.

MATERIALS AND METHODS

We performed a retrospective analysis of consecutive de-novo kidney transplants at Yale-New Haven Hospital (YNHH), who underwent screening for PyVAN screening and prevention, from January 2009 to December 2012. The Yale University Institutional Review Board approved this study and all procedures conducted were in accordance with the Helsinki Declaration of 1975. Individuals included in the study were above the age of 18 years and underwent primary kidney-only transplant. Medical records were reviewed for data on demographics,

comorbidities, and transplant parameters including type of transplant (deceased or living donor), CMV donor and recipient serostatus, induction and maintenance immunosuppression, presence of delayed graft function (DGF), biopsy-proven acute rejection (AR), graft loss and its etiology, last follow-up visit, deaths, BK viral load (copies/mL) and biopsy-proven PyVAN. Graft loss was censored for episodes of death with a functioning graft.

Presumptive PyVAN (PP) was defined as sustained BK viremia ≥ 4 -log copies/mL while definitive PyVAN (DP) required cytopathic changes on renal biopsy that were confirmed by positive BKV immunohistochemistry^[6]. Renal allograft rejection was graded in accordance with the BANFF working classification of renal allograft pathology^[13]. DGF was defined as acute renal failure requiring dialysis within 7 d of transplantation. Graft failure was defined as chronic allograft nephropathy leading to the resumption of chronic renal replacement therapy. Primary outcomes included both presumptive and definitive PyVAN while the secondary outcomes were graft survival and acute rejection.

The YNHH kidney transplant program has been active since 1967 and performs approximately 100 kidney transplantations annually. The standard maintenance immunosuppressive regimen consists of tacrolimus, mycophenolate mofetil and low dose prednisone (5-10 mg daily). The target tacrolimus trough level for the first 30 d post-transplant was 8-10 ng/mL and 5-7 ng/mL thereafter. As part of the institutional protocol for PyVAN screening and prevention, transplant recipient plasma is screened for BKV DNA *via* quantitative PCR. For the first two years of the study, an NIH-developed, real-time BKV PCR assay targeting the viral T antigen gene was used^[14]. Due to concerns about potential under-quantitation of some BKV subtypes, a multiplex real-time PCR assay developed at the University of Washington (UW) that targets both VP1 and T genes using two primer sets and three probes was implemented in December 2010^[15].

Per protocol, a serum BKV DNA viral load (VL) is obtained at months 1, 3, 6, 9 and 12 post-transplant and in case of worsening graft function. A BKV DNA VL between 3 and 3.99 log copies/mL prompted a 50% dose reduction of mycophenolate mofetil, a reduced target tacrolimus trough level of 5 ng/mL and monthly plasma BKVL until negative. Additionally, mycophenolate mofetil was discontinued and a renal biopsy was with immunohistochemical staining was performed if the serum BKV VL was above ≥ 4 -log copies/mL PP and DP were treated by reduction of immunosuppression, without adjunctive anti-viral treatment.

Statistical analysis

Statistical analysis was performed using SPSS software, version 16.0.0.0. In univariate analyses, χ^2 and Fisher's exact test (when appropriate) were used to evaluate categorical variables and Mann-Whitney *U* test was used for continuous variables. Predictors of PyVAN were identified using a multivariate logistic regression model. Only

variables with a *P*-value < 0.10 on univariate analysis were entered into a stepwise multivariate logistic regression model to identify factors independently associated with Presumptive PyVAN. All tests were double-tailed, with an assumed type 1 error risk a equal to 5%. Kaplan-Meier survival curves were plotted using GraphPad Prism version 6.03 (GraphPad Software, San Diego, CA, United States).

RESULTS

A total of 330 primary kidney transplant recipients were identified and were followed for a median time of 42 ± 14.7 mo. BK screening data on 11 patients was unavailable and thus they were excluded, leaving 319 patients available for analysis. The median age was 53 years (range 19-83), 65.8% were male, 58.9% were white, and 27.0% had diabetes mellitus. A CMV D+/R-serostatus was present in 18.2% of transplants and 54.5% of recipients underwent a deceased-donor kidney transplantation (DDKT). Induction immunosuppressive therapy consisted of basiliximab (44.8%), anti-thymocyte globulin (37.0%) or daclizumab (17.6%). Maintenance immunosuppressive therapy included both a calcineurin inhibitor and mycophenolate mofetil in 95% of cases and 95% received steroids. Biopsy-proven rejection was found in 18.5% ($n = 59$) of transplant recipients within 0-168 wk. Graft loss occurred in 5.3% ($n = 17$) and PyVAN-associated graft loss occurred in 0.6% ($n = 2$). Causes of graft loss included: AR ($n = 7$), antibody-mediated chronic rejection ($n = 2$), PyVAN ($n = 2$), CMV nephropathy ($n = 1$), hypoplastic kidney disease ($n = 1$), ureteral obstruction ($n = 1$), renal graft vein thrombosis ($n = 1$) and unknown cause ($n = 2$). Death ensued in 6.6% ($n = 21$) of the sample. A detailed list of demographics is found in Table 1.

BK viremia of ≥ 3 -log copies/mL was detected in 63 (19.7%) recipients. Of these, 47 (14.7%) were subsequently diagnosed with PP at a median time of 25 wk from initial screening. A renal biopsy was performed in 34 of these recipients and 18 (5.6% of the original sample) were confirmed to have DP. Two patients with DP progressed to graft failure and 4 developed AR within 90 d after reduction of immunosuppression.

The majority of the 319 patients included in the study (85.3%) never developed PyVAN. Time to first BK viremia was 190 d in patients with PyVAN and 235 d in those without. Graft loss occurred in 8.5% of patients with PyVAN vs. 4.8% of those without. Graft survival for 1-year, 3-year and 5-years were 99%, 95% and 92% respectively. A Kaplan-Meier curve of graft survival over time for recipients with and without PP (Figure 1) showed that survival was not significantly different between groups (logrank $P = 0.93$).

In a univariate analysis of recipients diagnosed with PP ($n = 47$) compared to recipients without PyVAN ($n = 272$), black race, DDKT and AR were significantly associated with PyVAN ($P < 0.10$). In a subsequent multivariate analysis, only DDKT (OR = 2.24; 95%CI

Table 1 Demographics and outcomes *n* (%)

Variable	All sample (<i>n</i> = 319)	PyVAN negative (<i>n</i> = 272)	PyVAN positive (<i>n</i> = 47)
Age (mean, yr)	53	51	53.1
Male	210 (65.8)	177 (65.1)	33 (70.2)
Black	86 (27.0)	67 (24.6)	19 (40.4)
Diabetes mellitus	86 (27.0)	71 (26.1)	15 (32.0)
CMV D+/R-	58 (18.2)	49 (18.0)	9 (19.1)
DDKT	174 (54.5)	140 (51.5)	34 (72.3)
Induction immunosuppression			
Thymoglobulin	118 (37.0)	97 (35.7)	21 (44.7)
Basiliximab	143 (44.8)	124 (45.6)	19 (40.4)
Daclizumab	56 (17.6)	50 (18.4)	6 (12.8)
Delayed graft function	58 (18.2)	47 (17.3)	11 (23.4)
Acute rejection	59 (18.5)	44 (16.2)	15 (32.0)
Graft loss	17 (5.3)	13 (4.8)	4 (8.5)
Death	21 (6.6)	15 (5.5)	6 (12.8)

DDKT: Deceased-donor kidney transplantation; PyVAN: Polyoma-virus associated nephropathy.




Figure 1 Kaplan-Meier Survival curve showing graft survival over time for recipients with and without presumptive polyoma-virus associated nephropathy. PyVAN: Polyoma-virus associated nephropathy.

= 1.1-4.54) and AR (OR = 2.42; 95%CI = 1.19-4.29) were significantly associated with PyVAN ($P < 0.05$). In this model, PyVAN was not associated with delayed graft function, graft loss or increased mortality. A full description covariates included in the fit-model is found in Table 2.

While the majority of patients with high-level viremia were found to have an initial BKVL above 4 log copies/mL, an initial BK viral load between 3 and 4-log copies/mL was reported in 27 transplant recipients, all of whom underwent reduction of immunosuppression, without administration of adjunctive anti-viral therapy. Of these, 16 (59%) never developed PyVAN while 11 (41%) developed PyVAN within a range of 11-39 wk. Among the 11 recipients with PyVAN, 4 were proven by renal biopsy and 7 were presumptive. Two of 27 recipients developed AR and none developed graft loss.

Since the BK PCR assay changed midway through the study, we compared the incidence of PyVAN when using the NIH assay (January 2009 to December 2010)

to the incidence when using the UW assay (January 2011 to December 2012). In a univariate chi-square analysis of all recipients with first BK viremia, 17/33 (52%) had PyVAN before the assay change versus 30/59 (51%) after the change ($P = 0.95$).

In addition, we reviewed BKVL data of patients with PP to evaluate for adherence or deviation from the set protocol for post-transplant viral load screening. Among 47 patients with PP, 16 patients underwent screening beyond the recommended interval during the study period and were found to have high-level viremia on belated screening. For 3 patients, screening was done within 10 d of recommended time point. A summary of patients in whom protocol deviation occurred is found in Table 3.

DISCUSSION

Though established guidelines recommend a reduction of immunosuppression at a sustained BK viremia of ≥ 4 -log copies/mL, studies vary significantly with regards to the implemented threshold. Cutoffs of any viremia^[16,17], of ≥ 3 -log copies/mL^[18] and of ≥ 4 -log copies/mL^[19] have been used with varying rates of PyVAN, graft loss and acute rejection. The incidence of PP in this study, using a lower threshold (≥ 3 -log copies/mL) of BK viremia for reduction of immunosuppression, was consistent with previously reported rates. In contrast to several investigations conducted in the last decade, which reported rates of graft loss of 15%-60% within 5 years of transplant, our incidence of graft loss, was 5.3%^[5,20-23]. The incidence of PyVAN-associated graft loss in this study (0.6%) was commensurate with more recently published data from 2009-2013, in which BK associated graft loss ranged from 0%-0.85%^[1,19,24-26]. Early reduction of immunosuppression in the setting of BK viremia, though potentially associated with decreased rates of graft loss due to BK nephropathy, carries the potential for increased rates of acute rejection. However, in this study, the incidence of acute rejection (18.5%) was also in keeping with previously reported rates^[19,25], suggesting that early reduction of

Table 2 Predictors of polyoma-virus associated nephropathy

Predictor	Univariate analysis			Multivariate analysis		
	PyVAN negative (<i>n</i> = 272)	PyVAN positive (<i>n</i> = 47)	<i>P</i> value	OR	CI	<i>P</i> value
Age (mean, yr)	51	53.1	0.343			
Male	177	33	0.493			
Black race	67	19	0.024 ^a	1.68	0.86-3.31	0.13
Diabetes mellitus	71	15	0.407			
CMV D+/R-	49	9	0.839			
DDKT	140	34	0.008 ^a	2.24	1.1-4.54	0.03 ^a
Thymoglobulin	97	21	0.237			
Basiliximab	124	19	0.511			
Daclizumab	50	6	0.412			
Delayed graft function	47	11	0.315			
Acute rejection	44	15	0.010 ^a	2.42	1.19-4.29	0.02 ^a
Graft loss	13	4				
Death	16	6	0.112			

^a*P* < 0.05. DDKT: Deceased-donor kidney transplantation; PyVAN: Polyoma-virus associated nephropathy.**Table 3 Protocol deviation among patients with presumptive polyoma-virus associated nephropathy¹**

Patient	Time point of protocol deviation post-transplant	Target days between serial screening	Actual days between serial screening	BKVL change (copies/mL)
1	Month 1 to month 3	60 d	100 d	ND to 1065190
2	Month 1 to month 3	60 d	138 d	ND to 17478
3	Month 9 to month 12	90 d	214 d	< 1000 to 1076120
4	Month 6 to month 9	90 d	393 d	< 1000 to 1269650
5	Month 6 to month 9	90 d	113 d	ND to 57982
6	Month 1 to month 3	60 d	134 d	ND to 392527
7	Month 3 to month 6	90 d	133 d	ND to 627218
8	Month 3 to month 6	90 d	108 d	ND to 82354
9	DOT to month 1	30 d	121 d	157939 at month 1
10	Month 3 to month 6	90 d	137 d	ND to 74389
11	Month 1 to month 3	60 d	179 d	ND to 28592
12	Month 1 to month 3	60 d	94 d	ND to 39000
13	DOT to month 1	30 d	58 d	17558 at month 1

¹Three patients with < 10 d of deviation from protocol were not included. ND: Not detectable.

immunosuppression may not necessarily increase the risk of acute rejection.

In the past decade, there has been a steady trend towards decreased rates of graft loss. This is thought to be the result of improved diagnostic tools including immunostaining and PCR, which better differentiate virus-induced nephropathy from acute rejection, as well as targeted interventions to promptly identify BK viremia and reduce immunosuppression earlier^[27]. A multitude of factors including the potency of induction and maintenance immunosuppressive regimens, demographic differences such as in age and race, frequency of BK viral load monitoring and use of adjunctive anti-virals may account for observed differences. The heterogeneity of these studies is further compounded by variation in the sensitivity, lower limit of detection of the BK virus PCR assay and most importantly a lack of equivalence of quantitation between different assays. Complicating matters is the presence of multiple viral subtypes, some of which (serotypes 3 and 4) are particularly prone to under-quantitation. In this study, we found no statistical difference in the number of patients with any viremia

when sequentially comparing two different molecular assays but results from laboratories using different assays have been shown to vary significantly, even when performed on the same sample^[15]. Since specific BK VL cutoffs are used to trigger interventions, assay variability is a critical issue and may indeed explain the variability in thresholds used across different transplant centers. A BK PCR standard that can be applied across laboratories is paramount in implementing a uniform BK viremia threshold for reduction of immunosuppression. In 2015, the World Health Organization (WHO) took steps to establish an international standard for BKV DNA nucleic acid amplification technique-based assays, using purified virions from BKV infected cell cultures^[28]. This standard, however, has not yet been widely adopted and additional *in-vitro* verification data and *in-vivo* clinical data are needed to ascertain its performance characteristics. Until then, performing serial testing on individual patients using the same assay within the same laboratory, eschewing over-interpretation of small viral load changes as biologically important and establishing center-specific viral load cutoffs to guide clinical decision making in local

patient populations will facilitate the interpretation of current BK viral load testing.

Certainly, adequate implementation of screening protocols is another critical factor in optimizing preventative strategies. In our study, a substantial number of patients with presumptive PyVAN did not adhere to the scheduled BKVL screening time-points. Strict adherence to screening protocol is likely to reduce the incidence of PyVAN by identifying viremia earlier and allowing for early interventions.

Instituting an early reduction of immunosuppression at ≥ 3 -log copies/mL, in the absence of adjunctive antivirals, was effective at preventing PyVAN in our center and may be associated with a lower incidence of graft-loss without an increased rate of acute rejection compared to published data. However, efforts to implement the WHO BK standard will be key in establishing a universal preventive strategy for PyVAN that is both highly effective and widely applicable.

COMMENTS

Background

Polyoma virus associated nephropathy (PyVAN) caused by BK virus (BKV) is a major complication occurring in 1%-10% of renal transplant recipients that is directly associated with graft loss and indirectly associated with graft rejection.

Research frontiers

Guidelines currently recommend prospective screening for BKV reactivation post-transplantation, with reduction of immunosuppression at > 4 -log copies/mL of BK virus. Additional research is needed to determine the best screening strategy.

Innovations and breakthroughs

The present study describes results of early reduction of immunosuppression (at ≥ 3 -log copies/mL) in the absence of antivirals. This strategy effective at preventing PyVAN and was associated with a lower incidence of graft-loss without a reciprocal increase in the incidence of acute rejection.

Applications

Early reduction of immunosuppression should be considered as a strategy for prospective screening for BKV reactivation post-transplantation.

Terminology

PyVAN is a disease of the kidney that results from reactivation of BK virus in the setting of immune suppression, leading to cytopathic effect on renal tubular cells and secondary inflammation.

Peer-review

The information provided by the authors adds to the existing knowledge on the subject and will prove useful to the transplant community.

REFERENCES

- Hirsch HH, Knowles W, Dickenmann M, Passweg J, Klimkait T, Mihatsch MJ, Steiger J. Prospective study of polyomavirus type BK replication and nephropathy in renal-transplant recipients. *N Engl J Med* 2002; **347**: 488-496 [PMID: 12181403 DOI: 10.1056/NEJMoa020439]
- Ramos E, Drachenberg CB, Portocarrero M, Wali R, Klassen DK, Fink JC, Farney A, Hirsch H, Papadimitriou JC, Cangro CB, Weir MR, Bartlett ST. BK virus nephropathy diagnosis and treatment: experience at the University of Maryland Renal Transplant Program. *Clin Transpl* 2002; 143-153 [PMID: 12971444]
- Hirsch HH. Polyomavirus BK nephropathy: a (re-)emerging complication in renal transplantation. *Am J Transplant* 2002; **2**: 25-30 [PMID: 12095052 DOI: 10.1034/j.1600-6143.2002.020106.x]
- Randhawa PS, Demetris AJ. Nephropathy due to polyomavirus type BK. *N Engl J Med* 2000; **342**: 1361-1363 [PMID: 10793170 DOI: 10.1056/NEJM200005043421809]
- Ramos E, Drachenberg CB, Papadimitriou JC, Hamze O, Fink JC, Klassen DK, Drachenberg RC, Wiland A, Wali R, Cangro CB, Schweitzer E, Bartlett ST, Weir MR. Clinical course of polyoma virus nephropathy in 67 renal transplant patients. *J Am Soc Nephrol* 2002; **13**: 2145-2151 [PMID: 12138148 DOI: 10.1097/01.ASN.0000023435.07320.81]
- Hirsch HH, Randhawa P; AST Infectious Diseases Community of Practice. BK polyomavirus in solid organ transplantation. *Am J Transplant* 2013; **13** Suppl 4: 179-188 [PMID: 23465010 DOI: 10.1111/ajt.12110]
- Josephson MA, Gillen D, Javaid B, Kadambi P, Meehan S, Foster P, Harland R, Thistlethwaite RJ, Garfinkel M, Atwood W, Jordan J, Sadhu M, Millis MJ, Williams J. Treatment of renal allograft polyoma BK virus infection with leflunomide. *Transplantation* 2006; **81**: 704-710 [PMID: 16534472 DOI: 10.1097/01.tp.0000181149.76113.50]
- Wu SW, Chang HR, Lian JD. The effect of low-dose cidofovir on the long-term outcome of polyomavirus-associated nephropathy in renal transplant recipients. *Nephrol Dial Transplant* 2009; **24**: 1034-1038 [PMID: 19059933 DOI: 10.1093/ndt/gfn675]
- Sharma BN, Li R, Bernhoff E, Gutteberg TJ, Rinaldo CH. Fluoroquinolones inhibit human polyomavirus BK (BKV) replication in primary human kidney cells. *Antiviral Res* 2011; **92**: 115-123 [PMID: 21798289 DOI: 10.1016/j.antiviral.2011.07.012]
- Wright AJ, Gill JS. Strategies to prevent BK virus infection in kidney transplant recipients. *Curr Opin Infect Dis* 2016; **29**: 353-358 [PMID: 27257795 DOI: 10.1097/QCO.0000000000000278]
- Bressollette-Bodin C, Coste-Burel M, Hourmant M, Sebillé V, Andre-Garnier E, Imbert-Marcille BM. A prospective longitudinal study of BK virus infection in 104 renal transplant recipients. *Am J Transplant* 2005; **5**: 1926-1933 [PMID: 15996241 DOI: 10.1111/j.1600-6143.2005.00934.x]
- Dall A, Hariharan S. BK virus nephritis after renal transplantation. *Clin J Am Soc Nephrol* 2008; **3** Suppl 2: S68-S75 [PMID: 18309005 DOI: 10.2215/CJN.02770707]
- Solez K, Axelsen RA, Benediktsson H, Burdick JF, Cohen AH, Colvin RB, Croker BP, Droz D, Dunnill MS, Halloran PF. International standardization of criteria for the histologic diagnosis of renal allograft rejection: the Banff working classification of kidney transplant pathology. *Kidney Int* 1993; **44**: 411-422 [PMID: 8377384 DOI: 10.1038/ki.1993.259]
- Ryschkewitsch C, Jensen P, Hou J, Fahle G, Fischer S, Major EO. Comparison of PCR-southern hybridization and quantitative real-time PCR for the detection of JC and BK viral nucleotide sequences in urine and cerebrospinal fluid. *J Virol Methods* 2004; **121**: 217-221 [PMID: 15381359 DOI: 10.1016/j.jviromet.2004.06.021]
- Hoffman NG, Cook L, Atienza EE, Limaye AP, Jerome KR. Marked variability of BK virus load measurement using quantitative real-time PCR among commonly used assays. *J Clin Microbiol* 2008; **46**: 2671-2680 [PMID: 18562587 DOI: 10.1128/JCM.00258-08]
- Knight RJ, Gaber LW, Patel SJ, DeVos JM, Moore LW, Gaber AO. Screening for BK viremia reduces but does not eliminate the risk of BK nephropathy: a single-center retrospective analysis. *Transplantation* 2013; **95**: 949-954 [PMID: 23545506 DOI: 10.1097/TP.0b013e31828423cd]
- Hardinger KL, Koch MJ, Bohl DJ, Storch GA, Brennan DC. BK-virus and the impact of pre-emptive immunosuppression reduction: 5-year results. *Am J Transplant* 2010; **10**: 407-415 [PMID: 20055811 DOI: 10.1111/j.1600-6143.2009.02952.x]
- Schaub S, Hirsch HH, Dickenmann M, Steiger J, Mihatsch MJ, Hopfer H, Mayr M. Reducing immunosuppression preserves allograft function in presumptive and definitive polyomavirus-associated nephropathy. *Am J Transplant* 2010; **10**: 2615-2623 [PMID: 21114642 DOI: 10.1111/j.1600-6143.2010.03310.x]

- 19 **Sood P**, Senanayake S, Sujeet K, Medipalli R, Zhu YR, Johnson CP, Hariharan S. Management and outcome of BK viremia in renal transplant recipients: a prospective single-center study. *Transplantation* 2012; **94**: 814-821 [PMID: 23018881 DOI: 10.1097/TP.0b013e31826690c6]
- 20 **Brennan DC**, Agha I, Bohl DL, Schnitzler MA, Hardinger KL, Lockwood M, Torrence S, Schuessler R, Roby T, Gaudreault-Keener M, Storch GA. Incidence of BK with tacrolimus versus cyclosporine and impact of preemptive immunosuppression reduction. *Am J Transplant* 2005; **5**: 582-594 [PMID: 15707414 DOI: 10.1111/j.1600-6143.2005.00742.x]
- 21 **Vasudev B**, Hariharan S, Hussain SA, Zhu YR, Bresnahan BA, Cohen EP. BK virus nephritis: risk factors, timing, and outcome in renal transplant recipients. *Kidney Int* 2005; **68**: 1834-1839 [PMID: 16164661 DOI: 10.1111/j.1523-1755.2005.00602.x]
- 22 **Wadei HM**, Rule AD, Lewin M, Mahale AS, Khamash HA, Schwab TR, Gloor JM, Textor SC, Fidler ME, Lager DJ, Larson TS, Stegall MD, Cosio FG, Griffin MD. Kidney transplant function and histological clearance of virus following diagnosis of polyomavirus-associated nephropathy (PVAN). *Am J Transplant* 2006; **6**: 1025-1032 [PMID: 16611340 DOI: 10.1111/j.1600-6143.2006.01296.x]
- 23 **Drachenberg CB**, Papadimitriou JC, Wali R, Nogueira J, Mendley S, Hirsch HH, Cangro CB, Klassen DK, Weir MR, Bartlett ST, Ramos E. Improved outcome of polyoma virus allograft nephropathy with early biopsy. *Transplant Proc* 2004; **36**: 758-759 [PMID: 15110653 DOI: 10.1016/j.transproceed.2004.03.040]
- 24 **Koukoulaki M**, Grispou E, Pistolas D, Balaska K, Apostolou T, Anagnostopoulou M, Tseleni-Kotsoyili A, Hadjiconstantinou V, Paniara O, Saroglou G, Legakis N, Drakopoulos S. Prospective monitoring of BK virus replication in renal transplant recipients. *Transpl Infect Dis* 2009; **11**: 1-10 [PMID: 18811631 DOI: 10.1111/j.1399-3062.2008.00342.x]
- 25 **Alm  ras C**, Vetromile F, Garrigue V, Szwarc I, Foulongne V, Mourad G. Monthly screening for BK viremia is an effective strategy to prevent BK virus nephropathy in renal transplant recipients. *Transpl Infect Dis* 2011; **13**: 101-108 [PMID: 21371220 DOI: 10.1111/j.1399-3062.2011.00619.x]
- 26 **Thakur R**, Arora S, Nada R, Minz M, Joshi K. Prospective monitoring of BK virus reactivation in renal transplant recipients in North India. *Transpl Infect Dis* 2011; **13**: 575-583 [PMID: 21507160 DOI: 10.1111/j.1399-3062.2011.00634.x]
- 27 **Hariharan S**. BK virus nephritis after renal transplantation. *Kidney Int* 2006; **69**: 655-662 [PMID: 16395271 DOI: 10.1038/sj.ki.5000040]
- 28 **Sheila Govind JH**, Clare Morris, Collaborative Study Group. Collaborative Study to establish the 1st WHO International Standard for BKV DNA for nucleic acid amplification technique (NAT)-based assays. *Biologicals* 2016; **44**: 423-433

P- Reviewer: Shrestha BM **S- Editor:** Kong JX **L- Editor:** A
E- Editor: Lu YJ





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

