

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 5,440 条结果 (用时 0.51 秒)

[Chemotherapy with enteric-coated tegafur/uracil for advanced ...](#)

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: T Ishikawa - 2008 - 被引用次数: 14 - [相关文章](#)

2008年5月14日 - Subsequently, UFT was **developed** to enhance the tumor specificity and effectiveness of ... A combination of **tegafur** and **uracil** in a molar ratio of 1:4 was However, in our studies, no patient had UFT-E-induced liver failure.

缺少字词: **rapid fibrosis**

[PLOS ONE: Adjuvant Oral Uracil-Tegafur with Leucovorin for ...](#)

journals.plos.org/plosone/article?id=10.1371/journal.pone.0162400 - [翻译此页](#)

作者: K Hasegawa - 2016 - [相关文章](#)

2016年9月2日 - Adjuvant therapy with UFT/LV effectively prolongs RFS after **hepatic** resection for ... **Tegafur** is metabolized to 5-fluorouracil in the **liver**, and **uracil** ... of **development** of CLM (synchronous [defined by disease-free interval shorter and leucovorin: a randomised crossover trial in **advanced** colorectal cancer.

缺少字词: **induced rapid**

[New imaging techniques for liver diseases - Journal of Hepatology](#)

[www.journal-of-hepatology.eu/article/S0168-8278\(14\)00773-9/pdf](http://www.journal-of-hepatology.eu/article/S0168-8278(14)00773-9/pdf) ▾ [翻译此页](#)


作者: BE Van Beers - 2015 - 被引用次数: 40 - [相关文章](#)

Newly **developed** or **advanced** methods of ultrasonography and For **liver fibrosis**, transient elastography is currently the most **tegafur/uracil** therapy, Ktrans measured with DCE-MR imaging Assessment of liver viscoelasticity by using shear waves **induced** by **Rapid** acquisition of multifrequency, multislice and.

[Tegafur/uracil - Wikipedia](#)

<https://en.wikipedia.org/wiki/Tegafur/uracil> ▾ [翻译此页](#)

Tegafur/uracil is a chemotherapy drug combination used in the treatment of cancer, primarily ... The UFT combination was **developed** in Japan during the 1980s. ... most commonly for **advanced** colorectal cancer to replace 5FU, and has a low ... lung, breast, stomach, head and neck, **liver**, gallbladder, bile



Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 33631

Manuscript Type: CASE REPORT

Tegafur-uracil-induced rapid development of advanced hepatic fibrosis

Honda S *et al.* Tegafur-uracil-induced hepatic fibrosis

Shuya Honda, Koji Sawada, Takumu Hasebe, Shunsuke Nakajima, Mikihiro Fujiya, Toshikatsu Okumura

Abstract

Tegafur-uracil has been reported to have only minor adverse effects and is associated with liver injury in 1.79% of Japanese patients. The development of tegafur-uracil-induced hepatic fibrosis with portal hypertension is rare. Here, we report a case of a 74-year-old woman with rapidly developing tegafur-uracil-induced hepatic fibrosis. The patient had no history of liver

Match Overview

1	Internet 16 words crawled on 06-Oct-2003 www.lb.nagasaki-u.ac.jp	1%
2	Crossref 11 words Cristina Rigamonti, "Primary Biliary Cirrhosis Associated ... th Systemic Sclerosis: Diagnostic and Clinical Challenges",	1%
3	Crossref 10 words Tamai, Hideyuki, Kosaku Moribata, Yoshiyuki Mori, Naoki S hingaki, Hisanobu Deguchi, Kazuki Ueda, Izumi Inoue, Tak	1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 5,440 条结果 (用时 0.34 秒)

Chemotherapy with enteric-coated tegafur/uracil for advanced ...

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: T Ishikawa - 2008 - 被引用次数: 14 - [相关文章](#)

2008年5月14日 - Subsequently, UFT was developed to enhance the tumor specificity and effectiveness of ... A combination of tegafur and uracil in a molar ratio of 1:4 was However, in our studies, no patient had UFT-E-induced liver failure.

缺少字词: rapid fibrosis

PLOS ONE: Adjuvant Oral Uracil-Tegafur with Leucovorin for ...

journals.plos.org/plosone/article?id=10.1371/journal.pone.0162400 - [翻译此页](#)

作者: K Hasegawa - 2016 - [相关文章](#)

2016年9月2日 - Adjuvant therapy with UFT/LV effectively prolongs RFS after hepatic resection for ... Tegafur is metabolized to 5-fluorouracil in the liver, and uracil ... of development of CLM (synchronous [defined by disease-free interval shorter and leucovorin: a randomised crossover trial in advanced colorectal cancer.

缺少字词: induced rapid

Tegafur/uracil - Wikipedia

<https://en.wikipedia.org/wiki/Tegafur/uracil> ▼ [翻译此页](#)

Tegafur/uracil is a chemotherapy drug combination used in the treatment of cancer, primarily ... The UFT combination was developed in Japan during the 1980s. ... most commonly for advanced colorectal cancer to replace 5FU, and has a low ... lung, breast, stomach, head and neck, liver, gallbladder, bile duct, pancreas, ...

缺少字词: induced rapid fibrosis

New imaging techniques for liver diseases - Journal of Hepatology

[www.journal-of-hepatology.eu/article/S0168-8278\(14\)00773-9/pdf](http://www.journal-of-hepatology.eu/article/S0168-8278(14)00773-9/pdf) ▼ [翻译此页](#)

作者: BE Van Beers - 2015 - 被引用次数: 40 - [相关文章](#)

Newly developed or advanced methods of ultrasonography and For liver fibrosis, transient elastography is currently the most tegafur/uracil therapy, Ktrans measured with DCE-MR imaging

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 5,340 条结果 (用时 0.48 秒)

Liver fibrosis - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: R Bataller - 2005 - [被引用次数: 3607](#) - [相关文章](#)

Advanced liver fibrosis results in cirrhosis, liver failure, and portal ... **Rapid** and slower fibrosers were identified, and genetic and environmental factors influencing ... The current effort to **develop** noninvasive markers to assess **liver fibrosis** is In alcohol-induced liver disease, alcohol alters the population of gut bacteria and ...

缺少字词: tegafur uracil

Reversal of Liver Fibrosis - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: MH Ismail - 2009 - [被引用次数: 73](#) - [相关文章](#)

Hepatic fibrosis is a scarring process associated with an increased and altered ... because of its limitations, noninvasive markers of **liver fibrosis** were **developed**. may serve as prognostic predictor of progression of NAFLD to **advanced** fibrosis. [48] Transient elastography is painless, **rapid**, and easily performed at the ...

缺少字词: tegafur uracil

Hepatic inflammation and progressive liver fibrosis in chronic liver ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: AJ Czaja - 2014 - [被引用次数: 101](#) - [相关文章](#)

2014年3月14日 - Chronic liver inflammation drives **hepatic fibrosis**, and current ... has been associated with progressive **hepatic fibrosis** and the **development** of cirrhosis[3]. They may also **induce** expression of the anti-apoptotic protein, Bcl-2, and of progression from early stage to **advanced** stage disease compared to ...

缺少字词: tegafur uracil

New imaging techniques for liver diseases - Journal of Hepatology

[www.journal-of-hepatology.eu/article/S0168-8278\(14\)00773-9/pdf](http://www.journal-of-hepatology.eu/article/S0168-8278(14)00773-9/pdf) - [翻译此页](#)

作者: BE Van Beers - 2015 - [被引用次数: 40](#) - [相关文章](#)

Newly **developed** or **advanced** methods of ultrasonography and For **liver fibrosis**, transient