

Outcome of a session of extracorporeal shock wave lithotripsy before ERCP for p

全部 图片 视频 新闻 地图 图书

找到约 15,700 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

[Extracorporeal shock wave lithotripsy for pancreatic and large ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3218149/>

21 Oct 2011 ... Extraction of **large** pancreatic and **common bile duct** (CBD) calculi has ... pancreatic and biliary calculi needed three or fewer **sessions** of **ESWL** with 5000 ... Small pancreatic duct (PD) **stones** can be extracted by the routine technique ... At our center, **ERCP**, pancreatic sphincterotomy and pancreatic ductal ...

[Endoscopic extraction of large common bile duct stones: A review ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3355239/>

16 May 2012 ... Keywords: **Large bile duct stones**, Endoscopic sphincterotomy, Papillary balloon ... intra or **extracorporeal shock wave lithotripsy** may offer adjuvant therapy in selected patients or ... The second **problem** is the size of the **stone** itself. .... for the management of difficult **bile duct stones** with a very good **outcome**.

[\[PDF\] Extracorporeal Shock Wave Lithotripsy \(ESWL\) in the ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/.../annsurg00189-0064.pdf>

in all patients; in two cases, two **ESWL sessions** were needed for **stone** disruption. ... ETAINED **STONES** after **common bile duct** exploration representa ...

[Laser lithotripsy of difficult bile duct stones: results in 60 patients ...](#)

[gut.bmj.com/content/43/6/823.full](http://gut.bmj.com/content/43/6/823.full)

Blind laser application of **giant common bile duct stones** before (A), during (B), and ... For blind laser application in the **bile duct** the standard **ERCP** catheter with a ... In 70/104 treatment **sessions** (67%) the fragmentation **result** was excellent or ... **Extracorporeal shock wave lithotripsy** (ESWL) is another treatment modality for ...

[\[DOC\] Use of intravenous cholecystokinin during extracorporeal shock ...](#)

[www.wjgnet.com/esps/DownloadFile.aspx?Type=M...doc](http://www.wjgnet.com/esps/DownloadFile.aspx?Type=M...doc)

18 Apr 2014 ... **ERCP** was performed 4 hours after a **session** of **ESWL**. ... CONCLUSION: CCK

## Match Overview

1	Internet 330 words crawled on 03-Mar-2017 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	11%
2	Crossref 112 words Manu Tandan. "Extracorporeal shock wave lithotripsy of large difficult common bile duct stones: Efficacy and an ...	4%
3	Crossref 52 words Heo, J.H.. "Endoscopic sphincterotomy plus large-balloon dilation versus endoscopic sphincterotomy for removal ...	2%
4	Internet 23 words crawled on 30-Jul-2010 <a href="http://lib.bioinfo.pl">lib.bioinfo.pl</a>	1%
5	Internet 22 words crawled on 10-Jan-2014 <a href="http://www.joplink.net">www.joplink.net</a>	1%
6	Crossref 22 words R. Den Toom. "Extracorporeal shock wave treatment of common bile duct stones: Experience with two different ...	1%
7	Internet 20 words crawled on 19-Mar-2013 <a href="http://www.turkgastro.org">www.turkgastro.org</a>	1%
8	Crossref 15 words Sauerbruch, T.. "Non-surgical management of bile duct stones refractory to routine endoscopic measures", Baill	<1%

Name of journal: *World Journal of Gastroenterology*

Manuscript NO: 33950

Manuscript Type: ORIGINAL ARTICLE

*Retrospective Study*

Outcome of a session of extracorporeal shock wave lithotripsy before endoscopic retrograde cholangiopancreatography for problematic and large common bile duct stones

Tao Tao, Ming Zhang, Qi-Jie Zhang, Liang Li, Tao Li, Xiao Zhu, Ming-Dong Li, Gui-Hua Li, Shu-Xia Sun

Abstract

AIM

[全部](#) [图片](#) [视频](#) [新闻](#) [地图](#) [图书](#)

找到约 22,000 条结果

## 时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

## 所有结果

精确匹配

[\[PDF\] extracorporeal shock wave lithotripsy for retained bile duct stones: a ...](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/.../pdf/gut00038-0128.pdf>

bile duct stones: a prospective randomised study. R Jakobs, H E ... less fragmentation sessions (1-29 v 2-82; ... after previouscholecystectomy and before ... patients because the calculi are too large, ... A major problem ... endoscopic retrograde cholangiopancreatography radiograph. ESWL. LISL ... CBD=Common bile duct.

[Extracorporeal shock wave lithotripsy for pancreatic and large ...](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3218149/>

21 Oct 2011 ... Extraction of large pancreatic and common bile duct (CBD) calculi has always .... due to biliary stricture are treated before subjecting the patient to ESWL. ... Repeat sessions are carried out on successive days until the stone ... ESWL followed by PD clearance on ERCP is therefore increasingly used in the ...

[\[PDF\] Extracorporeal Shock Wave Lithotripsy \(ESWL\) in the ... - NCBI](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/.../pdf/annsurg00189-0064.pdf>

in all patients; in two cases, two ESWL sessions were needed for stone disruption. ... ETAINED STONES after common bile duct exploration representa ...

[\[PDF\] Extracorporeal shock-wave lithotripsy and endoscopy: combined ...](#)[link.springer.com/content/pdf/10.1007/BF02653263.pdf](http://link.springer.com/content/pdf/10.1007/BF02653263.pdf)

Endoscopic treatment of bile duct stones is currently successful in 86% ... was performed in 4 cases for residual common bile duct ... pancreatography (ERCP) and associated endoscopic methods ... lithotripsy (ESWL) of problematic bile duct stones com- ... one subject in whom endoscopic stone removal failed as a result of.

[Laser lithotripsy of difficult bile duct stones: results in 60 patients ...](#)[gut.bmj.com/content/43/6/823](http://gut.bmj.com/content/43/6/823)

Introduction—Laser lithotripsy of bile duct stones has become a widely .... with a standard



[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 21,900 条结果 (用时 0.64 秒)

### extracorporeal shock wave lithotripsy for retained bile duct stones - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/.../pdf/gut00038-0128.pdf> - 翻译此页

作者：R Jakobs - 1997 - 被引用次数：72 - 相关文章

**bile duct stones**: a prospective randomised study. R Jakobs, H E ... less fragmentation **sessions** (1-29 v 2-82; ... after previous cholecystectomy and **before** ... patients because the calculi are too **large**, ... A major **problem** ... **endoscopic retrograde cholangiopancreatography** radiograph. **ESWL**. **LISL** ... **CBD**=**Common bile duct**.

### Extracorporeal Shock Wave Lithotripsy (ESWL) in the ... - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/.../pdf/annsurg00189-0064.pdf> - 翻译此页

作者：MC Taylor - 1988 - 被引用次数：20 - 相关文章

in all patients; in two cases, two **ESWL sessions** were needed for **stone** disruption. ... **ETAINED STONES** after **common bile duct** exploration representa ...

### Extracorporeal shock wave lithotripsy for pancreatic and large ...

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - 翻译此页

作者：M Tandan - 2011 - 被引用次数：49 - 相关文章

2011年10月21日 - Extraction of **large** pancreatic and **common bile duct** (CBD) calculi has always ... due to biliary stricture are treated **before** subjecting the patient to **ESWL**. ... Repeat **sessions** are carried out on successive days until the **stone** ... **ESWL** followed by PD clearance on **ERCP** is therefore increasingly used in the ...

### Extracorporeal shock-wave lithotripsy and endoscopy ... - Springer Link

<link.springer.com/content/pdf/10.1007/BF02653263.pdf> ▼ 翻译此页

作者：C Harz - 1991 - 被引用次数：17 - 相关文章

Endoscopic treatment of **bile duct stones** is currently successful in 86% ... was performed in 4 cases for residual **common bile duct** ... pancreatography (**ERCP**) and associated endoscopic methods ...