

## Match Overview

1

Internet 13 words  
crawled on 14-Sep-2010  
[www.sbm.ac.ir](http://www.sbm.ac.ir)

&lt;1%

**Name of Journal:** *World Journal of Hepatology***Manuscript NO:** 33999**Manuscript Type:** ORIGINAL ARTICLE*Retrospective study***Conjugated hyperbilirubinemia presenting in first 14 d in term neonates**

Fang Kuan Chiou, Christina Ong, Kong Boo Phua, Fares Chedid, Ajmal Kader



## Recurrence of kernicterus in term and near - term infants in Denmark

F Ebbesen - Acta paediatrica, 2000 - Wiley Online Library

... (μmol/L) Ratio conjugated/total plasma ... Pediatrics 1995; 96: 730–3 3. Rosenthal P. Direct bilirubin measurements in term newborns—a waste of time and ... American Academy of Pediatrics, provisional committee for quality improvement and subcommittee on hyperbillrubinemia. ... 被引用次数：174 相关文章 所有 5 个版本 引用 保存

## Guideline for the evaluation of cholestatic jaundice in infants: recommendations of the North American Society for Pediatric Gastroenterology, Hepatology and ...

V Moyer, DK Freese, PF Whittington... - Journal of pediatric ..., 2004 - journals.lww.com

For the primary care provider, cholestatic jaundice in infancy, defined as jaundice caused by an ele. 被引用次数：304 相关文章 所有 6 个版本 引用 保存

## Prospective surveillance study of severe hyperbilirubinaemia in the newborn in the UK and Ireland

D Manning, P Todd, M Maxwell, MJ Platt - ... Childhood-Fetal and Neonatal ..., 2007 - fn.bmj.com

... lactation, and relative immaturity of hepatic uptake and conjugation of billrubin. ... excluded some infants with encephalopathy associated with severe conjugated hyperbilirubinaemia.16 Also ... discharged early, supplemented by laboratory measurement of serum bilirubin and other ... 被引用次数：171 相关文章 所有 11 个版本 引用 保存

## Neonatal hyperbilirubinemia

PA Dennergy, DS Seidman... - New England Journal of ..., 2001 - Mass Medical Soc

... infants, unconjugated bilirubin is not readily excreted, and the ability to conjugate bilirubin is limited. ... Bilirubin then enters the liver and is modified to an excretable conjugated form that enters ... deficiency, there is an increased risk of hemolysis and impaired conjugation of bilirubin ... 被引用次数：582 相关文章 所有 11 个版本 引用 保存

## Identifying patients, on the first day of life, at high-risk of developing parenteral nutrition-associated liver disease

RD Christensen, E Henry, SE Wiedmeier... - Journal of ..., 2007 - search.proquest.com

... Parameter estimates are presented on the anti-logarithmic scale. ... Figure 2 Duration of the direct hyperbilirubinemia. ... Certainly, the incidence of late death was higher among those with the highest direct bilirubin concentrations. ... 被引用次数：133 相关文章 所有 9 个版本 引用 保存

## Biliary atresia

JL Hartley, M Davenport, DA Kelly - The Lancet, 2009 - Elsevier

... of affected infants have not been identified, although possibly all direct traces are ... jaundice have increased serum concentrations of unconjugated bilirubin, whereas conjugated bilirubin is raised in ... Caused by excretion of water-soluble bilirubin conjugates; urine might stain the ... 被引用次数：403 相关文章 所有 9 个版本 引用 保存

## Kernicterus in a full term infant

AA Penn, DR Enzmann, JS Hahn, DK Stevenson - Pediatrics, 1994 - Am Acad Pediatrics

... On arrival at Stanford, the physical exam was notable only for jaundice and ... The initial work-up did not reveal an obvious cause for the extreme indirect hyperbilirubinemia, but on the fourth hospital ... tinued after 2 days. Direct bilirubin peaked at 5.2 mg/dL at 8 days ... 被引用次数：163 相关文章 所有 6 个版本 引用 保存

## Development of bilirubin transport and metabolismin the newborn rhesus monkey

LM Gartner, K Lee, S Vaisman, D Lane, I Zarafu - The Journal of pediatrics, 1977 - Elsevier

... the results of this comparison strongly suggest that the capacity to conjugate bilirubin with glucuronic ... physiologic studies yields a mean value of 8.21 1.2 (SE) ~g bilirubin conjugated/100 gm ... in early Phase I animals, indicating that despite enhancement of conjugation during the ... 被引用次数：126 相关文章 所有 5 个版本 引用 保存

## Investigation of prolonged neonatal jaundice

S Hannam, M McDonnell, JM Rennie - Acta paediatrica, 2000 - Wiley Online Library

... atresia is likely (4). Bloods are taken for a PCV, total and conjugated bilirubin and a ... Hyperbilirubinaemia in breastfed vs formula-fed infants in the first six weeks of life: relationship to ... Kemper K, Forsyth B, McCarthy P. Jaundice, terminating breast- feeding and the vulnerable ... 被引用次数：69 相关文章 所有 4 个版本 引用 保存

## Understanding neonatal jaundice: a perspective on causation

RS Cohen, RJ Wong, DK Stevenson - Pediatrics & Neonatology, 2010 - Elsevier

... acid to form bilirubin glucuronide, also known as either “conjugated” or “direct” bilirubin. ... 7–10 days, and certainly beyond 14 days, conjugated hyperbilirubinemia must be ... Hemolysis and bilirubin conjugation in association with UDP-glucuronosyl- transferase 1A1 promoter ... 被引用次数：66 相关文章 所有 12 个版本 引用 保存



## 学术搜索

找到约 18,000 条结果 (用时0.19秒)

## 文章

**小提示：**只搜索中文(简体)结果，可在 [学术搜索设置](#) 指定搜索语言

## 我的图书馆

**Jaundice as an early diagnostic sign of urinary tract infection in infancy**[FJ García](#), [AL Nager](#) - [Pediatrics](#), 2002 - [Am Acad Pediatrics](#)

... The **hyperbilirubinemia** associated with UTIs can be unconjugated and related to hemolysis caused by ... coli 28,34 and other Gram-negative organisms, 29 or **conjugated** secondary to ... clear, but possible mechanisms include microcirculatory changes in the liver, **direct** effects from ...

被引用次数：171 [相关文章](#) [所有 4 个版本](#) [引用](#) [保存](#)

## 时间不限

2017以来

2016以来

2013以来

[自定义范围...](#)**Identifying patients, on the first day of life, at high-risk of developing parenteral nutrition-associated liver disease**[RD Christensen](#), [E Henry](#), [SE Wiedmeier](#)... - [Journal of ...](#), 2007 - [nature.com](#)

... model. Parameter estimates are **presented** on the anti-logarithmic scale. ... Duration of the **direct hyperbilirubinemia**. The ... mortality. Certainly, the incidence of late death was higher among those with the highest **direct bilirubin** concentrations. ...

被引用次数：127 [相关文章](#) [所有 11 个版本](#) [引用](#) [保存](#)

## 按相关性排序

[按日期排序](#)

## 不限语言

[中文网页](#)[简体中文网页](#)**Kernicterus in a full term infant**[AA Penn](#), [DR Enzmann](#), [JS Hahn](#), [DK Stevenson](#) - [Pediatrics](#), 1994 - [Am Acad Pediatrics](#)

... On arrival at Stanford, the physical exam was notable only for **jaundice** and ... The initial work-up did not reveal an obvious cause for the extreme indirect **hyperbilirubinemia**, but on the fourth hospital ... tinued after 2 **days**. **Direct bilirubin** peaked at 5.2 mg/dL at 8 **days** ...

被引用次数：165 [相关文章](#) [所有 5 个版本](#) [引用](#) [保存](#)☒ 包括专利☐ 包含引用**Guideline for the evaluation of cholestatic jaundice in infants: recommendations of the North American Society for Pediatric Gastroenterology, Hepatology and ...**[V Moyer](#), [DK Freese](#), [PF Whittington](#)... - [Journal of pediatric ...](#), 2004 - [journals.lww.com](#)

For the primary care provider, cholestatic jaundice in infancy, defined as jaundice caused by an ele...

被引用次数：304 [相关文章](#) [所有 6 个版本](#) [引用](#) [保存](#)☒ 创建快讯**Biliary atresia**[JL Hartley](#), [M Davenport](#), [DA Kelly](#) - [The Lancet](#), 2009 - [Elsevier](#)

... of affected **infants** have not been identified, although possibly all **direct** traces are ... **jaundice** have increased serum concentrations of unconjugated **bilirubin**, whereas **conjugated bilirubin** is raised



[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 245,000 条结果 (用时 0.55 秒)

### Neonatal Jaundice. Neonatal hyperbilirubinemia Information | Patient

<https://patient.info/doctor/neonatal-jaundice-pro> ▼ 翻译此页

Unconjugated bilirubin is metabolised by the liver to produce **conjugated bilirubin** ... Jaundice lasting for longer than **14 days in term infants** and 21 days in preterm infants: .... Any baby **presenting** with jaundice in the **first** 24 hours of life should ...

### Prolonged neonatal jaundice: When to worry and what to do

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: SM Gilmour - 2004 - 被引用次数: 19 - 相关文章

Search **term** ... **Neonatal** jaundice persisting beyond **14 days** of age is a common clinical ... clinical **presentation** is a group of **neonates** with **conjugated hyperbilirubinemia** and .... These patients typically present with a history of being well at birth and for their **first** few weeks, with jaundice appearing after the **first** week of life.

### Neonatal Cholestasis - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: AG Feldman - 被引用次数: 51 - 相关文章

2013年2月1日 - Cholestatic jaundice is a common **presenting** feature of **neonatal** .... **first 14 days** after birth, the cutoff for elevated **conjugated bilirubin** may be greater .... fibrotic component of PNAC and provide long-**term** benefit is not known.

### Unexplained neonatal jaundice as an early diagnostic sign of urinary ...

[www.sciencedirect.com/science/article/pii/S1201971212000707](http://www.sciencedirect.com/science/article/pii/S1201971212000707) ▼ 翻译此页

作者: M Shahian - 2012 - 被引用次数: 23 - 相关文章

**Hyperbilirubinemia** is one of the **presenting** signs of bacterial infection in ... Of 120 asymptomatic jaundiced **neonates** with a mean age of  $7 \pm 4$  **days**, ... About 60% of full **term** and 80% of premature **neonates** develop clinical **jaundice** in the **first** ... **direct** Coombs' test, blood typing of the **neonate** and