

World Journal of *Gastrointestinal Endoscopy*

World J Gastrointest Endosc 2017 October 16; 9(10): 499-539



MINIREVIEWS

- 499 Endoscopic ultrasound: Current roles and future directions
Friedberg SR, Lachter J
- 506 Endoscopic ultrasound elastography for solid pancreatic lesions
Chantarojanasiri T, Kongkam P

ORIGINAL ARTICLE

Retrospective Study

- 514 Oral esomeprazole vs injectable omeprazole for the prevention of hemorrhage after endoscopic submucosal dissection
Uchiyama T, Higurashi T, Kuriyama H, Kondo Y, Hata Y, Nakajima A

CASE REPORT

- 521 Two case reports of acute upper gastrointestinal bleeding from duodenal ulcers after Roux-en-Y gastric bypass surgery: Endoscopic diagnosis and therapy by single balloon or push enteroscopy after missed diagnosis by standard esophagogastroduodenoscopy
Hakim S, Reddy SRR, Batke M, Polidori G, Cappell MS
- 529 Cap polyposis refractory to *Helicobacter pylori* eradication treated with endoscopic submucosal dissection
Murata M, Sugimoto M, Ban H, Otsuka T, Nakata T, Fukuda M, Inatomi O, Bamba S, Kushima R, Andoh A
- 535 Endoscopic ultrasound-guided pancreaticogastrostomy for symptomatic pancreatic duct obstruction caused by migrated pancreatic stent
Lu L, Jin HB, Yang JF, Zhang XF

Contents

World Journal of Gastrointestinal Endoscopy
Volume 9 Number 10 October 16, 2017

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Endoscopy*, Dr. Stepan Suchanek, MD, PhD, Chief Doctor, Department of Internal Medicine, 1st Faculty of Medicine of the Charles University, Prague CZ16902, Czech Republic

AIM AND SCOPE

World Journal of Gastrointestinal Endoscopy (*World J Gastrointest Endosc*, *WJGE*, online ISSN 1948-5190, DOI: 10.4253) is a peer-reviewed open access (OA) academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJGE covers topics concerning gastroscopy, intestinal endoscopy, colonoscopy, capsule endoscopy, laparoscopy, interventional diagnosis and therapy, as well as advances in technology. Emphasis is placed on the clinical practice of treating gastrointestinal diseases with or under endoscopy.

We encourage authors to submit their manuscripts to *WJGE*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great clinical significance.

INDEXING/ABSTRACTING

World Journal of Gastrointestinal Endoscopy is now indexed in Emerging Sources Citation Index (Web of Science), PubMed, and PubMed Central.

FLYLEAF

I-III Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ya-Jing Lu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Gastrointestinal Endoscopy

ISSN
ISSN 1948-5190 (online)

LAUNCH DATE
October 15, 2009

FREQUENCY
Monthly

EDITORS-IN-CHIEF
Atsushi Imagawa, PhD, Director, Doctor, Department of Gastroenterology, Mitoyo General Hospital, Kan-onji, Kagawa 769-1695, Japan
Juan Manuel Herrerias Gutierrez, PhD, Academic Fellow, Chief Doctor, Professor, Unidad de Gestión Clínica de Aparato Digestivo, Hospital Universitario Virgen Macarena, Sevilla 41009, Spain

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com>

www.wjgnet.com/1948-5190/editorialboard.htm

EDITORIAL OFFICE
Xiu-Xia Song, Director
World Journal of Gastrointestinal Endoscopy
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.fjgpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.fjgpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
October 16, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.fjgpublishing.com>

Retrospective Study

Oral esomeprazole vs injectable omeprazole for the prevention of hemorrhage after endoscopic submucosal dissection

Takashi Uchiyama, Takuma Higurashi, Hitoshi Kuriyama, Yoshinobu Kondo, Yasuo Hata, Atsushi Nakajima

Takashi Uchiyama, Takuma Higurashi, Atsushi Nakajima, Department of Gastroenterology and Hepatology, Yokohama City University School of Medicine, Yokohama city, Kanagawa Prefecture 236-0042, Japan

Takashi Uchiyama, Hitoshi Kuriyama, Yasuo Hata, Department of Gastroenterology, Chigasaki Municipal Hospital, Chigasaki city, Kanagawa Prefecture 235-0042, Japan

Yoshinobu Kondo, Department of Endocrinology and Metabolism, Chigasaki Municipal Hospital, Chigasaki city, Kanagawa Prefecture 235-0042, Japan

Author contributions: Uchiyama T collected and analyzed the data, and drafted the manuscript; Higurashi T provided analytical oversight; Nakajima A supervised the study; Kuriyama H and Hata Y revised the manuscript for important intellectual content; Kondo Y offered the technical support; authors have read and approved the final version to be published.

Institutional review board statement: Ethics committee approval was received for this study from the ethics committee of Chigasaki Municipal Hospital.

Informed consent statement: All the treatment procedures were performed after obtaining informed consent in writing from the patients.

Conflict-of-interest statement: No conflict of interest was declared by the authors.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Takashi Uchiyama, MD, PhD, Department of Gastroenterology and Hepatology, Yokohama City University School of Medicine, 3-9 Fukuura, Kanazawaku, Yokohama city, Kanagawa 236-0004, Japan. tuchiyama-ykh@umin.ac.jp
Telephone: +81-45-7872640
Fax: +81-45-7843546

Received: March 22, 2017

Peer-review started: March 23, 2017

First decision: April 17, 2017

Revised: May 19, 2017

Accepted: July 14, 2017

Article in press: July 17, 2017

Published online: October 16, 2017

Abstract**AIM**

To evaluate the effectiveness of oral esomeprazole (EPZ) vs injectable omeprazole (OPZ) therapy to prevent hemorrhage after endoscopic submucosal dissection (ESD).

METHODS

A case-control study was conducted using a quasi-randomized analysis with propensity score matching. A total of 258 patients were enrolled in this study. Patients were treated with either oral EPZ or injectable OPZ. The endpoint was the incidence of hemorrhage after ESD.

RESULTS

Data of 71 subjects treated with oral EPZ and 172 subjects treated with injectable OPZ were analyzed. Analysis of 65 matched samples revealed no difference in the incidence of hemorrhage after ESD between the oral EPZ and injectable OPZ groups (OR = 0.89, 95%CI:

0.35-2.27, $P \geq 0.99$).

CONCLUSION

We conclude that oral EPZ therapy is a useful alternative to injectable PPI therapy for the prevention of hemorrhage after ESD.

Key words: Endoscopic submucosal dissection; Proton pump inhibitors; Hemorrhage

© The Author(s) 2017. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Proton pump inhibitors (PPIs) have been reported to be effective for suppressing hemorrhage after endoscopic submucosal dissection (ESD); however, it remains unclear whether oral PPI therapy or injectable PPI therapy is preferable. The results of the present study indicate that oral effectiveness of oral esomeprazole therapy is a useful alternative to injectable PPI therapy for the prevention of hemorrhage after ESD.

Uchiyama T, Higurashi T, Kuriyama H, Kondo Y, Hata Y, Nakajima A. Oral esomeprazole vs injectable omeprazole for the prevention of hemorrhage after endoscopic submucosal dissection. *World J Gastrointest Endosc* 2017; 9(10): 514-520 Available from: URL: <http://www.wjgnet.com/1948-5190/full/v9/i10/514.htm> DOI: <http://dx.doi.org/10.4253/wjge.v9.i10.514>

INTRODUCTION

Endoscopic submucosal dissection (ESD) allows *en-bloc* resection of even large and ulcerated gastric tumors^[1,2]. It enables accurate histopathological diagnosis and reduces the risk of local recurrence^[3], and is a standard treatment for selected gastric tumors. However, ESD is technically difficult and is associated with a higher risk of adverse events than conventional endoscopic mucosal resection (EMR)^[3-5]. Among the adverse events, hemorrhage is a frequently encountered and serious problem^[6]. Hemorrhage after ESD can occur at a later stage than other complications of ESD, such as perforation, sometimes occurring even after hospital discharge. Furthermore, hemorrhage after gastric ESD can be serious, as it can be massive and complicated by life-threatening hemorrhagic shock^[7]. Thus, the importance of preventing hemorrhage after ESD cannot be overemphasized. While some previous studies have reported the risk factors for hemorrhage after ESD^[6-14], no consensus has been arrived at yet in respect of the risk factors. Proton pump inhibitors (PPIs) have been reported to be effective for controlling hemorrhage after ESD^[15]. However, to the best of our knowledge, there have been no studies yet to compare the efficacy of oral PPI therapy vs injectable PPI therapy for the control of hemorrhage after ESD. It remains unclear whether oral PPI therapy or injectable PPI therapy is preferable for the

prevention of hemorrhage after ESD.

Esomeprazole (EPZ) is the S-isomer of omeprazole (OPZ) and has more favorable pharmacokinetic and pharmacodynamic profiles than OPZ^[16]. However, injectable EPZ is not available at present in our hospital. In the current study, therefore, we compared the efficacy of oral EPZ therapy with that of injectable OPZ (in place of EPZ) therapy for the prevention of hemorrhage after ESD by propensity score-matched analysis.

MATERIALS AND METHODS

Patients and methods

We conducted a retrospective study with propensity score-matched analysis. We registered patients who had undergone ESD for gastric tumors at our hospital between March 2008 and March 2014 ($n = 258$). The research protocol was approved by the Hospital Ethics Committee. Written informed consent was obtained from each of the participants of the study.

Treatment

Figure 1 shows the treatment protocol used. The patients received either oral EPZ (20 mg daily) for 8 wk after ESD (oral EPZ group) or injectable OPZ (20 mg twice daily) for the first 5 d, followed by oral OPZ (20 mg daily) from day 6 to the end of 8 wk after the ESD (injectable OPZ group). Additionally, all the patients underwent an endoscopic examination on day 2 and a third endoscopy on day 6 after the ESD. All patients were given sucralfate from day 2 to the end of 8 wk after the ESD. Antiplatelet/anticoagulant drugs were discontinued before the ESD.

ESD procedure

ESD was performed using a videoendoscope (GIF-Q260J), Electric scalpel for endoscopic surgery (IT-Knife2) (Olympus Corporation, Tokyo, Japan) and an electrosurgical unit (ICC 200) (ERBE Elektromedizin GmbH, Tübingen, Germany). After tumor resection, all the visible vessels in the created ulcer were coagulated using a coagulation device (Coagrasper) (Olympus Corporation).

Hemorrhage

Hemorrhage after ESD was defined as the presence of clinical evidence of hemorrhage, such as the occurrence of melena or hematemesis confirmed by the hospital staff, or confirmation of the presence of blood or bleeding spots in the post-ESD ulcer at the second or third endoscopy. Preventive hemostasis for visible vessels not showing evidence of hemorrhage during the second or third endoscopy was not included as evidence of hemorrhage after ESD. We also defined clinically significant hemorrhage after ESD as hemorrhage necessitating emergency endoscopy or blood transfusion.

Statistical analysis

Data are presented as mean \pm SD or number, and the diagnostic outcomes were examined using the χ^2 test.




Figure 1 Treatment protocol for the two groups. EGD: Esophagogastrroduodenoscopy; ESD: Endoscopic submucosal dissection; EPZ: Esomeprazole; OPZ: Omeprazole.

The variables and incidence of hemorrhage after ESD in the oral EPZ group were compared with those in the injectable OPZ group using the χ^2 test. Furthermore, propensity score matching was performed to control and reduce the selective bias^[17-19]. Ten variables that could potentially influence the incidence of hemorrhage after ESD, listed below, were used to generate propensity scores using logistic regression: Patient age, patient sex, history of use of antiplatelet/anticoagulant drugs, location of the lesion, lesion depth, presence/absence of ulceration, diameter of the lesion, duration of operation, macroscopic type of the lesion, and the operator experience (beginners: Surgeons who had performed < 50 gastric ESDs; experts: Surgeons who had performed > 50 gastric ESDs). A propensity score-matched cohort was created by trying to match each patient given oral EPZ with a patient given injectable OPZ (a 1:1 match), using the nearest pair method. After matching, a coarse comparison of the matched cohorts was performed using χ^2 test. *P* values of < 0.05 were considered to indicate significance. Statistical analyses were performed using the JMP 10.0 software (SAS, North Carolina, United States).

RESULTS

Background characteristics of the patients

A total of 258 patients were enrolled in this study. Fifteen following reasons: Failure of resection of the lesion *en-bloc* (*n* = 8); presence of perforation (*n* = 5); interruption of the ESD (*n* = 1); previous history of gastrectomy (*n* = 1). Data of the remaining 243 patients were evaluated. Of the 243 patients, 172 who had undergone ESD before November 2012 received injectable OPZ, and the remaining 71 patients who had received ESD after November 2012 received oral EPZ (Figure 2). The data of 71 patients of the oral EPZ group and 172 patients of the

injectable OPZ group were analyzed.

Among the 243 patients included in the analysis, 33 developed hemorrhage after the ESD (13.6%), with the hemorrhage being clinically significant in 10 of these cases (4.1%). Table 1 shows the baseline characteristics of the study population. A univariate analysis identified operator experience as the only risk factor for hemorrhage after ESD (beginner vs OR = 2.16; 95%CI: 1.03-4.55, *P* = 0.039). Hemorrhage after ESD was observed within 6 d of the procedure in all the cases.

Propensity score matching

A quasi-randomized experiment can be created using propensity score matching. That is, two subjects assigned to each group are equally likely to receive oral EPZ or injectable OPZ (Tables 2 and 3). Nearest-neighbor matches were performed using a caliper with 0.25 standard deviation of the propensity score (log odds scale). The predictive performance of the treatment model was evaluated using the χ^2 statistic that can take values from 0.5 for chance prediction to 1.0 for perfect prediction^[20]. The propensity score allowed clear distinction between cases with and without hemorrhage after ESD, with a *c*-statistic of 0.77.

Among the matched samples, the incidence of hemorrhage after ESD was 15.4% (10/65) in the oral EPZ group and 16.9% (11/65) in the injectable OPZ group, with no statistically significant difference seen between the two groups (EPZ group vs OPZ group, OR = 0.89, 95%CI: 0.35-2.27, *P* ≥ 0.99). The incidence of clinically significant hemorrhage was 6.2% (4/65) in the oral EPZ group and 4.6% (3/65) in the injectable OPZ group, with no significant difference of this parameter between the two groups either (EPZ group vs OPZ group, OR = 1.36, 95%CI: 0.29-6.31, *P* ≥ 0.99). No significant differences in any of the other variables examined were found between the two groups.




Figure 2 Selection of cases for study. ESD: Endoscopic submucosal dissection; EPZ: Esomeprazole; OPZ: Omeprazole.

Table 1 Background characteristics of the patients in relation to the bleeding status

	Hemorrhage after ESD		P value
	Negative	Positive	
Number	210	33	
Age (yr)	73.9 ± 8.0	71.7 ± 9.4	0.14
Gender (M/F)	151/59	8/25	0.65
Antiplatelet/anticoagulant drugs (-/+)	161/49	8/25	0.91
Hypertension (-/+)	123/87	20/13	0.83
Diabetes mellitus (-/+)	186/24	4/27	0.52
<i>H. pylori</i> infection (-/+)	90/120	13/20	0.89
Location (U/M/L)	28/96/86	2/15/16	0.45
Depth (m/sm)	190/20	5/28	0.35
Ulcer (-/+)	205/5	33/0	0.37
Diameter of the lesion (mm)	40.7 ± 15.3	40.4 ± 13.9	0.89
Duration of the operation (min)	106.9 ± 64.1	123.8 ± 98.5	0.2
Macroscopic type of the lesion (elevated or flat/combined/depressed)	130/29/51	5/10/2018	0.49
Pathological findings (Adenoma/Differentiated ca/Undifferentiated ca)	76/126/8	12/21/0	0.32
Lymphatic/venous invasion (-/+)	206/4	Jan-32	0.67
Anastomosis (-/+)	200/10	33/0	0.2
Operator type (expert/beginner)	129/81	14/19	0.0392
Bleeding during ESD (good/poor control)	164/46	9/24	0.49

P value calculated by the χ^2 test (Fisher's exact test or Pearson's test). ESD: Endoscopic submucosal dissection; *H. pylori*: *Helicobacter pylori*.

DISCUSSION

EPZ is the first optical isomer developed as a PPI. Previous studies have demonstrated the efficacy of oral EPZ in the treatment of GERD^[21,22]. Recently, Bunno *et al.*^[23] reported that oral EPZ was effective for ulcer healing after ESD. However, no studies have assessed the efficacy of oral EPZ for the control of hemorrhage after ESD. Recent studies have reported that oral EPZ therapy is a useful alternative to injectable PPI therapy to prevent recurrent hemorrhage in hemorrhagic gastric ulcer patients^[24,25]. Laine *et al.*^[26], who compared oral and injectable lansoprazole, showed a difference in the intragastric pH only during the first hour after PPI administration, with no difference in the intragastric

pH seen between the two groups at ≥ 1.5 h after the drug administration. Javid *et al.*^[27] demonstrated an equivalent ability of injectable and high oral doses of various PPIs in suppressing gastric acid secretion, and no significant difference in effect among various PPIs given through different routes on the gastric pH ≥ 6 for 72 h after successful endoscopic hemostasis. Our results were consistent with the findings of these previous studies. Oral EPZ therapy has the advantages of a lower cost and easier administration as compared to injectable PPI therapy, whereas injectable PPIs will still be needed for patients who cannot receive oral medications. Therefore, we conclude that oral EPZ therapy is a useful alternative to intravenous PPI therapy for the prevention of hemorrhage after ESD.

Table 2 Background characteristics of the patients according to the treatment group before the propensity score matching

	EPZ group	OPZ group	P value
Number	71	172	
Age (yr)	75.3 ± 7.1	72.9 ± 8.5	0.0361
Gender (M/F)	52/19	124/48	0.86
Antiplatelet/anticoagulant drugs (-/+)	50/21	136/36	0.15
Hypertension (-/+)	40/31	103/69	0.61
Diabetes Mellitus (-/+)	Sep-62	151/21	0.92
<i>H. pylori</i> infection (-/+)	29/42	55/117	0.19
Location (U/M/L)	10/34/27	20/77/75	0.7
Depth (m/sm)	57/14	161/11	0.0019
Ulcer (-/+)	71/0	167/5	0.15
Diameter of the lesion (mm)	42.2 ± 17.2	40.14 ± 14.1	0.31
Duration of the operation (min)	115.9 ± 87.3	106.4 ± 61.1	0.34
Macroscopic type of the lesion (elevated or flat/combined/depressed)	46/16/9	102/18/52	0.003
Pathological findings (Adenoma/Differentiated ca/Undifferentiated ca)	26/42/3	62/105/5	0.86
Lymphatic/venous invasion (-/+)	Jan-70	168/4	0.65
Anastomosis (-/+)	Mar-68	163/9	0.74
Operator type (expert/beginner)	31/40	112/60	0.002
Bleeding during ESD (good/poor control)	55/16	134/38	0.94
Hemorrhage after ESD (-/+)	149/23	Oct-61	0.88
Clinically significant bleeding (-/+)	165/7	Mar-68	0.96

P value calculated by the χ^2 test (Fisher's exact test or Pearson's test). EPZ: Esomeprazole; OPZ: Omeprazole; ESD: Endoscopic submucosal dissection.

Table 3 Background characteristics of the patients according to the treatment group after the propensity score matching

	EPZ group	OPZ group	P value
Number	65	65	
Age (yr)	75.2 ± 7.3	75.0 ± 7.5	0.9
Gender (M/F)	46/19	44/21	0.85
Antiplatelet/anticoagulant drugs (-/+)	46/19	44/21	0.85
Hypertension (-/+)	37/28	33/32	0.6
Diabetes Mellitus (-/+)	Aug-57	Dec-53	0.47
<i>H. pylori</i> infection (-/+)	26/39	23/42	0.72
Location (U/M/L)	5/33/27	10/24/31	0.19
Depth (m/sm)	Sep-56	Jun-59	0.58
Ulcer (-/+)	65/0	65/0	0
Diameter of the lesion (mm)	41.9 ± 18.5	39.6 ± 12.1	0.4
Duration of the operation (min)	110.6 ± 85.7	102.0 ± 54.9	0.5
Macroscopic type of the lesion (elevated or flat/combined/depressed)	42/14/9	10/8/1947	0.6
Pathological findings (Adenoma/Differentiated ca/Undifferentiated ca)	26/37/2	29/35/1	0.76
Lymphatic/venous invasion (-/+)	Jan-64	Feb-63	≥ 0.99
Anastomosis (-/+)	Mar-62	Mar-62	≥ 0.99
Operator type (expert/beginner)	29/36	27/38	0.86
Bleeding during ESD (good/poor control)	51/14	49/16	0.84
Hemorrhage after ESD (-/+)	Oct-55	Nov-54	≥ 0.99
Clinically significant bleeding (-/+)	Apr-61	Mar-62	≥ 0.99

P value calculated by the χ^2 test (Fisher's exact test or Pearson's test). EPZ: Esomeprazole; OPZ: Omeprazole; ESD: Endoscopic submucosal dissection.

Previous studies have reported the incidence and risk factors for hemorrhage after ESD, although the results are conflicting. We also assessed the risk factors for hemorrhage after ESD, and our analysis identified only the operator experience as a significant predictor of hemorrhage after ESD. Adequate coagulation of the vessels at the ulcer base after ESD is important to prevent delayed hemorrhage. As some experience is required for such coagulation, the incidence of hemorrhage after ESD differs between beginners and experts. A previous study also identified the operator experience as a significant risk factor for hemorrhage

after ESD^[11]. On the other hand, several studies have reported the absence of any significant effect of the operator experience on the risk of hemorrhage after ESD^[6-10,12-14]. These differences in the outcomes were likely caused by the diversity of the ESD procedures and treatments employed. Further investigation is required to clarify the unified risk factors for hemorrhage after ESD.

Our study had some limitations. First of all, injectable EPZ is not available at our hospital; therefore, we compared oral esomeprazole with injectable OPZ. Second, we could not carry out a non-inferiority study, because the number of cases was small. Third, hemorrhage after

ESD is usually defined as bleeding, including hematemesis or melena, that necessitates endoscopic treatment and has been reported to occur in 1.3% to 11.9% of patients undergoing ESD^[28]. We observed only 10 cases (10/243) with hemorrhage after ESD fulfilling this conventional definition, which made a reasonable comparison between oral EPZ and injectable OPZ difficult. Therefore, in this study, we defined hemorrhage after ESD as described in the text above. This was the reason why the frequency of hemorrhage after ESD was relatively high in this study, while the frequency of clinically significant hemorrhage was comparable to that reported from other studies. Fourth, this study was a 6-year clinical study. During this period, ESD has gradually become more and more popular. Individual learning curves, introduction of new devices, and establishment of education programs over the last few years make reasonable comparisons difficult.

In conclusion, in the present study, we assessed the frequency of hemorrhage after ESD after oral EPZ and injectable OPZ treatments. No difference was seen in the incidence of hemorrhage after ESD between the oral EPZ and injectable OPZ groups. This is the first study to investigate the effectiveness of oral EPZ therapy to prevent hemorrhage after ESD. Further large-scale trials are necessary to clarify the effectiveness of oral EPZ therapy. Oral PPI therapy shows a clear cost benefit and is easier to administer as compared to injectable PPI therapy. Thus, we conclude that oral EPZ therapy is a useful alternative to intravenous PPI therapy for the prevention of hemorrhage after ESD.

COMMENTS

Background

Endoscopic submucosal dissection (ESD) is technically difficult and is associated with a high risk of adverse events. Among the adverse events, hemorrhage is a frequently encountered and serious problem. Proton pump inhibitors (PPIs) have been reported to be effective for controlling hemorrhage after ESD. In this study, the authors compared the efficacy of oral effectiveness of oral esomeprazole (EPZ) therapy with that of injectable omeprazole (OPZ) therapy for the prevention of hemorrhage after ESD.

Research frontiers

No comparison of the efficacy of oral PPI therapy vs injectable PPI therapy for the control of hemorrhage after ESD has been carried out previously. Therefore, whether oral PPI therapy or injectable PPI therapy is preferable for the prevention of hemorrhage after ESD remains uncertain. The results of this study contributed to clarifying the efficacy of oral EPZ therapy vs injectable OPZ therapy for the prevention of hemorrhage after ESD.

Innovations and breakthroughs

A quasi-randomized experiment was created using propensity score matching. Among the matched samples, the incidence of hemorrhage after ESD was 15.4% (10/65) in the oral EPZ group and 16.9% (11/65) in the injectable OPZ group. No statistically significant difference was seen between these groups.

Applications

This study suggests that oral EPZ therapy is a useful alternative to intravenous PPI therapy for the prevention of hemorrhage after ESD. Oral PPI therapy shows a clear cost benefit and is easier to administer as compared to injectable PPI therapy.

Terminology

ESD: An endoscopic technique allows *en-bloc* resection even for large or ulcerated gastric tumors.

Peer-review

The study is very well described and the results are clear. In this study, the authors investigated to compare EPZ vs intravenous omeprazole therapy to prevent hemorrhage after ESD using a quasi-randomized analysis with propensity score matching.

REFERENCES

- 1 Ono H, Kondo H, Gotoda T, Shirao K, Yamaguchi H, Saito D, Hosokawa K, Shimoda T, Yoshida S. Endoscopic mucosal resection for treatment of early gastric cancer. *Gut* 2001; **48**: 225-229 [PMID: 11156645 DOI: 10.1136/gut.48.2.225]
- 2 Gotoda T, Yamamoto H, Soetikno RM. Endoscopic submucosal dissection of early gastric cancer. *J Gastroenterol* 2006; **41**: 929-942 [PMID: 17096062 DOI: 10.1007/s00535-006-1954-3]
- 3 Oda I, Saito D, Tada M, Ishi H, Tanabe S, Oyama T, Doi T, Otani Y, Fujisaki J, Ajioka Y, Hamada T, Inoue H, Gotoda T, Yoshida S. A multicenter retrospective study of endoscopic resection for early gastric cancer. *Gastric Cancer* 2006; **9**: 262-270 [PMID: 17235627 DOI: 10.1007/s10120-006-0389-0]
- 4 Oka S, Tanaka S, Kaneko I, Mouri R, Hirata M, Kawamura T, Yoshihara M, Chayama K. Advantage of endoscopic submucosal dissection compared with EMR for early gastric cancer. *Gastrointest Endosc* 2006; **64**: 877-883 [PMID: 17140890 DOI: 10.1016/j.gie.2006.03.932]
- 5 Fujishiro M. Endoscopic submucosal dissection for gastric cancer. *Curr Treat Options Gastroenterol* 2008; **11**: 119-124 [PMID: 18321439]
- 6 Mannen K, Tsunada S, Hara M, Yamaguchi K, Sakata Y, Fujise T, Noda T, Shimoda R, Sakata H, Ogata S, Iwakiri R, Fujimoto K. Risk factors for complications of endoscopic submucosal dissection in gastric tumors: analysis of 478 lesions. *J Gastroenterol* 2010; **45**: 30-36 [PMID: 19760133 DOI: 10.1007/s00535-009-0137-4]
- 7 Takizawa K, Oda I, Gotoda T, Yokoi C, Matsuda T, Saito Y, Saito D, Ono H. Routine coagulation of visible vessels may prevent delayed bleeding after endoscopic submucosal dissection--an analysis of risk factors. *Endoscopy* 2008; **40**: 179-183 [PMID: 18322872 DOI: 10.1055/s-2007-995530]
- 8 Okada K, Yamamoto Y, Kasuga A, Omae M, Kubota M, Hirasawa T, Ishiyama A, Chino A, Tsuchida T, Fujisaki J, Nakajima A, Hoshino E, Igarashi M. Risk factors for delayed bleeding after endoscopic submucosal dissection for gastric neoplasm. *Surg Endosc* 2011; **25**: 98-107 [PMID: 20549245 DOI: 10.1007/s00464-010-1137-4]
- 9 Jang JS, Choi SR, Graham DY, Kwon HC, Kim MC, Jeong JS, Won JJ, Han SY, Noh MH, Lee JH, Lee SW, Baek YH, Kim MJ, Jeong DS, Kim SK. Risk factors for immediate and delayed bleeding associated with endoscopic submucosal dissection of gastric neoplastic lesions. *Scand J Gastroenterol* 2009; **44**: 1370-1376 [PMID: 19891589 DOI: 10.3109/00365520903194609]
- 10 Jeon SW, Jung MK, Cho CM, Tak WY, Kweon YO, Kim SK, Choi YH. Predictors of immediate bleeding during endoscopic submucosal dissection in gastric lesions. *Surg Endosc* 2009; **23**: 1974-1979 [PMID: 18553202 DOI: 10.1007/s00464-009-0464-9]
- 11 Tsuji Y, Ohata K, Ito T, Chiba H, Ohya T, Gunji T, Matsushashi N. Risk factors for bleeding after endoscopic submucosal dissection for gastric lesions. *World J Gastroenterol* 2010; **16**: 2913-2917 [PMID: 20556838 DOI: 10.3748/wjg.v16.i23.2913]
- 12 Toyokawa T, Inaba T, Omote S, Okamoto A, Miyasaka R, Watanabe K, Izumikawa K, Horii J, Fujita I, Ishikawa S, Morikawa T, Murakami T, Tomoda J. Risk factors for perforation and delayed bleeding associated with endoscopic submucosal dissection for early gastric neoplasms: analysis of 1123 lesions. *J Gastroenterol Hepatol* 2012; **27**: 907-912 [PMID: 22142449 DOI: 10.1111/j.1440-1746.2011.07039.x]
- 13 Nakamura M, Nishikawa J, Hamabe K, Nishimura J, Satake M, Goto A, Kiyotoki S, Saito M, Fukagawa Y, Shirai Y, Okamoto T, Sakaida I. Risk factors for delayed bleeding from endoscopic submucosal

- dissection of gastric neoplasms. *Scand J Gastroenterol* 2012; **47**: 1108-1114 [PMID: 22783937 DOI: 10.3109/00365521.2012.699550]
- 14 **Higashiyama M**, Oka S, Tanaka S, Sanomura Y, Imagawa H, Shishido T, Yoshida S, Chayama K. Risk factors for bleeding after endoscopic submucosal dissection of gastric epithelial neoplasm. *Dig Endosc* 2011; **23**: 290-295 [PMID: 21951088 DOI: 10.1111/j.1443-1661.2011.01151.x]
- 15 **Uedo N**, Takeuchi Y, Yamada T, Ishihara R, Ogiyama H, Yamamoto S, Kato M, Tatsumi K, Masuda E, Tamai C, Yamamoto S, Higashino K, Iishi H, Tatsuta M. Effect of a proton pump inhibitor or an H2-receptor antagonist on prevention of bleeding from ulcer after endoscopic submucosal dissection of early gastric cancer: a prospective randomized controlled trial. *Am J Gastroenterol* 2007; **102**: 1610-1616 [PMID: 17403076 DOI: 10.1111/j.1572-0241.2007.01197.x]
- 16 **McKeage K**, Blick SK, Croxtall JD, Lyseng-Williamson KA, Keating GM. Esomeprazole: a review of its use in the management of gastric acid-related diseases in adults. *Drugs* 2008; **68**: 1571-1607 [PMID: 18627213]
- 17 **Rosenbaum PR**, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika* 1983; **70**: 41-55
- 18 **Rosenbaum PR**, Rubin DB. Reducing bias in observational studies using subclassification on the propensity score. *J Am Stat Assoc* 1984; **79**: 516-545
- 19 **Rosenbaum PR**, Rubin DB. The bias due to incomplete matching. *Biometrics* 1985; **41**: 103-116 [PMID: 4005368]
- 20 **Ash AS**, Shwartz M. Evaluating the performance of risk-adjustment methods: Dichotomous outcomes. In: Risk Adjustment for Measuring Health Care Outcomes. 2nd edition, edited by Iezzoni LI, Chicago, Health Administration Press, 1997
- 21 **Johnson DA**, Benjamin SB, Vakil NB, Goldstein JL, Lamet M, Whipple J, Damico D, Hamelin B. Esomeprazole once daily for 6 months is effective therapy for maintaining healed erosive esophagitis and for controlling gastroesophageal reflux disease symptoms: a randomized, double-blind, placebo-controlled study of efficacy and safety. *Am J Gastroenterol* 2001; **96**: 27-34 [PMID: 11197282 DOI: 10.1111/j.1572-0241.2001.03443.x]
- 22 **Richter JE**, Kahrilas PJ, Johanson J, Maton P, Breiter JR, Hwang C, Marino V, Hamelin B, Levine JG; Esomeprazole Study Investigators. Efficacy and safety of esomeprazole compared with omeprazole in GERD patients with erosive esophagitis: a randomized controlled trial. *Am J Gastroenterol* 2001; **96**: 656-665 [PMID: 11280530 DOI: 10.1111/j.1572-0241.2001.3600_b.x]
- 23 **Bunno M**, Gouda K, Yamahara K, Kawaguchi M. A Case-Control Study of Esomeprazole Plus Rebamipide vs. Omeprazole Plus Rebamipide on Post-ESD Gastric Ulcers. *Jpn Clin Med* 2013; **4**: 7-13 [PMID: 23966811 DOI: 10.4137/JCM.S11320]
- 24 **Sung JJ**, Suen BY, Wu JC, Lau JY, Ching JY, Lee VW, Chiu PW, Tsoi KK, Chan FK. Effects of intravenous and oral esomeprazole in the prevention of recurrent bleeding from peptic ulcers after endoscopic therapy. *Am J Gastroenterol* 2014; **109**: 1005-1010 [PMID: 24777150 DOI: 10.1038/ajg.2014.105]
- 25 **Yen HH**, Yang CW, Su WW, Soon MS, Wu SS, Lin HJ. Oral versus intravenous proton pump inhibitors in preventing re-bleeding for patients with peptic ulcer bleeding after successful endoscopic therapy. *BMC Gastroenterol* 2012; **12**: 66 [PMID: 22681960 DOI: 10.1186/1471-230X-12-66]
- 26 **Laine L**, Shah A, Bemanian S. Intra-gastric pH with oral vs intravenous bolus plus infusion proton-pump inhibitor therapy in patients with bleeding ulcers. *Gastroenterology* 2008; **134**: 1836-1841 [PMID: 18423628 DOI: 10.1053/j.gastro.2008.03.006]
- 27 **Javid G**, Zargar SA, U-Saif R, Khan BA, Yatoo GN, Shah AH, Gulzar GM, Sodhi JS, Khan MA. Comparison of p.o. or i.v. proton pump inhibitors on 72-h intra-gastric pH in bleeding peptic ulcer. *J Gastroenterol Hepatol* 2009; **24**: 1236-1243 [PMID: 19682194 DOI: 10.1111/j.1440-1746.2009.05900.x]
- 28 **Park CH**, Lee SK. Preventing and controlling bleeding in gastric endoscopic submucosal dissection. *Clin Endosc* 2013; **46**: 456-462 [PMID: 24143302 DOI: 10.5946/ce.2013.46.5.456]

P- Reviewer: Shu X, Suchanek S, Sugimoto M **S- Editor:** Kong JX
L- Editor: A **E- Editor:** Lu YJ





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

