

Name of Journal: *World Journal of Virology*

Manuscript NO: 34045

Manuscript Type: Field of Vision

Additional attention to combination antiretroviral therapy-related lipodystrophy

Norihiko Kobayashi, Masako Nakahara, Masako Oka, Kumiko Saeki

Abstract

The occurrence of lipodystrophy in patients taking anti-human immunodeficiency virus (HIV) medications is a serious problem as it is irreversible even after drug withdrawal. Although it was first recognized in patients taking proteinase inhibitors, other types of anti-HIV agents can also cause lipodystrophy. In a recent publication by Jones *et al* entitled "Highly active antiretroviral therapy (HAART) dysregulates proliferation and differentiation of human pre-adipocytes" in *World Journal of Virology*, it was reported that simultaneous treatment of human subcutaneous adipocytes with

Match Overview

1 **Crossref** 12 words **1%**
Shibata, Yukie, Yoshihisa Yamashita, Kanji Tsuru, Kazuhiko Ishihara, Kyoko Fukazawa, and Kunio Ishikawa

[全部](#)[图片](#)[购物](#)[新闻](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 21,500 条结果 (用时 0.77 秒)

[Response to ZDV/ddI Containing Combination Antiretroviral Therapy ...](#)

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: H Bussmann - 2009 - [被引用次数: 52](#) - [相关文章](#)

In recent years, **combination antiretroviral therapy (cART)** has been introduced ... below); an AIDS-related malignancy **other** than mucocutaneous Kaposi's sarcoma; ... **Additional** treatment steps, defined as protease-inhibitor (PI)-containing ... every two months, **lipodystrophy** and performance assessments every six months, ...

[Hypertriglyceridemia, Metabolic Syndrome, and Cardiovascular ...](#)

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: JPH van Wijk - 2012 - [被引用次数: 54](#) - [相关文章](#)

2011年8月22日 - The use of **combination antiretroviral therapy (cART)** in HIV-infected patients has ... In **addition**, HIV itself, as well as specific antiretroviral agents, may further increase Similarly, the presence of **CART-associated lipodystrophy** in **attention** to potential drug interactions with antiretroviral agents and ...

[Assessment of ultrasound for use in detecting lipodystrophy in HIV ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/19133752> - [翻译此页](#)

作者: K Viskovic - 2009 - [被引用次数: 14](#) - [相关文章](#)

... **lipodystrophy** in HIV-infected patients taking **combination antiretroviral therapy**. ... for at least 1 year with **combination antiretroviral therapy (cART)** in Zagreb, Croatia. ... **more** frequently found in patients with a history of stavudine treatment and in ... Inhibitors/therapeutic use; HIV-1/drug effects; HIV-Associated **Lipodystrophy** ...

缺少字词: attention

Scholar

About 256 results (0.06 sec)

Articles

[Adverse effects of antiretroviral therapy](#)[A Carr](#), DA Cooper - *The Lancet*, 2000 - Elsevier

Case law

... Several factors have **combined** to increase our **attention** on the toxicity of HAART. ... factor, including

My library

a history of drug hypersensitivity, has been identified that predicts the development of **antiretroviral**

Any time

hypersensitivity, including **more** severe reactions. ... **Other** features, Rechallenge. ...

Since 2017

Cited by 1197 Related articles All 14 versions Cite Save

Since 2016

Since 2013

Custom range...

[\[PDF\] Fat distribution and metabolic abnormalities in HIV-infected patients on first combination antiretroviral therapy including stavudine or zidovudine: role of physical ...](#)[P Domingo](#), MA Sambeat, A Pérez, [J Ordoñez](#)... - *Antiviral ...*, 2003 - researchgate.net

Sort by relevance

... In **addition**, since WHR was considered the main measure to define lipohypertrophy, it is possible ...

Sort by date

384 indicate that patients treated with stavudine and didanosine lose fat **more** rapidly than ... and lamivudine [9]. It is obvious that the specific responsibility of **other** components of the ... include patents

Cited by 63 Related articles All 10 versions Cite Save More

 include citations[Increased mitochondrial toxicity with ribavirin in HIV/HCV coinfection](#)[A Lafeuillade](#), G Hittinger, S Chadapaud - *The Lancet*, 2001 - Elsevier Create alert... coinfecting with HIV and hepatitis C virus who received interferon- α plus ribavirin in **addition** to HAART ... lactate concentration decreased within 2 weeks to 3.2 mmol/L. In the **other** 13 HIV ... Thetwo patients presented here were stable clinically for **more** than 3 years with HAART and ...

Cited by 388 Related articles All 9 versions Cite Save

[The case for more cautious, patient-focused antiretroviral therapy](#)[K Henry](#) - *Annals of internal medicine*, 2000 - Am Coll Physicians

... Author Info. Search: Advanced Search. < PREV ARTICLE; This Issue; NEXT ARTICLE >

Ideas and Opinions |15 February 2000. The Case for **More** Cautious, Patient-Focused**Antiretroviral** Therapy. Keith Henry, MD. Keith Henry, MD.

Cited by 81 Related articles All 7 versions Cite Save

[Hyperlipidaemia in patients with HIV-1 infection receiving highly active antiretroviral therapy: epidemiology, pathogenesis, clinical course and management](#)[L Calza](#), R Manfredi, F Chiodo - *International journal of antimicrobial agents*, 2003 - Elsevier



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 120,000 条结果 (用时 0.60 秒)

Google 学术: Additional attention to combination antiretroviral therapy-related lipodystrophy

Adverse effects of antiretroviral therapy - Carr - 被引用次数: 1197

... HIV-infected patients on first combination antiretroviral ... - Domingo - 被引用次数: 63

Increased mitochondrial toxicity with ribavirin in HIV/ ... - Lafeuillade - 被引用次数: 388

Development of Buffalo Hump in the course of antiretroviral therapy ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: G Ceccarelli - 2011 - 被引用次数: 5 - 相关文章

2011年2月17日 - For these reasons this combination of antiretroviral therapy based on RAL plus ATV attracts the attention of the scientific ... Some reports indicate that Buffalo Hump is associated with other physical features of the lipodystrophy ...

Lipodystrophy in HIV patients: its challenges and management ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: R Singhanian - 2011 - 被引用次数: 22 - 相关文章

跳到Therapeutic options - In addition, the effect sizes of the various treatments vary considerably. ... Other studies demonstrated that the discontinuation of HAART was associated with adverse ... of different antiretroviral combinations in the management of ... side effects, lipoatrophy has received the most attention.

Human Immunodeficiency Virus and Highly Active Antiretroviral ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: E Anuurad - 2009 - 被引用次数: 56 - 相关文章

Furthermore, in addition to side effects from treatment of HIV, there is a ... As the HIV-positive population ages, the combination of cardiovascular risk factors ... were the first class of antiretroviral drugs associated with lipodystrophy when the initial ... and low bone mineral density (BMD) have not