

Match Overview

1	Internet 77 words crawled on 03-Dec-2014 www.jimmunol.org	2%
2	Crossref 72 words Imaeda, H., K. Takahashi, T. Fujimoto, E. Kasumi, H. Ban, S. Bamba, H. Sonoda, T. Shimizu, Y. Fujiyama, and A. An	2%
3	Internet 57 words crawled on 23-May-2014 gut.bmj.com	2%
4	Crossref 53 words T. Sugihara. "The increased mucosal mRNA expressions of complement C3 and interleukin-17 in inflammatory b ...	1%
5	Crossref 37 words Takahashi, Kenichiro, Atsushi Nishida, Makoto Shioya, Hirotsugu Imaeda, Shigeki Bamba, Osamu Inatomi, Tomoh	1%
6	Crossref 35 words J. Dambacher. "The role of the novel Th17 cytokine IL-26 in intestinal inflammation", Gut, 5/15/2008.	1%
7	Crossref 27 words Sara Tengvall, Karlhans Fru Che, Anders Lindén. "Interleukin-26: An Emerging Player in Host Defense and Inflam	1%
8	Internet 26 words crawled on 17-Jun-2017 journal.frontiersin.org	1%
9	Internet 26 words crawled on 09-Apr-2016	1%

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 34197

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Expression of Interleukin-26 is upregulated in inflammatory bowel disease

Makoto Fujii, Atsushi Nishida, Hirotsugu Imaeda, Masashi Ohno, Kyohei Nishino, Shigeki Sakai, Osamu Inatomi, Shigeki Bamba, Masahiro Kawahara, Tomoharu Shimizu, Akira Andoh

Abstract

AIM

To investigate IL-26 expression in the inflamed mucosa of patients with

Expression of Interleukin-26 is upregulated in inflammatory bowel disease

全部 图片 视频 新闻 地图 图书

找到约 5,830,000 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

Google 学术: Expression of Interleukin-26 is upregulated in inflammatory bowel disease

... the novel Th17 cytokine IL-26 in intestinal inflammation - Dambacher - 被引用次数: 153

... Th17 cells in the pathogenesis of Crohn's disease - Brand - 被引用次数: 475

Development, cytokine profile and function of human ... - Wilson - 被引用次数: 1725

Intestinal expression of the anti-inflammatory interleukin-1 ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/24732025>

2014 Aug;59(2):e18-26. doi: 10.1097/MPG.0000000000000387. Intestinal expression of the anti-inflammatory interleukin-1 homologue IL-37 in ... in the intestine of pediatric patients with chronic inflammatory bowel disease (IBD). ... mRNA expression of IL-8, IL-17, and IL-10 was upregulated in patients with CD and UC.

Role of cytokines in inflammatory bowel disease - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2731177/>

21 Jul 2008 ... TNF- α expression in human macrophages was discovered in the colonic tissue and ... More specifically, sTNF-RI is up-regulated in the serum of IBD patients ... Interferon-gamma and TNF-alpha potently induced TRAIL in IEC and ... blocking the recruitment of leukocytes and increasing T-cell apoptosis[26].

Interleukins Involved in Inflammatory Bowel Disease as New ...

https://www.researchgate.net/.../271542495_Interleukins_Involved_in_Inflammatory_Bowel_Disease_as_New_Therapeutic_Targets

The Inflammatory Bowel Disease (IBD) comprises two forms namely, Ulcerative Colitis (UC) ... inflammation in IBD. Qiyang W et al. (2011). Andoh A et al. (2009). IL-26 ... expression and its expression is up-regulated in active Crohn's disease .

Epithelial expression of interleukin-37b in inflammatory bowel disease

<onlinelibrary.wiley.com/doi/10.1111/cei.12061/full>

18 Apr 2013 ... In the inflamed mucosa of IBD patients, epithelial IL-37b expression was ... and can be up-regulated by inflammatory stimuli and cytokines [6]. ... inhibitors: p42/ 44 MAPK inhibitors (PD98059 and U0216) [25, 26], a p38 MAPK ...

Expression of Interleukin-26 is upregulated in inflammatory bowel disease

全部 图片 视频 新闻 地图 图书

找到约 11,200,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Expression of Interleukin-26 is upregulated in inflammatory bowel disease](#)

[... the novel Th17 cytokine IL-26 in intestinal inflammation](#) - Dambacher - 被引用次数: 164

[... Th17 cells in the pathogenesis of Crohn's disease](#) - Brand - 被引用次数: 492

[Development, cytokine profile and function of human ...](#) - Wilson - 被引用次数: 1791

[Increased expression of interleukin 17 in inflammatory bowel disease](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1773503/>

Background and aim: **Interleukin** (IL) 17 is a cytokine which exerts strong ... It is likely that IL-17 **expression** in IBD may be associated with altered immune and ... IL-17 gene were constructed according to published sequence data: 5'-AGA GAT ...

[Intestinal expression of the anti-inflammatory interleukin-1 ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pubmed/24732025>

2014 Aug;59(2):e18-26. doi: 10.1097/MPG.0000000000000387. Intestinal **expression** of the anti-inflammatory **interleukin**-1 homologue IL-37 in ... in the intestine of pediatric patients with chronic **inflammatory bowel disease** (IBD). ... mRNA **expression** of IL-8, IL-17, and IL-10 was **upregulated** in patients with CD and UC.

[The Role of the Novel Th17 Cytokine IL-26 in Intestinal Inflammation ...](#)

<https://www.ncbi.nlm.nih.gov/labs/articles/18483078/>

15 May 2008 ... Background and aims: **Interleukin** 26 (IL-26), a novel IL-10-like ... and its **expression** is **upregulated** in active Crohn's disease, indicating a role ...

[Cytokines in inflammatory bowel disease : Nature Reviews ...](#)

www.nature.com/nri/journal/v14/n5/abs/nri3661.html

22 Apr 2014 ... Loss of **interleukin**-10 signaling and infantile **inflammatory bowel disease**: ... is up-regulated in Crohn's disease; **expression** and localization in intestinal mucosal cells. The increased **expression** of IL-23 in **inflammatory bowel disease** The role of the novel Th17 cytokine IL-26 in intestinal inflammation.

[Interleukins Involved in Inflammatory Bowel Disease as New](#)

Expression of Interleukin-26 is upregulated in inflammatory bowel disease

全部 图片 视频 新闻 地图 图书

找到约 13,900,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

Google 学术: [Expression of Interleukin-26 is upregulated in inflammatory bowel disease](#)

... [the novel Th17 cytokine IL-26 in intestinal inflammation](#) - Dambacher - 被引用次数: 164

... [Th17 cells in the pathogenesis of Crohn's disease](#) - Brand - 被引用次数: 492

[Development, cytokine profile and function of human ...](#) - Wilson - 被引用次数: 1795

[The Role of the Novel Th17 Cytokine IL-26 in Intestinal Inflammation ...](#)

<https://www.ncbi.nlm.nih.gov/labs/articles/18483078/>

15 May 2008 ... Background and aims: **Interleukin 26** (IL-26), a novel IL-10-like ... and its expression is **upregulated** in active **Crohn's disease**, indicating a role ...

[Increased expression of interleukin 17 in inflammatory bowel disease](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1773503/>

Background and aim: **Interleukin** (IL) 17 is a cytokine which exerts strong ... It is likely that IL-17 **expression** in IBD may be associated with altered immune and ... IL-17 gene were constructed according to published sequence data: 5'-AGA GAT ...

[Intestinal expression of the anti-inflammatory interleukin-1 ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pubmed/24732025>

2014 Aug;59(2):e18-26. doi: 10.1097/MPG.0000000000000387. Intestinal **expression** of the anti-inflammatory **interleukin-1** homologue IL-37 in ... in the intestine of pediatric patients with chronic **inflammatory bowel disease** (IBD). ... mRNA **expression** of IL-8, IL-17, and IL-10 was **upregulated** in patients with CD and UC.

[Cytokines in inflammatory bowel disease : Nature Reviews ...](#)

www.nature.com/nri/journal/v14/n5/abs/nri3661.html

22 Apr 2014 ... Loss of **interleukin-10** signaling and infantile **inflammatory bowel disease**: ... is **up-regulated** in **Crohn's disease**: **expression** and localization in intestinal mucosal cells. ...