

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 35028

Manuscript Type: ORIGINAL ARTICLE

Integrated MELD model improves ¹short-term prognosis of HBV-related
acute-on-chronic liver failure

Wei Chen, Jia You, Jing Chen, Qi Zheng, Jia-Ji Jiang, Yue-Yong Zhu

Abstract

AIM

To investigate whether the short-term prognosis of HBV-related acute-on-chronic liver failure (ACLF) could be improved by using an integrated MELD model including serum lactate.


METHODS

This clinical study was conducted on the First Affiliated Hospital, Fujian Medicine University, China. From 2009 to 2015, 236 patients diagnosed with

Match Overview

- | Match Number | Source | Words Matched | Similarity |
|--------------|--|---------------|------------|
| 1 | Crossref 54 words
"Abstracts of the 26th Annual Conference of APASL, February 15–19, 2017, Shanghai, China", Hepatology International | 54 | 2% |
| 2 | Crossref 34 words
"Abstracts—APASL 2013", Hepatology International, 2013. | 34 | 1% |

Risk factors for partial and full work disability in Dutch patients with inflammato



全部 图片 视频 新闻 地图 图书

找到约 11,000 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

显示的是以下查询字词的结果: [Risk factors for partial and full work disability in Dutch patients with inflammatory bowel disease](#) [An integrated MELD model including serum lactate improves the short-term prognosis of HBV-related acute-on-chronic liver failure](#)

仍然搜索: [Risk factors for partial and full work disability in Dutch patients with inflammatory bowel disease](#) [An integrated MELD model including serum lactate improves the short-term prognosis of HBV-related acute-on-chronic liver failure](#)

[Static and dynamic prognostic factors for hepatitis-B-related acute ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4612284/>

30 Sep 2015 ... [Hepatitis-B-related acute-on-chronic liver failure](#) has a poor ... factors including the [Model for End-Stage Liver Disease \(MELD\)](#) scores ... Identification of [prognostic](#) factors for [ACLF-HBV patients](#) is critical, ... The [term](#) ascites, in all definitions, refers to grade 2 or 3 ... [Risk factors](#) for an unfavorable [prognosis](#).

[AST to Platelet Ratio Index Predicts Mortality in Hospitalized ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4782890/>

7 Mar 2016 ... Search [term](#) ... A total of 193 hospitalized [chronic HBV-infected patients](#) (cirrhosis , ... and the [model](#) for end-stage liver [disease](#) score (odds ratio: 1.194, ... Liver-related [mortality](#) is largely determined by [liver failure](#), which can ... Baseline Characteristics and [Factors](#) Associated With APRI in Cirrhotic [Patients](#).

[Serum C-reactive protein predicts early mortality in hospitalized ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5287978/>

27 Jan 2017 ... Search [term](#) ... The [serum](#) C-reactive protein (CRP) is an [inflammatory](#) marker. ... in hospitalized [patients](#) with [HBV-related](#) decompensated [cirrhosis](#) (HBV-DeCi). ... In addition, the [model](#) for end-stage liver [disease \(MELD\)](#) score and ... were independent [risk factors](#) for 30-day [mortality](#) in HBV-DeCi [patients](#).

[\[PDF\] HEP_v60_iS2_Rev1 FINAL.pdf - AASLD](#)

https://www.aasld.org/sites/default/.../HEP_v60_iS2_Rev1%20FINAL_0.pdf

5 Feb 2015 ... who have discontinued anti-hepatitis B therapy, including entecavir. ... used to

Integrated model for end-stage liver disease model improves short-term progn



全部 图片 视频 新闻 地图 图书

找到约 582,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[A new multiparameter integrated MELD model for prognosis of HBV ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5400346/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5400346/>

26 Aug 2016 ... Hepatitis B virus related acute-on-chronic liver failure (HBV-ACLF) is one ... model for end-stage liver disease (MELD) assesses well the short-term ... In recent years, a number of improved models based on MELD, like serum ...

[A comprehensive validation of HBV-related acute-on-chronic liver ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5025883/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5025883/>

16 Sep 2016 ... Caused by the acute exacerbation of chronic hepatitis B (CHB), hepatitis B virus-related acute-on-chronic liver failure (HBV-ACLF) is a severe ... patients have a poor prognosis with short-term mortality ranging from 30% to 70%. MELD: model of end-stage liver disease; MELD-Na: sodium MELD; MESO: ...

[Potent antiviral therapy improves survival in acute on chronic liver ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4239490/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4239490/>

21 Nov 2014 ... ACLF secondary to reactivation of chronic hepatitis B virus is a distinct ... Keywords: Acute on chronic liver failure, Chronic hepatitis B Spontaneous or treatment-related inflammatory flares are seen in patients with chronic HBV. and model for end-stage liver disease scores for predicting short term ...

[\[PDF\] Different models in predicting the short-term prognosis of patients ...](http://annalsofhepatology.com.mx/PDF/vol11n3/HP123-05-Different.pdf)

annalsofhepatology.com.mx/PDF/vol11n3/HP123-05-Different.pdf

hepatitis B virus-related acute-on-chronic liver failure ... King's College Hospital (KCH) criteria, model for end-stage liver disease (MELD), MELD combined.

[Change in model for end-stage liver disease score at two weeks, as ...](https://academic.oup.com/.../Change-in-model-for-end-stage-liver-disease-score)

<https://academic.oup.com/.../Change-in-model-for-end-stage-liver-disease-score>

11 Nov 2014 ... Background: Acute-on-chronic liver failure (ACLF) is characterised by a ... Absolute 'model for end-stage liver disease' (MELD) score and We used a Kaplan-Meier survival model to demonstrate that a MELD score that improves at 2 short-term prognosis of hepatitis B virus-related acute-on-chronic ...

[Population-representative Incidence of Acute-On-Chronic Liver Failure](#)



全部

图片

新闻

购物

地图

更多

设置

工具

找到约 594,000 条结果 (用时 0.68 秒)

A new multiparameter integrated MELD model for prognosis of HBV ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5400346/> - 翻译此页

作者: Y Luo - 2016 - 被引用次数: 2 - 相关文章

2016年8月26日 - Hepatitis B virus related acute-on-chronic liver failure (HBV-ACLF) is one ... model for end-stage liver disease (MELD) assesses well the short-term ... In recent years, a number of improved models based on MELD, like serum ...

A comprehensive validation of HBV-related acute-on-chronic liver ...

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMed Central \(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) - 翻译此页

作者: Y Shen - 2016 - 被引用次数: 1 - 相关文章

2016年9月16日 - Caused by the acute exacerbation of chronic hepatitis B (CHB), hepatitis B virus-related acute-on-chronic liver failure (HBV-ACLF) is a severe ... patients have a poor prognosis with short-term mortality ranging from 30% to 70%. MELD: model of end-stage liver disease; MELD-Na: sodium MELD; MESO: ...

[PDF] Different models in predicting the short-term prognosis of patients wit...

annalsofhepatology.com.mx/PDF/vol11n3/HP123-05-Different.pdf - 翻译此页

作者: WB Yang - 被引用次数: 24 - 相关文章

hepatitis B virus-related acute-on-chronic liver failure ... King's College Hospital (KCH) criteria, model for end-stage liver disease (MELD), MELD combined.

Short-term efficacy of treating hepatitis B virus-related acute-on ...

link.springer.com/article/10.1007/s11655-016-2582-2 - 翻译此页

2016年5月24日 - Secondary outcome measures included Model for End-Stage Liver ... The integrative treatment was found to significantly improve the MELD score (t=2.353, P=0.020). ... hepatitis B virus-related acute-on-chronic liver failure mortality A model to predict survival in patients with end-stage liver disease.