



Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 35066

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Prognostic value of lymphovascular invasion in Bismuth-Corlette type IV hilar cholangiocarcinoma

Li B *et al.* Association between lymphovascular invasion and prognosis

Bei Li, Xian-Ze Xiong, Yong Zhou, Si-Jia Wu, Zhen You, Jiong Lu, Nan-Sheng Cheng


Abstract

AIM

To investigate the prognostic value of lymphovascular invasion (LVI) in Bismuth-Corlette type IV hilar cholangiocarcinoma (HC) patients.

Match Overview

1	Crossref 79 words Jong Hoon Lee. "Lymphovascular Invasion is a Significant Prognosticator in Rectal Cancer Patients Who Receive...	2%
2	Internet 57 words crawled on 05-Feb-2017 e-sciencedcentral.org	2%
3	Crossref 25 words Gurleyik, G., E. Gurleyik, F. Aker, A. Aktekin, S. Emir, O. Gungor, and A. Saglam. "Lymphovascular Invasion, as a ...	1%
4	Crossref 25 words Wang, X., X. Hong, H. Pang, H. Dai, L. You, W. Wu, and Y. Zhao. "Selecting optimal surgical procedures for intrad...	1%
5	Internet 18 words crawled on 15-Mar-2016 www.wjgnet.com	1%
6	Crossref 18 words Syed A. Hoda. "Issues Relating to Lymphovascular Invasion in Breast Carcinoma", Advances in Anatomic Patholo...	1%
7	Internet 15 words crawled on 31-Dec-2016 pdfs.semanticscholar.org	<1%
8	Crossref 14 words Shao-Hua Li, Qiao-Xuan Wang, Zhong-Yuan Yang, Wu Jiang, Cong Li, Peng Sun, Wei Wei, Ming Shi, Rong-Ping	<1%
9	Crossref 12 words "Abstract: Abstract", Diseases of the Esophagus, 2014.	<1%



找到约 2,830 条结果 (用时 0.73 秒)

Google 学术: Prognostic value of lymphovascular invasion for Bismuth-Corlette Type IV hilar cholangiocarcinoma patients with radical resection

... survival after resection for hilar cholangiocarcinoma. - Su - 被引用次数: 299

Extended resections for hilar cholangiocarcinoma - Neuhaus - 被引用次数: 617

... in type IIIA and IIIB hilar cholangiocarcinoma: a 15-year ... - Kow - 被引用次数: 49

Hilar cholangiocarcinoma: diagnosis, treatment options, and ...

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: KC Soares - 2014 - 被引用次数: 68 - 相关文章

Hilar cholangiocarcinoma (HC) is a rare disease with a poor prognosis which Unlike the Bismuth-Corlette and MSKCC staging systems, the AJCC system ... Direct invasion into hilar structures as well as lymphatic and perineural invasion is ... Patients are generally eligible for resection 4-6 weeks after biliary drainage and ...

Extended Resections for Hilar Cholangiocarcinoma - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: P Neuhaus - 1999 - 被引用次数: 612 - 相关文章

Surgical resection of hilar cholangiocarcinoma is the only potentially curative treatment. ... Radical resections are currently considered as optimal treatment, but <20% of ... In principle, patients with Bismuth-Corlette type I or type II tumors can ... and peripancreatic tumor extension into lymphatic vessels or perineural sheaths ...

Outcome of surgical resection in Klatskin tumors - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: A Serrablo - 2013 - 被引用次数: 23 - 相关文章

2013年7月15日 - Hilar cholangiocarcinoma (HCC) is the most common type, and ... patients with regard to prognosis and that depth of tumor invasion is a ... et al[51] reported better survival for patients with Bismuth types III/IV ... a statistically significance of radiotherapy for R1 radical resection of LYMPHATIC SPREAD.

找到约 3,230 条结果 (用时 0.56 秒)

Google 学术: Prognostic value of lymphovascular invasion in Bismuth-Corlette type IV hilar cholangiocarcinoma

... survival after resection for hilar cholangiocarcinoma. - Su - 被引用次数: 300

Surgical treatment of hilar cholangiocarcinoma in the ... - Lee - 被引用次数: 177

... for hilar and peripheral cholangiocarcinoma - Robles - 被引用次数: 218

Hilar cholangiocarcinoma: diagnosis, treatment options, and ...

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: KC Soares - 2014 - 被引用次数: 77 - 相关文章

Hilar cholangiocarcinoma (HC) is a rare disease with a poor prognosis The Bismuth-Corlette classification of hilar cholangiocarcinoma. ... Direct invasion into hilar structures as well as lymphatic and perineural invasion is commonly seen. ... Patients are generally eligible for resection 4-6 weeks after biliary drainage and ...

Extended Resections for Hilar Cholangiocarcinoma - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: P Neuhaus - 1999 - 被引用次数: 620 - 相关文章

Surgical resection of hilar cholangiocarcinoma is the only potentially curative treatment. ... peripancreatic tumor extension into lymphatic vessels or perineural sheaths ... Few patients with Bismuth-Corlette type II carcinomas underwent right or left In type IV tumors, R0 resections were achieved in 18 of 25 patients (72%; ...

Management of perihilar cholangiocarcinoma in the era of multimodal ...

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: DC Vicente Valero III - 2012 - 被引用次数: 32 - 相关文章

MDCT has been shown to be accurate for the assessment of tumor invasion and to the confluence and extend into the main left hepatic duct and type IV tumors ... TNM staging for hilar cholangiocarcinoma: anatomic stage/prognostic groups. hepatic duct (Bismuth-Corlette type IIIb) that necessitate a left hepatectomy.

[全部](#) [图片](#) [新闻](#) [购物](#) [视频](#) [更多](#)[设置](#) [工具](#)

找到约 3,220 条结果 (用时 0.98 秒)

Google 学术: Prognostic value of lymphovascular invasion in Bismuth-Corlette type IV hilar cholangiocarcinoma

[... survival after resection for hilar cholangiocarcinoma.](#) - Su - 被引用次数: 301

[Surgical treatment of hilar cholangiocarcinoma in the ...](#) - Lee - 被引用次数: 178

[... for hilar and peripheral cholangiocarcinoma](#) - Robles - 被引用次数: 221

Hilar cholangiocarcinoma: diagnosis, treatment options, and ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: KC Soares - 2014 - 被引用次数: 78 - 相关文章

Hilar cholangiocarcinoma (HC) is a rare disease with a poor prognosis The Bismuth-Corlette classification of hilar cholangiocarcinoma. ... Direct **invasion** into hilar structures as well as **lymphatic** and **perineural invasion** is commonly seen. ... Patients are generally eligible for resection 4-6 weeks after biliary drainage and ...

Management of perihilar cholangiocarcinoma in the era of multimodal ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: DC Vicente Valero III - 2012 - 被引用次数: 32 - 相关文章

MDCT has been shown to be accurate for the assessment of tumor **invasion** and to the confluence and extend into the main left hepatic duct and **type IV** tumors ... TNM staging for **hilar cholangiocarcinoma**: anatomic stage/prognostic groups. hepatic duct (**Bismuth-Corlette** type IIIb) that necessitate a left hepatectomy.

Hilar cholangiocarcinoma: diagnosis and staging - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: W Jarnagin - 2005 - 被引用次数: 48 - 相关文章

Cancer arising from the proximal biliary tree, or **hilar cholangiocarcinoma**, ... although rarely other malignant cell types may arise primarily in the biliary tree. (2) **vascular invasion**; (3) hepatic lobes