

SCORING SYSTEMS FOR PEPTIC ULCER BLEEDING: WHICH ONE TO US

全部 图片 视频 新闻 地图 图书

找到约 285,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Upper gastrointestinal bleeding risk scores: Who, when and why?](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4753192/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4753192/>

15 Feb 2016 ... The use of risk scoring systems in early assessment of patients suffering ... cited ones are the Rockall score and the Glasgow Blatchford score (GBS). with bleeding peptic ulcer who had undergone endoscopic therapy[38].

[Hospitalization for peptic ulcer bleeding: evaluation of a risk scoring ...](https://www.ncbi.nlm.nih.gov/pubmed/11142555)

<https://www.ncbi.nlm.nih.gov/pubmed/11142555>

Hospitalization for peptic ulcer bleeding: evaluation of a risk scoring system in ... all patients admitted to our unit for bleeding peptic ulcer over a one-year period. need for surgery as well as mortality in the two periods, using Student t test.

[Management of Patients with Ulcer Bleeding | American College of ...](https://gi.org/.../management-of-patients-with-ulcer-bleeding/)

<https://gi.org/.../management-of-patients-with-ulcer-bleeding/> ▼

For the subset of patients with actively bleeding ulcers, thermal therapy or In general, risk assessment with scoring systems such as Blatchford or Rockall is not able to (41) assessed the histological characteristics of gastric ulcers with visible The one study using therapy matching current recommendations (vigorous ...

[Current Management of Peptic Ulcer Bleeding - Medscape](http://www.medscape.com/viewarticle/521189_2)

www.medscape.com/viewarticle/521189_2 ▼

12 Jan 2006 ... Current Management of Peptic Ulcer Bleeding ... and validated its ability to predict recurrent bleeding and death using data from 1,625 patients (Table 1). Their scoring system is based on age, comorbidities, the presence of ...

[Forrest classification - Wikipedia](https://en.wikipedia.org/wiki/Forrest_classification)

https://en.wikipedia.org/wiki/Forrest_classification ▼

Forrest classification is a classification of upper gastrointestinal hemorrhage used for purposes ... of endoscopic therapy and urgent surgery in bleeding peptic ulcers". ... RISK SCORING SYSTEM IN PREDICTION OF REBLEEDING IN PEPTIC ULCER ... By using this site, you agree to the Terms of Use and Privacy Policy.

[Forrest classification - Application](#)

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 35383

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

Scoring systems for peptic ulcer bleeding: Which one to use?

Ivan Budimir, Sanja Stojšavljević, Neven Baršić, Alen Biščanin, Gorana Mirošević,
Sven Bohnec, Lora Stanka Kirigin, Tajana Pavić, Neven Ljubičić

Abstract

AIM

To compare the Glasgow Blatchford score (GBS), Rockall score (RS) and Baylor

1

Match Overview

1	Crossref 358 words Budimir, Ivan, Marina Gradišer, Marko Nikolić, Neven Baršić, Neven Ljubičić, Dominik Kralj, and Ivan Budimir jr.. "Glas	9%
2	Crossref 78 words Bryant, Robert V., Paul Kuo, Kate Williamson, Chantelle Yam, Mark N. Schoeman, Richard H. Holloway, and Nam Q. N	2%
3	Internet 74 words crawled on 18-Feb-2017 www.wjgnet.com	2%
4	Crossref 69 words Hyett, Brian H., Marwan S. Abougergi, Joseph P. Charpentier, Navin L. Kumar, Suzana Brozovic, Brian L. Claggett, An	2%
5	Crossref 32 words M.E. Leerdam. "Acute upper GI bleeding: did anything change? Time trend analysis of incidence and outcome of ac	1%
6	Crossref 24 words Fischer, Anja, Marcus Fischer, Robert A. Nicholls, Stephanie Lau, Jana Poettgen, Kostas Patas, Christoph Heesen, an	1%
7	Crossref 10 words Kim, Jung-Wook, Hyun Jin Park, Jae-Young Jang, Chi Hyuk Oh, Jae-Jun Shim, Chang Kyun Lee, and Young Woon Cha	<1%



Scoring systems for peptic ulcer bleeding: Which one to use?



学术搜索

找到约 23,600 条结果 (用时0.12秒)

我的著作引用情况

文章

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

我的图书馆

Validation of the Rockall risk scoring system in upper gastrointestinal bleeding

[HTML] bmj.com

EM Vreeburg, CB Terwee, P Snel, EAJ Rauws... - Gut, 1999 - gut.bmj.com

... results of a logistic regression analysis and includes three clinical variables (age, shock, and comorbidity) and two endoscopic variables (diagnosis and major SRH), each categorised and scored with 0 ... We used this scoring system to assess the individual risk score for each ...

被引用次数: 256 相关文章 所有 12 个版本 引用 保存

Management of patients with ulcer bleeding

[HTML] gi.org

L Laine, DM Jensen - The American journal of ..., 2012 - search.proquest.com

... endoscopy may be considered in very low-risk patients with Blatchford scores of 0. ... Risk scoring systems to predict need for clinical intervention for patients with nonvariceal upper ... Comparing the Blatchford and pre-endoscopic Rockall score in predicting the need for endoscopic ...

被引用次数: 387 相关文章 所有 25 个版本 引用 保存

Discriminant value of dyspeptic symptoms: a study of the clinical presentation of 221 patients with dyspepsia of unknown cause, peptic ulceration, and cholelithiasis.

[PDF] bmj.com

NJ Talley, D McNeil, DW Piper - Gut, 1987 - gut.bmj.com

... dyspepsia was 94%, but only prospective studies will determine if this scoring system is of ... specificity, and predictive values were calculated for the diagnostic symptom scores based on ... Critical score values for the symptoms were then constructed based on their sensitivity and ...

被引用次数: 121 相关文章 所有 11 个版本 引用 保存

[引用] Long-term nonsteroidal antiinflammatory drug use and Helicobacter pylori infection

[PDF] gastrojournal.org

DY Graham, MD Lidsky, AM Cox, DJ Evans, DG Evans... - Gastroenterology, 1991 - Elsevier

被引用次数: 207 相关文章 所有 8 个版本 引用 保存

Successful outpatient management of acute upper gastrointestinal hemorrhage: use of practice guidelines in a large patient series

GF Longstreth, SP Feitelberg - Gastrointestinal endoscopy, 1998 - Elsevier

... 9 To use the scoring system as closely as possible according to its developers' intentions, we obtained examples of scored comorbidity from the principal author. ... Esophageal malignancy, 2 (1). Normal, 4 (2). Miscellaneous, 3 (2). Risk score, 0, 14 (8), 1, 55 (31), 2, 50 (28). ...

被引用次数: 166 相关文章 所有 8 个版本 引用 保存

[HTML] International consensus recommendations on the management of patients

[HTML] annals.org

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☒ 创建快讯

[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 254,000 条结果 (用时 0.60 秒)

Upper gastrointestinal bleeding risk scores: Who, when and why?

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: S Monteiro - 2016 - 被引用次数: 14 - 相关文章

2016年2月15日 - The use of risk scoring systems in early assessment of patients suffering ... cited ones are the Rockall score and the Glasgow Blatchford score (GBS). with bleeding peptic ulcer who had undergone endoscopic therapy[38].

Treatment and prognosis in peptic ulcer bleeding. - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/24547604> - 翻译此页

作者: SB Laursen - 2014 - 被引用次数: 13 - 相关文章

Use of risk scoring systems is shown to be advantageous in the primary assessment of patients presenting with symptoms of peptic ulcer bleeding. Studies ... Numbers needed to treat in order to avoid one case of rebleeding was 10. Study III ...

Comparison of AIMS65 Score and Other Scoring Systems for ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: SM Park - 2016 - 被引用次数: 2 - 相关文章

2016年7月15日 - Upper gastrointestinal bleeding (UGIB) is a common medical ... The main cause of UGIB is peptic ulcer disease, which accounts for ... AIMS65 score with other scoring systems; and validated the use of ... (6) could not obtain at least one of the four scoring systems; and (7) followed up for fewer than 30 days.

Scoring systems for outcome prediction in patients with perforated ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3626602/> - 翻译此页

作者: K Thorsen - 2013 - 被引用次数: 41 - 相关文章

2013年4月10日 - Keywords: Perforated peptic ulcer, Scoring systems, Outcome prediction, Mortality, Morbidity ... Perforation is the second most frequent complication after bleeding [1]. ... The ROC curve