

Increased intestinal mucosal leptin levels in patients with diarrhea-predominant iri

全部 图片 视频 新闻 地图 图书

找到约 124,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

Google 学术: Increased intestinal mucosal leptin levels in patients with diarrhea-predominant irritable bowel syndrome

Gluten causes gastrointestinal symptoms in subjects ... - Biesiekierski - 被引用次数: 623

The human gut microbiome: implications for future ... - Kinross - 被引用次数: 125

Sleisenger and Fordtran's Gastrointestinal and Liver ... - Feldman - 被引用次数: 979

PubMed Result - NCBI

https://www.ncbi.nlm.nih.gov/pubmed?linkname=pubmed_pubmed...

Plasma and gastric mucosal 5-hydroxytryptamine concentrations following ... Increased serum free tryptophan in patients with diarrhea-predominant ... Decreased levels of kynurenic acid in the intestinal mucosa of IBS patients: ... Moradi A, Joshaghani H. Serum leptin levels and irritable bowel syndrome: a new hypothesis.

Subclinical mucosal inflammation in diarrhea-predominant irritable ...

<https://www.ncbi.nlm.nih.gov/pubmed/22486731>

10 Apr 2012 ... Subclinical mucosal inflammation in diarrhea-predominant irritable bowel syndrome (IBS) in a tropical setting. ... we investigated 49 patients with IBS-D (cases; median age 34 years (range 18-59); M:F ... In cases, there were significantly more mucosal mast cells in the ileum and all segments of colon and ...

Serum biomarkers for irritable bowel syndrome - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4632880/>

1 Jul 2015 ... Irritable bowel syndrome (IBS) is one of the most frequent and ... Besides mastocytes, in the intestinal mucosa of IBS patients also an elevated number of ... to evaluate serum levels of leptin and their relation with IBS found that ... in differentiating a subset of IBS - diarrhea predominant IBS (IBS-D) [44].

Patients with constipation-predominant irritable bowel syndrome ...

<https://www.ncbi.nlm.nih.gov/pubmed/11351146>

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 36452

Manuscript Type: ORIGINAL ARTICLE

Case Control Study

Increased intestinal mucosal leptin levels in patients with
diarrhea-predominant irritable bowel syndrome

De-Rong Liu, Xiao-Juan Xu, Shu-Kun Yao

Abstract

AIM

To measure leptin levels and analyze the relationship of leptin with clinical features, visceral sensitivity, mast cells, and nerve fibers.

METHODS

Forty-two patients with diarrhea-predominant irritable bowel syndrome (IBS-D) fulfilling the Rome III criteria and 20 age- and sex-matched healthy

Match Overview

- 1** **Crossref** 35 words
X. J. Xu, Y. L. Zhang, L. Liu, L. Pan, S. K. Yao. "Increased expression of nerve growth factor correlates with visceral hyp... 1%
- 2** **Crossref** 22 words
Likun Zhong. "Pathophysiologic Findings of Irritable Bowel... Syndrome in China", Journal of Neurogastroenterology and 1%

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

The human gut microbiome: implications for future health care

JM Kinross, AC von Roon, E Holmes, A Darzi... - Current ..., 2008 - Springer

... of germ-free mice with an "obese microbiota" results in a significantly greater increase in total body ... Increased levels of formate, a microbial co-metabolite, were significantly associated with higher blood pressure in multiple regres- sion analyses, as were high alanine and ...

☆ 99 被引用次数: 126 相关文章 所有 9 个版本

[图书] Sleisenger and Fordtran's Gastrointestinal and Liver Disease E-Book: Pathophysiology, Diagnosis, Management, Expert Consult Premium Edition- Enhanced ...

M Feldman, LS Friedman, LJ Brandt - 2010 - books.google.com

... Kaiser Permanente, San Rafael, California Diverticular Disease of the Colon Amy E ... Diverticula of the Pharynx, Esophagus, Stomach, and Small Intestine; Abdominal Hernias and ... Attending Physician, Fletcher Allen Health Care, Burlington, Vermont Intestinal Protozoa Steven H ...

☆ 99 被引用次数: 988 相关文章 所有 6 个版本

Gluten causes gastrointestinal symptoms in subjects without celiac disease: a double-blind randomized placebo-controlled trial

JR Biesiekierski, ED Newnham, PM Irving... - The American journal of ..., 2011 - nature.com

... This may lead to increased smooth muscle contractility and indirectly have effects on luminal ... of the tight junction (16), induce apoptosis (14,30), and increase oxidative stress (15 ... of short-chain carbohydrates in common Australian vegetables and fruits by high-performance liquid ...

☆ 99 被引用次数: 636 相关文章 所有 14 个版本

Traditional thoughts on the pathophysiology of irritable bowel syndrome

WL Hasler - Gastroenterology Clinics of North America, 2011 - Elsevier

... However, using wireless motility capsules, increases in colonic contractile activity were noted in C ... gastrocolonic responses lasting up to 3 hours, especially after high-fat meals ... Patients with IBS show increased sigmoid colon and plasma cholecystokinin and vasoactive intestinal ...

☆ 99 被引用次数: 52 相关文章 所有 7 个版本

Probiotics, irritable bowel syndrome, and inflammatory bowel disease

MH Floch - Current treatment options in gastroenterology, 2003 - Springer

... of the flora was very high in patients with Crohn's disease, with an increase in the ... Nejdfor P, Wang Q, Ekelund M, et al.: Increased colonic permeability in patients with ulcerative colitis: an ... Mimura T, Pizzello E, Schreiber S, et al.: Once daily high dose probiotic therapy maintain...

[PDF] uj.edu.pl

[PDF] foodtest.com.tr

[全部](#)[新闻](#)[图片](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 24,700 条结果 (用时 0.55 秒)

Google 学术 : Increased intestinal mucosal leptin levels in patients with diarrhea-predominant irritable bowel syndrome

The human gut microbiome: implications for future ... - Kinross - 被引用次数 : 126

Sleisenger and Fordtran's Gastrointestinal and Liver ... - Feldman - 被引用次数 : 988

Gluten causes gastrointestinal symptoms in subjects ... - Biesiekierski - 被引用次数 : 636

Serum biomarkers for irritable bowel syndrome - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4632880/> - 翻译此页

作者 : A CHIRA - 2015 - 被引用次数 : 2 - 相关文章

2015年7月1日 - Irritable bowel syndrome (IBS) is one of the most frequent and ... Besides mastocytes, in the intestinal mucosa of IBS patients also an elevated number of ... to evaluate serum levels of leptin and their relation with IBS found that ... in differentiating a subset of IBS – diarrhea predominant IBS (IBS-D) [44].

[Abstract](#) · [Introduction](#) · [Materials and methods](#) · [Discussion](#)

Increased serum levels of lipopolysaccharide and anti-flagellin ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/26387872> - 翻译此页

作者 : A Dlugosz - 2015 - 被引用次数 : 19 - 相关文章

2015年9月21日 - Increased serum levels of lipopolysaccharide and anti-flagellin antibodies in patients with diarrhea-predominant irritable bowel syndrome. ... are likely important in microbial-host interactions and intestinal homeostasis. ... and activation of mucosal immunity against common microbial antigens might be ...

缺少字词 : leptin

Subclinical mucosal inflammation in diarrhea-predominant irritable ...

<https://www.ncbi.nlm.nih.gov/pubmed/22486731> - 翻译此页

作者 : AP De Silva - 2012 - 被引用次数 : 15 - 相关文章

2012年4月10日 - Subclinical mucosal inflammation in diarrhea-predominant irritable bowel syndrome (IBS) in a tropical setting. ... we investigated 49 patients with IBS-D (cases; median age 34 years (range 18-59); M:F ... In cases, there were significantly more mucosal mast cells in the ileum and all segments of colon and ...

缺少字词 : leptin levels

Immunological Biomarkers in Postinfectious Irritable Bowel Syndrome ...

[eys.com](#) > ... > Vol 22 Issue 4 > Abstract - 翻译此页