

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 36787

Manuscript Type: Original Article

#### Case Control Study

Risk factors for hepatic steatosis in adults with cystic fibrosis: Similarities to non-alcoholic fatty liver disease

Fares Ayoub, Cesar A Trillo-Alvarez, Giuseppe Morelli, Jorge Lascano

#### ABSTRACT

##### AIM

To investigate the clinical, biochemical and imaging characteristics of adult cystic fibrosis (CF) patients with hepatic steatosis as compared to normal CF controls.

##### METHODS

We performed a retrospective review of adult CF patients in an academic outpatient setting during 2016. Baseline characteristics, genetic mutation analysis as well as laboratory values were collected. Abdominal imaging (ultrasound, computed tomography, magnetic resonance) was used to determine presence of hepatic steatosis.

### Match Overview

1	<b>Crossref</b> 192 words Fares Ayoub, Jorge Lascano, Giuseppe Morelli. "Proton Pump Inhibitor Use Is Associated With an Increased Frequency (...)	6%
2	<b>Internet</b> 20 words crawled on 01-Apr-2014 <a href="http://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a>	1%
3	<b>Internet</b> 17 words crawled on 06-Jun-2009 <a href="http://www.sonoraquest.com">www.sonoraquest.com</a>	1%
4	<b>Internet</b> 16 words crawled on 02-Jul-2017 <a href="http://www.rug.nl">www.rug.nl</a>	1%
5	<b>Internet</b> 15 words crawled on 12-Sep-2017 <a href="http://www.infona.pl">www.infona.pl</a>	<1%


找到约 99,100 条结果 (用时 0.47 秒)

## Nonalcoholic Fatty Liver Disease - Cleveland Clinic Center for ...

[www.clevelandclinicmeded.com/.../diseasemanagement/.../nonalcoholic-fatt...](http://www.clevelandclinicmeded.com/.../diseasemanagement/.../nonalcoholic-fatt...)  翻译此页

Nonalcoholic fatty liver disease (NAFLD) is one of the most common causes of chronic liver disease. ... Cystic fibrosis; Abetalipoproteinemia; Others ... NASH cirrhosis is a risk factor for development of hepatocellular carcinoma (HCC). .... In adults, similar histologic findings can be found in a number of conditions (Table 1).


## Non-alcoholic fatty liver disease and non-alcoholic steatohepatitis

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral\(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) -  翻译此页

作者: R Vuppalanchi - 2009 - 被引用次数: 618 - 相关文章

Nonalcoholic fatty liver disease (NAFLD) is one of the most common causes of elevated liver ... Its incidence in adults and children is rising rapidly due to the ongoing ... alcoholic fatty liver. However, in individuals with metabolic risk factors such as ... It remains controversial whether ballooning or pericellular fibrosis that ...


## Cystic Fibrosis Related Liver Disease—Another Black Box in Hepatology

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral\(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) -  翻译此页

作者: K Stauffer - 2014 - 被引用次数: 9 - 相关文章

2014年8月4日 · Cystic fibrosis (CF) is the most frequent fatal autosomal recessive disorder in Caucasians. ... to be an independent risk factor for mortality and lung transplantation, .... Non-alcoholic fatty liver disease (NAFLD) is now considered one of ... [37,39] adding to the similarities between these two disease entities,

## Non alcoholic fatty liver disease and metabolic syndrome - NCBI - NIH

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral\(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) -  翻译此页

作者: P Paschos - 2009 - 被引用次数: 314 - 相关文章



[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 173,000 条结果 (用时 0.54 秒)

**Google 学术 : Risk factors for hepatic steatosis in adults with cystic fibrosis: Similarities to non-alcoholic fatty liver disease**

... and management of non-alcoholic fatty liver disease: ... - [Chalasan](#) - 被引用次数 : 971

... and management of non-alcoholic fatty liver disease: ... - [Chalasan](#) - 被引用次数 : 2153

Prevalence of fatty liver in children and adolescents - [Schwimmer](#) - 被引用次数 : 1026

**Non-Invasive Evaluation of Cystic Fibrosis Related Liver Disease in ...**

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3405060/> - 翻译此页

作者 : T Karlas - 2012 - 被引用次数 : 27 - 相关文章

2012年7月25日 - Non-Invasive Evaluation of Cystic Fibrosis Related Liver Disease in Adults with ARFI, Transient Elastography and Different Fibrosis Scores. Thomas Karlas,<sup>#1</sup> Marie ... Non-cirrhotic CFLD individuals showed elastography values similar to CF patients without liver involvement. Cases with liver cirrhosis ...

**Pediatric non alcoholic fatty liver disease: old and new concepts on ...**

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3620555/> - 翻译此页

作者 : V Giorgio - 2013 - 被引用次数 : 110 - 相关文章

2013年3月25日 - It includes a spectrum of disease ranging from intrahepatic fat accumulation (steatosis) to various degrees of necrotic inflammation and fibrosis (non-alcoholic steatohepatitis [NASH]). NAFLD is associated, in children as in adults, with severe metabolic impairments determining an increased risk of ...

[全部](#)[新闻](#)[图片](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 165,000 条结果 (用时 0.47 秒)

## Cystic Fibrosis Related Liver Disease—Another Black Box in Hepatology

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4159809/> ▼ 翻译此页

作者: K Staufer - 2014 - 被引用次数: 9 - 相关文章

2014年8月4日 - CF-related liver disease (CFLD) was recently shown to be an independent risk factor for mortality and lung transplantation, and has become one of the .... Non-alcoholic fatty liver disease (NAFLD) is now considered one of the most important causes of liver disease in Western countries, ranging from simple ...

## Fatty liver in lean patients: is it a different disease? - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/3959355/> - 翻译此页

作者: MV Machado - 2012 - 被引用次数: 11 - 相关文章

2011年11月29日 - Nonalcoholic fatty liver disease (NAFLD) has widely been considered a manifestation of metabolic syndrome in close relation to obesity [1]. In fact, although obesity is undoubtedly one of the main risk factors for fatty liver, since its first description it is known that it can also occur in lean subjects. Studies in ...

## Non-Invasive Evaluation of Cystic Fibrosis Related Liver Disease in ...

[journals.plos.org/plosone/article?id=10.1371/journal.pone.0042139](https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0042139) ▼ 翻译此页

作者: T Karlas - 2012 - 被引用次数: 27 - 相关文章

2012年7月25日 - Healthy probands and patients with alcoholic liver cirrhosis served as controls. Results Fourteen CF patients met CFLD criteria, six had liver cirrhosis. Elastography acquisition was successful in >89% of cases. Non-cirrhotic CFLD individuals showed elastography values similar to CF patients without liver ...

## Nonalcoholic fatty liver disease: Definitions, risk factors, and workup ...

[onlinelibrary.wiley.com/doi/10.1002/cld.81/full](https://onlinelibrary.wiley.com/doi/10.1002/cld.81/full) ▼ 翻译此页