

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☒ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

ARFI elastography in patients with chronic autoimmune liver diseases: A preliminary study

[HTML] nih.gov

S Righi, E Fiorini, C De Molo, V Cipriano, F Cassani... - Journal of ultrasound, 2012 - Elsevier

... In our opinion, the correlation between ARFI elastographic data and hepatic fibrosis varies depending on the type of liver ... The diagnostic performance of the latter method is good in patients with PBC or PSC [17] and [32], and the same might be true of ARFI elastography. ...

☆ 99 被引用次数: 22 相关文章 所有 8 个版本

[HTML] Unmet clinical need in autoimmune liver diseases

[HTML] sciencedirect.com

JK Dyson, G Webb, GM Hirschfield, A Lohse... - Journal of ..., 2015 - Elsevier

... to PBC, with unmet clinical need particularly relating to extra-hepatic disease and ... the very least allow identification of high-risk patients in whom timely liver transplantation can ... clinical practice is based around CA19-9 measurement, conventional radiological assessment and/or ...

☆ 99 被引用次数: 35 相关文章 所有 23 个版本

Autoimmune hepatitis in southern Israel: A 15 - year multicenter study

JS Delgado, A Vodonos, S Malnick... - Journal of digestive ..., 2013 - Wiley Online Library

... Previous article in issue: Performance of transient elastography (TE) and ... Email: delgado@bgu.ac.il. Abstract. Objective. In this study we aimed to assess the incidence, prevalence and clinical outcomes of patients with autoimmune hepatitis (AIH) in southern Israel. Methods. ...

☆ 99 被引用次数: 38 相关文章 所有 6 个版本

Noninvasive markers in the assessment and management of autoimmune liver diseases

[PDF] semanticscholar.org

D Grunwald, D Kothari, R Malik - European journal of ..., 2014 - journals.lww.com

... Skip Navigation Links Home > October 2014 - Volume 26 - Issue 10 > Noninvasive markers in the assessment and management of auto...

☆ 99 被引用次数: 10 相关文章 所有 5 个版本

5

Name of Journal: *World Journal of Gastroenterology***Manuscript NO:** 36949**Manuscript Type:** ORIGINAL ARTICLE*Retrospective Study***Performance of transient elastography in assessing liver fibrosis in patients with autoimmune hepatitis-primary biliary cirrhosis overlap syndrome**Wu HM *et al.* Transient elastography and AIH-PBC overlap syndromeHui-Min Wu, Li Sheng, Qi Wang, Han Bao, Qi Miao, Xiao Xiao, Can-Jie Guo,
Hai Li, Xiong Ma, De-Kai Qiu, Jing Hua**Abstract****AIM**

To investigate the performance of transient elastography (TE) for diagnosis of fibrosis in patients with autoimmune hepatitis and primary biliary cirrhosis

Match Overview

1	Internet 53 words crawled on 28-Sep-2016 www.pubfacts.com	2%
2	Internet 39 words crawled on 08-Dec-2010 www.biomedcentral.com	1%
3	Internet 27 words crawled on 14-Jun-2017 repository.um.edu.my	1%
4	Internet 16 words crawled on 22-Jul-2017 www.wjgnet.com	1%
5	Internet 15 words crawled on 05-Oct-2017 real.mtak.hu	1%
6	Crossref 14 words Nahon, P. "Assessment of liver fibrosis using transient ... astography in patients with alcoholic liver disease", Jour	<1%
7	Crossref 14 words "Abstracts of the 25th Annual Conference of APASL, Fe ... ruary 20-24, 2016, Tokyo, Japan", Hepatology Internatio	<1%
8	Internet 14 words crawled on 16-Sep-2017 diposit.ub.edu	<1%
9	Internet 13 words crawled on 26-Oct-2017 www.e-sciencecentral.org	<1%

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 18,300 条结果 (用时 0.50 秒)

Transient elastography to assess hepatic fibrosis in primary biliary ...

onlinelibrary.wiley.com/doi/10.1111/j.1365-2036.2007.03585.x/full - 翻译此页

作者: E GÓMEZ - DOMINGUEZ - 2008 - 被引用次数: 56 - 相关文章

2007年12月10日 - Methods Liver stiffness measurements were generated in 80 patients with primary biliary cirrhosis by applying transient elastography; however, as there ... (PBC) and primary sclerosing cholangitis (PSC), as well as a number of variants such as autoimmune cholangitis (AIC) and certain overlap syndromes.

Assessment of biliary fibrosis by transient elastography in patients with ...

onlinelibrary.wiley.com/doi/10.1002/hep.21151/full - 翻译此页

作者: C Corpechot - 2006 - 被引用次数: 426 - 相关文章

2006年4月20日 - The aim of the current study was to assess the diagnostic performance of transient elastography for the evaluation of fibrosis and histological stages in chronic cholestatic diseases. Abbreviations. PBC, primary biliary sclerosis; PSC, primary sclerosing cholangitis; LB, liver biopsy; LSM, liver stiffness ...

Transient Elastography (FibroScan) Performs Better Than Non ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5669534/> - 翻译此页

作者: L Guo - 2017

2017年10月26日 - Transient Elastography (FibroScan) Performs Better Than Non-Invasive Markers in Assessing Liver Fibrosis and Cirrhosis in Autoimmune Hepatitis Patients We excluded all patients with hepatitis B or C virus infection, excessive alcohol consumption, primary biliary cirrhosis (PBC), drug-induced liver ...

Non-invasive assessment of liver fibrosis using two-dimensional shear ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5514650/> - 翻译此页

作者: J Zeng - 2017

2017年7月14日 - Non-invasive assessment of liver fibrosis using two-dimensional shear wave elastography in patients with autoimmune liver diseases ... Autoimmune hepatitis (AIH), primary biliary cholangitis (PBC), and primary sclerosing cholangitis (PSC) are the three major forms of AILD, which differ according to the ...

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 2,930 条结果 (用时 0.42 秒)

Assessment of biliary fibrosis by transient elastography in patients with ...

onlinelibrary.wiley.com/doi/10.1002/hep.21151/full - 翻译此页

作者: C Corpechot - 2006 - 被引用次数: 427 - 相关文章

2006年4月20日 - The aim of the current study was to assess the diagnostic performance of transient elastography for the evaluation of fibrosis and histological stages in chronic cholestatic diseases.

Abbreviations. PBC, primary biliary sclerosis; PSC, primary sclerosing cholangitis; LB, liver biopsy; LSM, liver stiffness ...

Transient elastography to assess hepatic fibrosis in primary biliary ...

onlinelibrary.wiley.com/doi/10.1111/j.1365-2036.2007.03585.x/full - 翻译此页

作者: E GÓMEZ-DOMINGUEZ - 2008 - 被引用次数: 56 - 相关文章

2007年12月10日 - Methods Liver stiffness measurements were generated in 80 patients with primary biliary cirrhosis by applying transient elastography; however, as there ... (PBC) and primary sclerosing cholangitis (PSC), as well as a number of variants such as autoimmune cholangitis (AIC) and certain overlap syndromes.

Transient Elastography (FibroScan) Performs Better Than Non ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5669534/> - 翻译此页

作者: L Guo - 2017

2017年10月26日 - Transient Elastography (FibroScan) Performs Better Than Non-Invasive Markers in Assessing Liver Fibrosis and Cirrhosis in Autoimmune Hepatitis Patients We excluded all patients with hepatitis B or C virus infection, excessive alcohol consumption, primary biliary cirrhosis (PBC), drug-induced liver ...

ARFI elastography in patients with chronic autoimmune liver diseases ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3558243/> - 翻译此页

作者: S Righi - 2012 - 被引用次数: 23 - 相关文章

2012年10月25日 - We enrolled 9 adult patients (8 women, 1 man; age 48.1 ± 12.8 years) with chronic autoimmune disease (primary biliary cirrhosis (PBC, $n = 3$), autoimmune hepatitis (AIH, $n = 2$), primary sclerosing cholangitis (PSC, $n = 1$) and overlap syndromes, ($n = 3$) who underwent a liver biopsy and 11 healthy ...

Performance and Utility of Transient Elastography and Non-Invasive ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5203728/> - 翻译此页

作者: OE Anastasiou - 2016 - 被引用次数: 2 - 相关文章

2016年10月31日 - Performance and Utility of Transient Elastography and Non-Invasive Markers of Liver