

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 3,950 条结果 (用时 0.43 秒)

Benefits of long-term therapy with nucleos(t)ide analogues in ...

www.tandfonline.com/doi/full/10.1080/03007995.2016.1264932 - 翻译此页

作者: L Wei - 2017 - 被引用次数: 3 - 相关文章

Benefits of long-term therapy with nucleos(t)ide analogues in treatment-naïve patients with chronic hepatitis B in particular hepatitis B vaccination of infants and people considered at high risk of HBV infection (e.g. those who frequently require blood products or dialysis, prisoners in crowded jails, intravenous drug abusers ...

ide analogues in treatment-naïve patients with chronic hepatitis B

www.tandfonline.com/doi/pdf/10.1080/03007995.2016.1264932 - 翻译此页

作者: L Wei - 2017 - 被引用次数: 3 - 相关文章

improved the clinical status of patients with decompensated cirrhosis, the risk of cirrhotic complica- ... Although the prevalence of HBV infection has decreased ... Benefits of long-term nucleos(t)ide analogue therapy in treatment-naïve patients with chronic hepatitis B. Literature search strategy. Because the majority of studies ...

13-year nationwide cohort study of chronic kidney disease risk among ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4508999/> ▼ 翻译此页

作者: YC Chen - 2015 - 被引用次数: 14 - 相关文章

2015年7月22日 - 13-year nationwide cohort study of chronic kidney disease risk among treatment-naïve patients with chronic hepatitis B in Taiwan. Yi-Chun Chen In Taiwan, the NHI program has reimbursed nucleos(t)ide analogues for patients with chronic HBV infection since October 1, 2003 [22]. Thus, we could identify ...
缺少字词: decreases acquiring

A Taiwanese Nationwide Cohort Study Shows Interferon-Based ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4616715/> ▼ 翻译此页

作者: YC Chen - 2015 - 被引用次数: 5 - 相关文章

2015年8月14日 - A Taiwanese Nationwide Cohort Study Shows Interferon-Based Therapy for Chronic Hepatitis C Reduces the Risk of Chronic Kidney Disease. Yi-Chun Chen, MD, In conclusion, IBT, especially for 6 or more months, is associated with reduced risk of CKD in HCV-infected patients. Hyperlipidemia, diabetes

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 37011

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Nationwide cohort study suggests nucleos(t)ide analogues decreases dialysis risk in Taiwanese chronic kidney disease patients acquiring hepatitis B virus infection

Chen YC *et al.* Dialysis risk of HBV-infected CKD patients treated with and without nucleos(t)ide analogues

Yi-Chun Chen, Chung-Yi Li, Shiang-Jiun Tsai, Yen-Chun Chen

Abstract

AIM

To investigate the risk of end-stage renal disease (ESRD) in HBV-infected

Match Overview

1	Crossref 201 words Chen, Yi-Chun, Shang-Jyh Hwang, Chung-Yi Li, Chia-Pin Wu, and Li-Chu Lin. "A Taiwanese Nationwide Cohort	4%
2	Internet 76 words crawled on 25-Nov-2014 jama.jamanetwork.com	2%
3	Internet 67 words crawled on 22-Nov-2017 hub.hku.hk	1%
4	Crossref 53 words Chen, Yi-Chun, Yu-Chieh Su, Chung-Yi Li, Chia-Pin Wu, and Moon-Sing Lee. "A nationwide cohort study suggest	1%
5	Internet 49 words crawled on 05-Oct-2016 bmcnephrol.biomedcentral.com	1%
6	Internet 34 words crawled on 22-Dec-2015 tndt.org	1%
7	Crossref 18 words B M Brenner. "Predicting initiation and progression of c ... onic kidney disease: Developing renal risk scores", Kidn	<1%
8	Internet 17 words crawled on 04-Aug-2017 linknovate.com	<1%
9	Internet 17 words crawled on 10-Jul-2016 www.citeulike.org	<1%

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 1,020 条结果 (用时 0.52 秒)

[A nationwide cohort study suggests chronic hepatitis B virus infection ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/25426815> ▾ [翻译此页](#)

作者: YC Chen - 2015 - 被引用次数: 18 - 相关文章

2014年11月26日 - A nationwide cohort study suggests chronic hepatitis B virus infection increases the risk of end-stage renal disease among patients in Taiwan. ... We identified 17,758 adults who had chronic HBV infection and had not taken nucleos(t)ide analogs from 1999 to 2010 and randomly selected 71,032 matched ...

[Nucleos\(t\)ide analogues associated with a reduced risk of ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pubmed/25537961> - [翻译此页](#)

作者: JP Wang - 2015 - 被引用次数: 12 - 相关文章

2014年12月23日 - Nucleos(t)ide analogues associated with a reduced risk of hepatocellular carcinoma in hepatitis B patients; a population-based cohort study. Wang JP(1), Kao FY, Wu CY, ... Branch), Taipei, Taiwan. BACKGROUND: Hepatocellular carcinoma (HCC) is a major complication of hepatitis B virus (HBV) infection.

[A Taiwanese Nationwide Cohort Study Shows Interferon-Based ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4616715/> ▾ [翻译此页](#)

作者: YC Chen - 2015 - 被引用次数: 5 - 相关文章

2015年8月14日 - Hepatitis C virus (HCV) infection is a risk factor for chronic kidney disease (CKD). However, it remains unclear whether interferon-based therapy (IBT) for HCV was associated with reduced risk of CKD. From the Taiwan National Health Insurance Research Database, we identified 919 patients who received ...

[\[PDF\] Association Between Hepatitis B Virus and Chronic Kidney Disease: ...](#)

www.medigraphic.com/pdfs/hepato/ah-2017/ah171g.pdf ▾ [翻译此页](#)

作者: F Fabrizi - 被引用次数: 1 - 相关文章

Hepatitis B virus infection and chronic kidney disease are prevalent and remain a major public health problem world- are at increased risk for acquiring HBV. et al. Does nucleos(t)ide analogues treatment affect renal function in chronic hepatitis B patients who have already decreased eGFR? A longitudinal study.

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 4,870 条结果 (用时 0.73 秒)

Nucleos(t)ide analogues associated with a reduced risk of ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/25537961> - 翻译此页

作者: JP Wang - 2015 - 被引用次数: 12 - 相关文章

2014年12月23日 - Nucleos(t)ide analogues associated with a reduced risk of hepatocellular carcinoma in hepatitis B patients: a population-based cohort study. Wang JP(1), Kao FY, Wu CY ... Branch), Taipei, Taiwan. BACKGROUND: Hepatocellular carcinoma (HCC) is a major complication of hepatitis B virus (HBV) infection.

A nationwide cohort study suggests chronic hepatitis B virus infection ...

<https://www.ncbi.nlm.nih.gov/pubmed/25426815> - 翻译此页

作者: YC Chen - 2015 - 被引用次数: 18 - 相关文章

2014年11月26日 - A nationwide cohort study suggests chronic hepatitis B virus infection increases the risk of end-stage renal disease among patients in Taiwan. ... We identified 17,758 adults who had chronic HBV infection and had not taken nucleos(t)ide analogs from 1999 to 2010 and randomly selected 71,032 matched ...

A Taiwanese Nationwide Cohort Study Shows Interferon-Based ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4616715/> - 翻译此页

作者: YC Chen - 2015 - 被引用次数: 6 - 相关文章

2015年8月14日 - Hepatitis C virus (HCV) infection is a risk factor for chronic kidney disease (CKD). However, it remains unclear whether interferon-based therapy (IBT) for HCV was associated with reduced risk of CKD. From the Taiwan National Health Insurance Research Database, we identified 919 patients who received ...

[PDF] Association Between Hepatitis B Virus and Chronic Kidney Disease: ...

www.medigraphic.com/pdfs/hepato/ah-2017/ah171g.pdf - 翻译此页

作者: F Fabrizi - 被引用次数: 1 - 相关文章

Hepatitis B virus infection and chronic kidney disease are prevalent and remain a major public health problem world- are at increased risk for acquiring HBV. et al. Does nucleos(t)ide analogues treatment affect renal function in chronic hepatitis B patients who have already decreased eGFR? A longitudinal study.

Association Between Hepatitis B Virus and Chronic Kidney Disease: a ...

www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S1665... - 翻译此页

作者: F Fabrizi - 2017 - 被引用次数: 1 - 相关文章

Hepatitis B virus infection and chronic kidney disease are prevalent and remain a major public health