

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 37464

Manuscript type: ORIGINAL ARTICLE

Basic Study

Dysregulation of PARP1 is involved in the development of Barrett's esophagus

Zhang C *et al.* PARP1 in the development of Barrett's esophagus

Chao Zhang, Teng Ma, Tao Luo, Ang Li, Xiang Gao, Zhong-Gao Wang, Fei Li

Chao Zhang, Tao Luo, Ang Li, Xiang Gao, ³Zhong-Gao Wang, Fei Li,
Department of General Surgery, Xuanwu Hospital, Capital Medical
University, Beijing 100053, China

Teng Ma, Beijing Institute of Radiation Medicine, Beijing 100053, China

Match Overview

1	Internet 111 words crawled on 30-Jul-2015 bmbreports.org	3%
2	Internet 107 words crawled on 01-Jan-2018 gut.bmj.com	3%
3	Internet 84 words crawled on 16-Jun-2017 www.wjgnet.com	3%
4	Crossref 67 words Bo Qi, Yan Wang, Zhi-Jun Chen, Xiang-Nan Li, Yu Qi, Yang Yang, Guang-Hui Cui, Hai-Zhou Guo, Wei-Hao Li, Song Zh	2%
5	Internet 28 words crawled on 09-Jul-2016 mdpi.com	1%
6	Crossref 26 words X. Huo. "Deoxycholic Acid Causes DNA Damage while Inducing Apoptotic Resistance through NF- B Activation in Ber ...	1%
7	Crossref 26 words Pham, Thai H., Robert M. Genta, Stuart Jon Spechler, Rhonda F. Souza, and David H. Wang. "Development and Chara	1%
8	Internet 12 words crawled on 01-Oct-2016 cancerres.aacrjournals.org	<1%
9	Crossref 12 words "8th Annual Meeting of Chinese College of Surgeons and ... 9th Annual Meeting of the European Society of Surgery : Bei	<1%

获得 6 条结果 (用时 0.47 秒)

MicroRNA profiling of Barrett's oesophagus and oesophageal ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/20301167> ▼ 翻译此页

作者 : BP Wijnhoven - 2010 - 被引用次数 : 144 - 相关文章

The aberrant expression of microRNAs (miRNAs) is involved in the development of cancer. This study examined whether miRNAs play a role in the development of oesophageal adenocarcinoma. METHODS: RNA was extracted from mucosa of normal oesophageal squamous epithelium, normal gastric epithelium, Barrett's ...

缺少字词 : parp1

Novel Aberrations Uncovered in Barrett's Esophagus and Esophageal ...

<https://www.ncbi.nlm.nih.gov/pubmed/28751461> - 翻译此页

作者 : JLV Maag - 2017

2017年7月27日 - Novel Aberrations Uncovered in Barrett's Esophagus and Esophageal Adenocarcinoma Using Whole Transcriptome Sequencing. Maag JLV(1)(2), Fisher ... Barrett's esophagus (BE) is the main risk factor for EAC; however, the mechanisms driving EAC development remain poorly understood. Here ...

缺少字词 : parp1

Progression of Barrett's esophagus toward esophageal ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5198232/> ▼ 翻译此页

作者 : N Schoofs - 2017 - 相关文章

2016年9月30日 - In this review we will focus on the known risk factors for this progression, as well as the molecular pathways involved. We searched ... This is most likely to be attributable to the higher incidence of GERD in this population, since this is a well-known causal factor for the development of BE. The incidence of ...

缺少字词 : dysregulation parp1

MicroRNAs, development of Barrett's esophagus, and progression to ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2816263/> ▼ 翻译此页

作者 : CM Smith - 2010 - 被引用次数 : 50 - 相关文章

2010年2月7日 - Key areas of interest for miRNAs in Barrett's esophagus include their involvement in transdifferentiation and its links with the development of columnar metaplasia, and also cancer development within Barrett's esophagus. miRNAs are useful biomarkers for tumor classification, and their expression pattern ...

缺少字词 : dysregulation parp1

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 7,980 条结果 (用时 0.42 秒)

MicroRNA profiling of Barrett's oesophagus and oesophageal ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/20301167> ▼ [翻译此页](#)

作者: BP Wijnhoven - 2010 - 被引用次数: 144 - [相关文章](#)

The aberrant expression of microRNAs (miRNAs) is involved in the development of cancer. This study examined whether miRNAs play a role in the development of oesophageal adenocarcinoma.

METHODS: RNA was extracted from mucosa of normal oesophageal squamous epithelium, normal gastric epithelium, Barrett's ...

缺少字词: `parp4`

Novel Aberrations Uncovered in Barrett's Esophagus and Esophageal ...

<https://www.ncbi.nlm.nih.gov/pubmed/28751461> - [翻译此页](#)

作者: JLV Maag - 2017

2017年7月27日 - Novel Aberrations Uncovered in Barrett's Esophagus and Esophageal

Adenocarcinoma Using Whole Transcriptome Sequencing. Maag JLV(1)(2), Fisher ... Barrett's

esophagus (BE) is the main risk factor for EAC; however, the mechanisms driving EAC development remain poorly understood. Here ...

缺少字词: `parp4`

Biology of Barrett's Esophagus and Esophageal Adenocarcinoma

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3052949/> ▼ [翻译此页](#)

作者: DH Wang - 2011 - 被引用次数: 49 - [相关文章](#)

跳到Key Growth Regulatory Pathways that Contribute to Carcinogenesis - Although the data are inconclusive, it appears that Rb itself is targeted for inactivation in the latter stages of Barrett's

carcinogenesis (i.e. dysplasia and carcinoma), but not in non-dysplastic Barrett's metaplasia.

Inactivation of Rb is not the only ...

缺少字词: `parp4`

MicroRNAs, development of Barrett's esophagus, and progression to ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2816263/> ▼ [翻译此页](#)

作者: CM Smith - 2010 - 被引用次数: 50 - [相关文章](#)

2010年2月7日 - Key areas of interest for miRNAs in Barrett's esophagus include their involvement in transdifferentiation and its links with the development of columnar metaplasia, and also cancer