

Name of Journal: *World Journal of Nephrology*

Manuscript NO: 37509

Manuscript Type: Original Article

Basic Study

Genetic defects in ciliary genes in autosomal dominant polycystic kidney disease

Skalická K *et al.* Ciliary genes in polycystic kidney disease

Katarína Skalická, Gabriela Hřčková, Anita Vaská, Ágnes Baranyaiová, László Kovács

Abstract

AIM

To evaluate the genetic defects of ciliary genes causing the loss of primary cilium in autosomal dominant polycystic kidney disease (ADPKD).

Match Overview

1	Internet 129 words crawled on 12-Nov-2017 link.springer.com	4%
2	Internet 77 words crawled on 17-Nov-2016 www.omicsonline.org	2%
3	Crossref 26 words Ma, Ming, Xin Tian, Peter Igarashi, Gregory J Pazour and Stefan Somlo "Loss of cilia suppresses	1%
4	Crossref 25 words Bu, F., T. Maga, N. C. Meyer, K. Wang, C. P. Thomas C. M. Nester and R. J. H. Smith "Compreh	1%
5	Internet 20 words crawled on 30-Nov-2017 www.ncbi.nlm.nih.gov	1%
6	Internet 18 words crawled on 22-Jul-2017 www.wjgnet.com	<1%
7	Crossref 16 words Newton, Richard, and Lorenz Wernisch: "A Meta-Analysis of Multiple Matched Conv Number ar	<1%
8	Internet 16 words crawled on 19-Jun-2017 orca.cf.ac.uk	<1%
9	Internet 14 words crawled on 29-Oct-2003 www.grupoaulamedica.com	<1%
10	Internet 14 words crawled on 20-Sep-2017 eprints.gla.ac.uk	<1%

Autosomal Dominant Polycystic Kidney Disease Induced by Ciliary ...

<https://www.ncbi.nlm.nih.gov/books/NBK373370/>

Jump to **PKD** and **ciliary defects** - Among the known ciliopathies, **PKD** is one of the most common **renal genetic disorders**. Recent studies have demonstrated that mutations in **ciliary genes** are closely related to the onset of cystic **kidneys** (12). Therefore, cilia-defective mouse models were produced to elucidate ...

Autosomal Dominant Polycystic Kidney Disease Induced by Ciliary ...

<https://www.ncbi.nlm.nih.gov/pubmed/27512773>

by X Li - 2015

Autosomal dominant polycystic kidney disease (ADPKD) is a common **genetic** renal disease, which is caused by pathogenic mutations of either PKD1 (85%) or PKD2 (15%) genes, encoding for ... Also, animal models targeting **ciliary genes** show abnormal phenotypes such as polycystic kidneys and developmental **defects**.

Cyst growth, polycystins, and primary cilia in autosomal dominant ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4714135/> ▼

by SH Lee - 2014 - Cited by 17 - Related articles

Jump to **ADPKD genes** and proteins – PKD1, PKD2, and PC1 and PC2 - Functional **defects** in this complex caused by mutation of PKD1 or PKD2 result in **autosomal dominant polycystic kidney disease** (ADPKD) [7], [8]. ... Recent studies have generated novel information regarding the **genetic** and molecular implications of ADPKD, its pathogenesis, and new potential strategies for targeted treatment.

[Abstract](#) · [Introduction](#) · [PC1 dosage in ADPKD](#) · [Novel inter-relationship](#) ...

Mouse models of polycystic kidney disease induced by defects of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4133850/> ▼

by JY Ko - 2013 - Cited by 16 - Related articles

Polycystic kidney disease (PKD) is a common **genetic disorder** and is manifested by **defects** in **renal** epithelial cells, resulting in an increase of cell proliferation and cyst formation (1). **PKD** is a systemic **disorder** which is characterized by cysts in the kidney, liver and pancreas (2).



文章

找到约 5,410 条结果 (用时0.10秒)

时间不限

2018以来

2017以来

2014以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☐ 包括专利☐ 包含引用☒ 创建快讯

seek68文献馆--各种中外期刊数据库!

不限IP | 包月下载 | 可开发票 | 九年金牌服务

您是不是要找: Genetic defects in *cilia* genes in autosomal dominant polycystic kidney disease

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[HTML] The Caenorhabditis elegans **autosomal dominant polycystic kidney disease gene** homologs lov-1 and pkd-2 act in the same pathway
[MM Barr](#), J DeModena, D Braun, CQ Nguyen, DH Hall... - Current Biology, 2001 - Elsevier
... or incompletely rescues (two out of six stable lines) pkd-2(sy606Δ) response **defects** (Table 2 ...
encodes the C. elegans homolog of Tg737, a murine **autosomal** recessive PKD **gene**, and is ... Our
results provide direct **genetic** evidence that the C. elegans polycystins, LOV-1 and PKD ...
☆ 99 被引用次数: 256 相关文章 所有 10 个版本

[HTML] sciencedirect.com

Genetics and pathogenesis of polycystic kidney disease
P Igarashi, S Somlo - Journal of the American Society of ..., 2002 - Am Soc Nephrol
... By **birth**, the **kidneys** are massively replaced with cysts ... Although the two-hit hypothesis explains
many features of the **disease**, other **genetic** mechanisms, such as haploinsufficiency or ...
Polycystin-1, the product of the PKD1 **gene**, contains 4302 amino acids and has a molecular ...
☆ 99 被引用次数: 541 相关文章 所有 5 个版本

[HTML] asnjournals.org

Polycystin-2 localizes to **kidney cilia** and the **ciliary** level is elevated in orpk mice
with **polycystic kidney disease**
[GJ Pazour](#), JT San Agustín, JA Folliot, JL Rosenbaum... - Current Biology, 2002 - cell.com
... connection between PKD caused by **defects** in the polycystins, and PKD caused by **defects** in
the ... **Autosomal** recessive **polycystic kidney disease**: clinical and **genetic** profiles ... A **polycystic**
kidney-disease gene homologue required for male mating behaviour in C. elegans ...
☆ 99 被引用次数: 454 相关文章 所有 7 个版本

[PDF] cell.com

The **polycystic kidney disease** proteins, polycystin-1, polycystin-2, polaris, and
il **cilia**

[HTML] asnjournals.org

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 152,000 条结果 (用时 0.54 秒)

Google 学术: Genetic defects in ciliary genes in autosomal dominant polycystic kidney disease

... autosomal dominant polycystic kidney disease gene ... - Barr - 被引用次数: 256

Genetics and pathogenesis of polycystic kidney ... - Igarashi - 被引用次数: 541

... is elevated in orpk mice with polycystic kidney disease - Pazour - 被引用次数: 454

Autosomal Dominant Polycystic Kidney Disease Induced by Ciliary ...

<https://www.ncbi.nlm.nih.gov/pubmed/27512773> - 翻译此页

作者: X Li - 2015 - 被引用次数: 9 - 相关文章

Autosomal dominant polycystic kidney disease (ADPKD) is a common genetic renal disease, which is caused by pathogenic mutations of either PKD1 (85%) or PKD2 (15%) genes, encoding for ... Also, animal models targeting ciliary genes show abnormal phenotypes such as polycystic kidneys and developmental defects.

Cyst growth, polycystins, and primary cilia in autosomal dominant ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4714135/> - 翻译此页

作者: SH Lee - 2014 - 被引用次数: 18 - 相关文章

跳到ADPKD genes and proteins – PKD1, PKD2, and PC1 and PC2 - PKD2, the gene mutated in 15% of polycystic kidney disease cases, encodes PC2 [5], [27]. ... to renal epithelial proliferation and cyst growth in animal models and many human genetic diseases that are classified as ciliopathies [2], [3], [34], [35].

Autosomal Dominant Polycystic Kidney Disease Induced by Ciliary ...

<https://www.ncbi.nlm.nih.gov/books/NBK373370/> - 翻译此页

跳到PKD and ciliary defects - Among the known ciliopathies, PKD is one of the most common renal genetic disorders. Recent studies have demonstrated that mutations in ciliary genes are closely related to the onset of cystic kidneys [12]. Therefore, cilia defective mouse models were produced to elucidate the