

Name of journal: World Journal of Gastroenterology

Manuscript NO: 37585

Manuscript Type: RETROSPECTIVE STUDY

10
The prognostic value of lymph nodes count on survival of patients
with distal cholangiocarcinomas

Huapeng Lin et al. Prognostic value of retrieved LNs

Huapeng Lin^a, Shengwei Li^a, Ye Liu^b, Shiji Zhou^{c,*}

Abstract:

Match Overview

1	Crossref 16 words Peschaud Frédérique. "The Ratio of Metastatic to Exan ... ned Lymph Nodes Is a Powerful Independent Prognosti	<1%
2	Crossref 13 words Sanchuan Lai, Tingting Su, Xingkang He, Zhenghua Lin, Shujie Chen. "Prognostic value of resected lymph nodi ...	<1%
3	Publications 13 words "Higher continuity of statin use associated with decrea ... ed all-cause mortality.(News Capsules: WHAT'S", Form	<1%
4	Crossref 13 words Liu, Q., Z. Tan, P. Lin, H. Long, L. Zhang, T. Rong, Y. Men g, and G. Ma. "Impact of the number of resected lymph n	<1%
5	Internet 12 words crawled on 17-Jun-2017 f1000.com	<1%
6	Internet 12 words crawled on 23-Sep-2017 pure.uva.nl	<1%

[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 36,900 条结果 (用时 0.40 秒)

Prognostic impact of lymph node metastasis in distal ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/25611179> ▼ [翻译此页](#)

作者: M Kiriama - 2015 - 被引用次数: 39 - 相关文章

2015年1月22日 - The Cox proportional hazards model was used to identify independent predictors of survival. χ^2 scores were calculated to determine the cut-off value of the number of involved nodes, lymph node ratio (LNR) and total lymph node count (TLNC) for discriminating survival. RESULTS: Some 370 patients were ...

The prognostic value of lymph nodes count on survival of patients with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5190052/> ▼ [翻译此页](#)

作者: W Zheng - 2016 - 被引用次数: 7 - 相关文章

2016年6月6日 - The retrieved lymph node (LN) count has been validated as a prognostic factor in various cancers. However, the interaction between LN counts and patients' prognosis in gastric cancer with negative-LN metastasis is not fully studied. Surveillance, Epidemiology, and End Results Program (SEER)-registered ...

缺少字词: distal cholangiocarcinomas

Prognostic significance of lymph node ratio after resection of peri-hilar ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3081624/> ▼ [翻译此页](#)

作者: A Guglielmi - 2011 - 被引用次数: 45 - 相关文章



Prognostic value of lymph nodes count on survival of patients with distal



全部

图片

视频

新闻

购物

更多

设置

工具

找到约 36,700 条结果 (用时 0.41 秒)

Prognostic impact of lymph node metastasis in distal ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/25611179> ▼ 翻译此页

作者: M Kiriya - 2015 - 被引用次数: 41 - 相关文章

2015年1月22日 - The Cox proportional hazards model was used to identify independent predictors of survival. χ^2 scores were calculated to determine the cut-off value of the number of involved nodes, lymph node ratio (LNR) and total lymph node count (TLNC) for discriminating survival. RESULTS: Some 370 patients were ...

The prognostic value of lymph nodes count on survival of patients with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5190052/> ▼ 翻译此页

作者: W Zheng - 2016 - 被引用次数: 7 - 相关文章

2016年6月6日 - The retrieved lymph node (LN) count has been validated as a prognostic factor in various cancers. However, the interaction between LN counts and patients' prognosis in gastric cancer with negative-LN metastasis is not fully studied. Surveillance, Epidemiology, and End Results Program (SEER)-registered ...

缺少字词: distal cholangiocarcinomas

Prognostic significance of lymph node ratio after resection of peri-hilar ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3081624/> ▼ 翻译此页

作者: A Guglielmi - 2011 - 被引用次数: 45 - 相关文章

The aims of the present study were to evaluate, in a cohort of patients submitted to surgical resection with curative intent for perihilar cholangiocarcinoma, the relationship between lymph nodes status and survival, the prognostic significance of the total number of resected lymph nodes and to evaluate the prognostic ...

Prognostic impact of lymph node metastasis in distal ...

onlinelibrary.wiley.com/doi/10.1002/bjs.9752/abstract - 翻译此页

作者: M Kiriya - 2015 - 被引用次数: 39 - 相关文章

2015年1月22日 - Patients who underwent pancreaticoduodenectomy for distal cholangiocarcinoma in 24 hospitals in Japan between 2001 and 2010 were included. Survival was calculated by means of the Kaplan-Meier method and differences between subgroups were assessed with the log rank test. The