

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 37708

Manuscript Type: CASE REPORT

Obeticholic acid for severe bile acid diarrhoea with intestinal failure: case report

Christian L Hvas, Peter Ott, Peter Paine, Simon Lal, Søren Peter Jørgensen, Jens F Dahlerup

Abstract

Bile acid diarrhoea results from excessive amounts of bile acids that enter the colon following bile acid malabsorption in the terminal ileum. Main therapies

Match Overview

- | | | |
|---|--|----|
| 1 | Crossref 21 words
Walters, J. R. F., I. M. Johnston, J. D. Nolan, C. Vassie, M. E. Pruzanski, and D. A. Shapiro. "The response of patients | 1% |
| 2 | Crossref 12 words
Borghede, M.K.. "Bile acid malabsorption investigated by selenium-75-homocholeic acid taurine (⁷⁵ SeHCAT) scans: | 1% |

全部

新闻

购物

图片

地图

更多

设置

工具

找到约 4,220 条结果 (用时 0.60 秒)

The response of patients with bile acid diarrhoea to the farnesoid X ...

<https://www.ncbi.nlm.nih.gov/pubmed/25329562> ▼ 翻译此页

作者: JR Walters - 2015 - 被引用次数: 57 - 相关文章

2014年10月20日 - BACKGROUND: Bile acid diarrhoea is a common cause of chronic diarrhoea, occurring as a primary condition or secondary to ileal disease or ... AIM: To investigate whether obeticholic acid, a potent FXR agonist, could increase FGF19 in patients with bile acid diarrhoea, and produce clinical benefits.

缺少字词: failure case review literature

Bile acid malabsorption in chronic diarrhea: Pathophysiology and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3816948/> ▼ 翻译此页

作者: A Barkun - 2013 - 被引用次数: 35 - 相关文章

Bile acid malabsorption (BAM) is a common but frequently under-recognized cause of chronic diarrhea, with an estimated prevalence of 4% to 5%. ... review, the PubMed database was searched using a combination of controlled vocabulary and text words to identify reports related to bile acid diarrhea for the period October ...

缺少字词: obeticholic failure

Obeticholic Acid in Bile Acid Diarrhoea - Full Text View - ClinicalTrials ...

<https://clinicaltrials.gov/ct2/show/NCT01585025> ▼ 翻译此页

2012年4月25日 - This study is a pilot, proof-of-concept, open-label study to investigate whether OCA

找到约 4,230 条结果 (用时 0.66 秒)

The response of patients with bile acid diarrhoea to the farnesoid X ...

<https://www.ncbi.nlm.nih.gov/pubmed/25329562> ▼ 翻译此页

作者: JR Walters - 2015 - 被引用次数: 57 - 相关文章

2014年10月20日 - BACKGROUND: **Bile acid diarrhoea** is a common cause of **chronic** diarrhoea, occurring as a primary condition or secondary to ileal disease or ... AIM: To investigate whether **obeticholic acid**, a potent FXR agonist, could increase FGF19 in patients with **bile acid diarrhoea**, and produce clinical benefits.

缺少字词: *failure case review literature*

Obeticholic Acid in Bile Acid Diarrhoea - Full Text View - ClinicalTrials ...

<https://clinicaltrials.gov/ct2/show/NCT01585025> ▼ 翻译此页

2012年4月25日 - This **study** is a pilot, proof-of-concept, open-label **study** to investigate whether OCA can stimulate FGF19 in **bile acid diarrhoea** patients to provide a safe and effective treatment. ...

Primary **Bile Acid Malabsorption** Secondary **Bile Acid Malabsorption Chronic** Diarrhoea, Drug: **Obeticholic acid**, Phase 2 ...

缺少字词: *literature*

[PDF] 207999Orig1s000 - FDA

https://www.accessdata.fda.gov/drugsatfda_docs/nda/.../207999Orig1s000MedR.pdf ▼

2016年4月22日 - Clinical **Review**. Ruby Mehta. NDA 207999. OCALIVA [**Obeticholic acid** (OCA)].

CDER Clinical **Review** Template 2015 Edition. 1. Version date June 25 ... Primary **Biliary**

Cholangitis/Cirrhosis FDA **reviewed** the **case report** forms (CRFs) that were to be used for collecting the data for the Global PBC **study**.

[PDF] Gastrointestinal Drug Advisory Committee (GIDAC) Meeting - FDA

<https://www.fda.gov/.../GastrointestinalDrugsAdvisoryCommittee/UCM494108.pdf> ▼

2016年4月7日 - The **Obeticholic acid** (OCA) Phase 2 and Phase 3 clinical trials in Primary **Biliary**

Cirrhosis. The OCA PBC is a rare and potentially fatal disease leading to liver **failure** and the need for liver transplant to avoid ... FDA **reviewed** the **case report** forms (CRFs) that were to be used for collecting the data for the.