

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 38588

Manuscript Type: ORIGINAL ARTICLE

**mitochondrial pathway mediated by reactive oxygen species involvement
in α -Hederin-induced apoptosis in hepatocellular carcinoma cells**

Jiao Li, Dan-dan Wu, Ji-xiang Zhang, Jing-jing Ma, Xue Hu

Abstract

AIM

To investigate the antitumor activity of α -hederin in hepatocellular carcinoma (HCC) cells and its underlying mechanisms *in vitro* and *in vivo*.

Match Overview

There are no matching sources for this report.

文章

获得 10 条结果 (用时0.08秒)

时间不限

2018以来

2017以来

2014以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利☐ 包含引用☒ 创建快讯

您是不是要找: mitochondrial pathway mediated by reactive oxygen species involvement in hepatocellular carcinoma cells

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Intracellular glutathione depletion and reactive oxygen species generation are important in α -hederin-induced apoptosis of P388 cells

BTK Huat, SMK Swamy - Molecular and cellular biochemistry, 2003 - Springer

... Abundant evidence has indicated that caspase activation is a common pathway for the execution of apoptosis [50 ... Fulda S, Scaffidi C, Susin SA, Krammer PH, Kroemer G, Peter ME, Debatin KM: Activation of mitochondria and release of mitochondrial apoptogenic factors ...

☆ 99 被引用次数: 86 相关文章 所有 11 个版本

[HTML] The anticancer effect and mechanism of α -hederin on breast cancer cells

L Cheng, TS Xia, YF Wang... - International ..., 2014 - spandidos-publications.com

... Mitochondrial apoptotic pathway was reported widely for the actions of triterpenoid saponins in ... reduced the mitochondrial membrane potential and decreased mitochondrial Apaf-1 ... Consistent with these results, α -hederin induced mitochondria-mediated apoptosis of MCF-7 and ...

☆ 99 被引用次数: 20 相关文章 所有 9 个版本

Pentacyclic triterpenoids and their saponins with apoptosis-inducing activity

SR Wang, WS Fang - Current topics in medicinal chemistry, 2009 - ingentaconnect.com

... The C-21 amphipathic side chain is crucial to the channel formation in the mitochondrial outer membrane, which may ... groups in these saponins are crucial to their apoptotic activity, and 52 induced apoptosis via mitochondria-mediated intrinsic pathway, and caspase-9 ...

☆ 99 被引用次数: 32 相关文章 所有 4 个版本

Therapeutic implications of black seed and its constituent thymoquinone in the



文章

获得 10 条结果 (用时0.07秒)

时间不限

2018以来

2017以来

2014以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利☐ 包含引用☒ 创建快讯

您是不是要找: Mitochondrial pathway mediated by reactive oxygen species involvement in hepatocellular carcinoma cells

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Intracellular glutathione depletion and reactive oxygen species generation are important in α -hederin-induced apoptosis of P388 cells

BTK Huat, SMK Swamy - Molecular and cellular biochemistry, 2003 - Springer

... Abundant evidence has indicated that caspase activation is a common pathway for the execution of apoptosis [50 ... Fulda S, Scaffidi C, Susin SA, Krammer PH, Kroemer G, Peter ME, Debatin KM: Activation of mitochondria and release of mitochondrial apoptogenic factors ...

☆ 99 被引用次数: 88 相关文章 所有 11 个版本

[HTML] The anticancer effect and mechanism of α -hederin on breast cancer cells

L Cheng, TS Xia, YF Wang... - International ..., 2014 - spandidos-publications.com

... Mitochondrial apoptotic pathway was reported widely for the actions of triterpenoid saponins in ... reduced the mitochondrial membrane potential and decreased mitochondrial Apaf-1 ... Consistent with these results, α -hederin induced mitochondria-mediated apoptosis of MCF-7 and ...

☆ 99 被引用次数: 20 相关文章 所有 9 个版本

Pentacyclic triterpenoids and their saponins with apoptosis-inducing activity

SR Wang, WS Fang - Current topics in medicinal chemistry, 2009 - ingentaconnect.com

... The C-21 amphipathic side chain is crucial to the channel formation in the mitochondrial outer membrane, which may ... groups in these saponins are crucial to their apoptotic activity, and 52 induced apoptosis via mitochondria-mediated intrinsic pathway, and caspase-9 ...

☆ 99 被引用次数: 33 相关文章 所有 4 个版本

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 2,900 条结果 (用时 0.59 秒)

Google 学术: Mitochondrial pathway mediated by reactive oxygen species involvement in α -hederin-induced apoptosis in hepatocellular carcinoma cells

... important in α -hederin-induced apoptosis of P388 cells - Huat - 被引用次数: 88

... and mechanism of α -hederin on breast cancer cells - Cheng - 被引用次数: 20

... triterpenoids and their saponins with apoptosis- ... - Wang - 被引用次数: 33

Hesperetin induces the apoptosis of hepatocellular carcinoma cells ...

<https://www.ncbi.nlm.nih.gov/pubmed/25737432> - 翻译此页

作者: J Zhang - 2015 - 被引用次数: 12 - 相关文章

Med Oncol. 2015 Apr;32(4):101. doi: 10.1007/s12032-015-0516-z. Epub 2015 Mar 4. Hesperetin induces the apoptosis of hepatocellular carcinoma cells via mitochondrial pathway mediated by the increased intracellular reactive oxygen species, ATP and calcium. Zhang J(1), Song J, Wu D, Wang J, Dong W. 缺少字词: involvement α hederin

PubMed Result - NCBI

https://www.ncbi.nlm.nih.gov/pubmed?linkname=pubmed_pubmed... - 翻译此页

Intracellular glutathione depletion and reactive oxygen species generation are important in α -hederin-induced apoptosis of P388 cells. Mol Cell Biochem. 2003 Mar;245(1-2):127-39. PubMed PMID: 12708752. 2: Liu J, Shen HM, Ong CN. Role of intracellular thiol depletion, mitochondrial dysfunction and reactive oxygen ...

Carnosic acid induces apoptosis of hepatocellular carcinoma cells via ...

<https://www.ncbi.nlm.nih.gov/pubmed/28918123> - 翻译此页

作者: X Zhang - 2017 - 被引用次数: 3 - 相关文章

2017年9月13日 - Carnosic acid induces apoptosis of hepatocellular carcinoma cells via ROS-mediated mitochondrial pathway. ... Specifically, CA led to a decreased mitochondrial membrane potential (MMP) and increases in intracellular reactive oxygen species (ROS) levels and apoptosis-related protein expression.

缺少字词: involvement α hederin