

World Journal of *Gastroenterology*

World J Gastroenterol 2018 August 14; 24(30): 3313-3468





REVIEW

- 3313 MicroRNAs as non-invasive diagnostic biomarkers for gastric cancer: Current insights and future perspectives
Link A, Kupcinskas J
- 3330 Nutritional issues in patients with obesity and cirrhosis
Schiavo L, Busetto L, Cesaretti M, Zelber-Sagi S, Deutsch L, Iannelli A
- 3347 Host genetic factors affecting hepatitis B infection outcomes: Insights from genome-wide association studies
Akçay IM, Katrinli S, Ozdil K, Dinler Doganay G, Doganay L

MINIREVIEWS

- 3361 Current guidelines for the management of non-alcoholic fatty liver disease: A systematic review with comparative analysis
Leoni S, Tovoli F, Napoli L, Serio I, Ferri S, Bolondi L
- 3374 Form confers function: Case of the 3'X region of the hepatitis C virus genome
Dutkiewicz M, Ciesiolka J

ORIGINAL ARTICLE

Basic Study

- 3384 Herb-partitioned moxibustion alleviates colon injuries in ulcerative colitis rats
Zhang D, Ren YB, Wei K, Hong J, Yang YT, Wu LJ, Zhang J, Shi Z, Wu HG, Ma XP
- 3398 Novel sericin-based hepatocyte serum-free medium and sericin's effect on hepatocyte transcriptome
Huang Y, Peng Q, Li HY, Jia ZD, Li Y, Gao Y
- 3414 Total flavone of *Abelmoschus manihot* suppresses epithelial-mesenchymal transition *via* interfering transforming growth factor- β 1 signaling in Crohn's disease intestinal fibrosis
Yang BL, Zhu P, Li YR, Xu MM, Wang H, Qiao LC, Xu HX, Chen HJ
- 3426 Identification of a five-long non-coding RNA signature to improve the prognosis prediction for patients with hepatocellular carcinoma
Zhao QJ, Zhang J, Xu L, Liu FF
- 3440 Application of modified primary closure of the pelvic floor in laparoscopic extralevator abdominal perineal excision for low rectal cancer
Wang YL, Zhang X, Mao JJ, Zhang WQ, Dong H, Zhang FP, Dong SH, Zhang WJ, Dai Y

Retrospective Study

**Observational Study**

- 3448** Altered oral microbiota in chronic hepatitis B patients with different tongue coatings

Zhao Y, Mao YF, Tang YS, Ni MZ, Liu QH, Wang Y, Feng Q, Peng JH, Hu YY

CASE REPORT

- 3462** Large heterotopic gastric mucosa and a concomitant diverticulum in the rectum: Clinical experience and endoscopic management

Chen WG, Zhu HT, Yang M, Xu GQ, Chen LH, Chen HT

Contents

World Journal of Gastroenterology
Volume 24 Number 30 August 14, 2018

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Herwig Cerwenka, MD, Professor, Department of Surgery, Medical University of Graz, Graz A-8036, Austria

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2018 edition of Journal Citation Reports® cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35th among 80 journals in gastroenterology and hepatology (quartile in category Q2).

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Shu-Yu Yin*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Rao-Yu Ma*
Proofing Editorial Office Director: *Ze-Mao Gong*

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski, MD, PhD, DSc (Med),
Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach, CA 90822, United States

EDITORIAL BOARD MEMBERS

All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE

Ze-Mao Gong, Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER

Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE

August 14, 2018

COPYRIGHT

© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS

Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION

<http://www.f6publishing.com>

Basic Study

Identification of a five-long non-coding RNA signature to improve the prognosis prediction for patients with hepatocellular carcinoma

Qiu-Jie Zhao, Jiao Zhang, Lin Xu, Fang-Feng Liu

Qiu-Jie Zhao, Jiao Zhang, Lin Xu, Department of Gastroenterology, Shandong Provincial Hospital affiliated to Shandong University, Jinan 250021, Shandong Province, China

Fang-Feng Liu, Department of Hepatobiliary Surgery, Shandong Provincial Hospital affiliated to Shandong University, Jinan 250021, Shandong Province, China

ORCID number: Qiu-Jie Zhao (0000-0003-4053-8406); Jiao Zhang (0000-0002-9178-1003); Lin Xu (0000-0003-0253-608X); Fang-Feng Liu (0000-0002-3745-5012).

Author contributions: Zhao QJ conceived the study, drafted and revised the manuscript; Zhang J and Xu L helped with the statistical analysis; Liu FF helped participated in data mining; all authors read and approved the final manuscript.

Supported by the National Nature Science Foundation of China, No. 81702816 (to Zhao QJ); and Shandong Provincial Natural Science Foundation, No. ZR2017PH030 (to Zhao QJ).

Institutional review board statement: This studied mined the TCGA database and doesn't involve any experiments with animals or human beings. Because the TCGA data are a community resource project, additional ethical approval was not acquired.

Conflict-of-interest statement: The authors declare that there is no conflict of interest related to this study.

Data sharing statement: The datasets supporting the conclusions of this article are included within the article.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

licenses/by-nc/4.0/

Manuscript source: Unsolicited manuscript

Correspondence to: Qiu-Jie Zhao, PhD, Chief Doctor, Department of Gastroenterology, Shandong Provincial Hospital affiliated to Shandong University, No. 324, Jingwuwei 7th Road, Jinan 250021, Shandong Province, China. zhaoqijiesdu@126.com
Telephone: +86-531-68772952

Received: March 27, 2018

Peer-review started: March 28, 2018

First decision: May 9, 2018

Revised: May 24, 2018

Accepted: June 22, 2018

Article in press: June 22, 2018

Published online: August 14, 2018

Abstract

AIM

To construct a long non-coding RNA (lncRNA) signature for predicting hepatocellular carcinoma (HCC) prognosis with high efficiency.

METHODS

Differentially expressed lncRNAs (DELs) between HCC specimens and peritumor liver specimens were identified using the edgeR package to analyze The Cancer Genome Atlas (TCGA) LIHC dataset. Univariate Cox proportional hazards regression was performed to obtain the DELs significantly associated with overall survival (OS) in a training set. These OS-related DELs were further analyzed using a stepwise multivariate Cox regression model. Those lncRNAs fitted in the multivariate Cox regression model and independently associated with overall survival were chosen to build a prognostic risk formula. The prognostic value of

this formula was then validated in the test group and the entire cohort and further compared with two previously identified prognostic signatures for HCC. Gene ontology and Kyoto Encyclopedia of Genes and Genomes pathway analyses were performed to explore the potential biological functions of the lncRNAs in the signature.

RESULTS

Based on lncRNA expression profiling of 370 HCC patients from the TCGA database, we constructed a 5-lncRNA signature (AC015908.3, AC091057.3, TMCC1-AS1, DCST1-AS1 and FOXD2-AS1) that was significantly associated with prognosis. HCC patients with high-risk scores based on the expression of the 5 lncRNAs had significantly shorter survival times compared to patients with low-risk scores in both the training and test groups. Multivariate Cox regression analysis demonstrated that the prognostic value of the 5 lncRNAs was independent of clinicopathological parameters. A comparison study involving two previously identified prognostic signatures for HCC demonstrated that this 5-lncRNA signature showed improved prognostic power compared with the other two signatures. Functional enrichment analysis indicated that the 5 lncRNAs were potentially involved in metabolic processes, fibrinolysis and complement activation.

CONCLUSION

Our present study constructed a 5-lncRNA signature that improves survival prediction and can be used as a prognostic biomarker for HCC patients.

Key words: Long non-coding RNA; Hepatocellular carcinoma; Prognosis; Survival prediction; Prognostic biomarker

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: In the present study, we developed a 5-long non-coding RNA (lncRNA) signature for predicting the prognosis of hepatocellular carcinoma (HCC) patients based on The Cancer Genome Atlas database. The signature was reproducible and robust in another independent large-scale HCC cohort, supporting its utility and effectiveness. In addition, the prognostic value of the 5-lncRNA signature was independent of clinicopathological variables. When compared with two previously identified signatures for HCC survival prediction, this 5-lncRNA signature showed superior prognostic power. Our study indicates that the 5-lncRNA signature could improve survival prediction and could be used as a prognostic biomarker for HCC patients.

Zhao QJ, Zhang J, Xu L, Liu FF. Identification of a five-long non-coding RNA signature to improve the prognosis prediction for patients with hepatocellular carcinoma. *World J Gastroenterol* 2018; 24(30): 3426-3439 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v24/i30/3426.htm>

INTRODUCTION

Hepatocellular carcinoma (HCC) is the sixth most commonly diagnosed cancer in the world^[1]. According to previous epidemiologic studies, the incidence of HCC varies strikingly worldwide and is particularly high in eastern Asian countries, including China, and sub-Saharan Africa^[2,3]. The 5-year overall survival rate for HCC is lower than 20%^[4], and the ratio of its mortality to morbidity is 0.95^[1]. Because of its poor prognosis, HCC ranks as the second leading cause of cancer-related deaths worldwide^[1]. An estimated 782500 new liver cancer cases and 745500 deaths occurred worldwide in 2012, among which 50% occurred in China^[3]. There are multiple risk factors related to HCC, including hepatitis B or C viral infection, chronic alcohol abuse, nonalcoholic fatty liver disease and smoking^[5,6]. Although treatment for HCC, including surgical resection, has improved over the past decades, the overall survival rate for this disease remains devastatingly high due to its high recurrence rate (50%-70% at 5 years)^[7-9]. Because HCC is a heterogeneous disease with substantially variable clinical outcomes, the search for effective biomarkers to predict recurrence and prognosis is indispensable. To date, no widely accepted molecular biomarkers for HCC aggressiveness are available. In the past 40 years, serum alpha fetoprotein (AFP) levels have been utilized for the diagnosis of HCC and for predicting its response to therapy. However, AFP levels can be influenced by tumor size and cancer stage, and they are not reliable in clinical applications^[10]. In addition, the American Association for the Study of Liver Diseases concluded that the use of AFP levels lacks sufficient sensitivity and specificity to effectively monitor or diagnose HCC^[11].

With the development of high-throughput sequencing technologies, it has become easy to acquire whole genome profiles for specific cancers and develop more reliable prognostic signatures. Long non-coding RNAs (lncRNAs) are mRNA-like transcripts of more than 200 nucleotides (nt) with little or no protein-coding capacity^[12,13]. In the past, they were previously thought to be redundant segments of the genome, but in recent decades, emerging studies have indicated the importance of lncRNAs in cellular physiological and pathological processes^[14,15]. Increasing evidence suggests that dysregulated lncRNAs are associated with various human diseases, particularly the initiation and progression of various human cancers^[16,17]. Prognostic lncRNA signatures have been examined in many cancer types, including renal cancer, glioblastoma, colorectal cancer, lymphoma, and others^[18-21]. For HCC, most of the published gene signatures associated with prognosis have focused on mRNAs and microRNAs^[22-25]. To the

Table 1 Clinicopathological parameters of hepatocellular carcinoma patients in each cohort

Variables		Training group (n = 184)	Test group (n = 186)	Entire group (n = 370)
Age, yr	< 60	84	85	169
	≥ 60	100	101	201
Sex	Male	129	120	249
	Female	55	66	121
Weight, kg	< 70	92	85	177
	≥ 70	78	89	167
	NA	14	12	26
Child-Pugh grade	A	109	107	216
	B	9	12	21
	C	0	1	1
Fibrosis ishak score	NA	66	66	132
	0	32	42	74
	1-5	38	31	69
	6	36	33	69
Vascular tumor invasion	NA	78	80	158
	None	100	106	206
	Micro	45	46	91
	Macro	8	9	17
Serum FAP level, ng/mL	NA	31	25	56
	< 100	102	90	192
	≥ 100	37	48	85
	NA	45	48	93
Tumor grade	1	37	18	55
	2	82	95	177
	3 + 4	61	72	133
	NA	4	1	5
Pathologic stage	I	92	79	171
	II	43	42	85
	III + IV	37	53	90
	NA	12	12	24

best of our knowledge, very few lncRNA signatures have been developed for HCC prognosis prediction^[26]. Thus, it is necessary to identify a more effective lncRNA signature for HCC prognosis. In the present study, we aimed to construct a lncRNA signature capable of predicting HCC prognosis with high efficiency.

In this work, we analyzed a cohort of 370 HCC patients from The Cancer Genome Atlas (TCGA) to identify a potential lncRNA signature for predicting the survival of HCC patients. We identified a five-lncRNA prognostic signature from the TCGA dataset and determined that its prognostic value was independent from clinical factors. The identification of prognostic lncRNAs suggests the potential roles of lncRNAs in HCC pathogenesis and progression.

MATERIALS AND METHODS

Data and patients

Level 3 RNA-seq data (HTSeq-counts) from 374 HCC tumor specimens and 50 peritumoral liver specimens and their corresponding clinicopathological information were downloaded from the TCGA project (<https://cancergenome.nih.gov/>) on June 2, 2017. Because TCGA data are a community resource project, additional ethical approval was not acquired, and the present study adhered to TCGA publication guidelines and data access policies. After excluding the data without complete survival information, a total of 370 HCC

patients with complete follow up data were enrolled in our study and then randomly divided into a training set ($n = 184$) and test set ($n = 186$) using SPSS software (version 24.0). The clinicopathological parameters of the HCC patients in each group are listed in Table 1.

lncRNA expression profile in the TCGA LIHC cohort

Only lncRNAs with a description in NCBI or Ensemble were selected for further study in this paper. We obtained the expression profiles of 6929 lncRNAs from the RNA-seq data of the TCGA LIHC cohort. Differentially expressed lncRNAs (DELs) between the HCC specimens and peritumor liver specimens were identified with the edgeR package, using an adjusted $P < 0.05$ and \log_2 |fold change| > 1 . The expression level of each lncRNA was \log_2 transformed for the downstream analyses.

Identification of prognostic lncRNAs and construction of the risk formula for overall survival prediction

Univariate Cox proportional hazards regression was performed to obtain the DELs that were significantly associated with the overall survival (OS) of HCC patients in the training group. After acquiring survival-related lncRNAs ($P < 0.01$), we excluded those not expressed in at least 10% of the samples. The remaining OS-related lncRNAs were then adjusted using the stepwise multivariate Cox regression model. Finally, those lncRNAs fitted in the multivariate Cox regression model and independently associated with OS were chosen. A




Figure 1 Volcano plot of the differentially expressed long non-coding RNAs between hepatocellular carcinoma tumor specimens and peritumoral liver specimens. The X-axis represents the adjusted FDR, and the Y-axis represents the value of the \log_2 fold change. Aberrantly expressed long non-coding RNAs (lncRNAs) were calculated using the EdgeR package. Red dots represent upregulated lncRNAs in the hepatocellular carcinoma (HCC) tumor specimens, while green dots indicate downregulated lncRNAs compared with the peritumoral liver specimens. Black dots show the lncRNAs without significant differences between the HCC tumor and peritumoral liver specimens. Altogether, 2240 upregulated and 353 downregulated lncRNAs were found. This volcano plot was conducted using the ggplot2 package of R language.

prognostic risk formula was established based on a linear combination of the expression level of these lncRNAs multiplied by the regression coefficient derived from the multivariate Cox regression model as previously described^[18-21]. The subjects in each dataset were classified into a high-risk group and low-risk group according to the median risk score of the risk formula derived from the training set.

Statistical analysis

Univariate Cox proportional hazards regression was performed to obtain survival-related DELs, and the stepwise multivariate Cox regression model was performed for further selection. Overall survival analyses in the high-risk and low-risk groups were performed using Kaplan-Meier survival curves and a log-rank test. Receiver operating curve analyses were performed to assess the specificity and sensitivity of the prognosis prediction. The above analyses were performed using R (version 3.3.1). To verify the independence of the prognostic value of the 5-lncRNA signature and clinicopathological parameters, univariate and multivariate Cox regression analyses were performed using SPSS software (version 24.0). In the comparison study, Kaplan-Meier survival analysis and receiver operating curve (ROC) analysis were also performed using SPSS (version 24.0).

Functional enrichment analyses

To identify co-expressed lncRNA-mRNA pairs, we performed Person correlation analyses with R (version

3.3.1) for each of the five lncRNAs with protein-coding genes based on the RNA-seq data of the TCGA LIHC cohort. The protein-coding genes with a correlation coefficient > 0.5 and a $P < 0.01$ were considered to be significantly correlated genes. For functional enrichment analysis, the correlated protein-coding genes were subjected to gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analyses using DAVID Bioinformatics Resources (version 6.8)^[27,28]. Significant functional categories were identified and limited to GO terms in the "Biological Process" (GOTERM-BP-DIRECT) and KEGG pathway categories, using the human whole genome as the background. Significantly enriched GO terms with similar functions were visualized using the EnrichmentMap plugin in Cytoscape (version 3.5.1)^[29].

RESULTS

Determining prognostic lncRNAs from the training set

Using the edgeR package, we identified a total of 2593 lncRNAs differentially expressed ($|\log_2|\text{fold change}| > 1$ and adjusted $P < 0.05$) between 374 HCC tumor specimens and 50 peritumor liver specimens, including 2240 upregulated and 353 downregulated lncRNAs (Figure 1). A total of 370 HCC samples with complete survival information were subjected to further analyses. For the training set, univariate Cox proportional hazards regression analyses revealed 82 lncRNAs significantly correlated with OS ($P < 0.01$) among the 2593 differentially expressed lncRNAs. Among the 82 OS-related lncRNAs, we further excluded those expressed in less than 10% of the HCC specimens, and the remaining 30 lncRNAs were subjected to further selection.

Construction of a lncRNA-based prognostic signature and validation in the training group

Stepwise multivariable Cox proportional hazards regression analyses were performed to identify the optimal prognostic lncRNAs among the 30 candidate lncRNAs. Based on this model, a final 5 lncRNAs were found to be significantly and independently related to prognosis. We then constructed a prognostic signature based on the expression levels of these 5 lncRNAs and their coefficients derived from the multivariable Cox model. The formula is as follows: risk score = $(-0.1900 \times \text{the expression level of AC015908.3}) + (0.1764 \times \text{the expression level of FOXD2-AS1}) + (0.3588 \times \text{the expression level of AC091057.3}) + (0.5615 \times \text{the expression level of TMCC1-AS1}) + (0.4877 \times \text{the expression level of DCST1-AS1})$. Detailed information for the 5 lncRNAs is listed in Table 2. The risk score for each patient in the training group was calculated using the formula. The training set was then divided into a high-risk group ($n = 92$) and a low-risk group ($n = 92$) according to the median risk score. Kaplan-Meier analysis revealed that the high-risk group had a significantly poorer prognosis than that of the low-

Table 2 The 5 long non-coding RNAs significantly associated with overall survival of hepatocellular carcinoma patients

Gene ID	Gene symbol	Coefficient	Hazard ratio	P value
ENSG00000264016	AC015908.3	-0.1900	0.7792	0.000305
ENSG00000237424	FOXD2-AS1	0.1764	1.2865	0.007317
ENSG00000269974	AC091057.3	0.3588	1.4682	0.000375
ENSG00000271270	TMCC1-AS1	0.5615	1.5417	0.000287
ENSG00000232093	DCST1-AS1	0.4877	1.3909	0.001632

risk group ($P = 1.3\text{e-}09$, log-rank test, Figure 2A). The median survival time for the high-risk group and the low-risk group was 2.096 and 6.811 years, respectively. Additionally, the 3- and 5-year survival rates of the high-risk group were 40% and 23.5%, whereas the corresponding survival rates were 90% and 71.8%, respectively, in the low-risk group. To evaluate the performance of the 5-lncRNA signature for predicting the prognosis of HCC patients, a time-dependent ROC analysis was conducted. The area under the ROC curve (AUC) for the 5-lncRNA signature was 0.857, which indicated good performance (Figure 2B). The risk scores of patients in the training group were also ranked, and survival status was plotted for each patient on a dot plot (Figure 2C). The mortality for patients in the high-risk group was much higher than that in the low-risk group. A heat map displays the expression profiles of these five lncRNAs in the samples from the training group; the expression profiles are ranked according to risk score (Figure 2D). Among the 5 lncRNAs, AC015908.3 showed a negative coefficient derived for the multivariate Cox regression model and seemed to be a protective factor, as its high expression predicts a low risk. The other 4 lncRNAs with positive coefficients, including FOXD2-AS1, AC091057.3, TMCC1-AS1 and DCST1-AS1, seemed to be risk factors and all were upregulated in the high-risk group compared to the low-risk group within the training set.

Validation of the prognostic value of the 5-lncRNA signature for the test set and the entire cohort

To further verify the prognostic value of the 5-lncRNA signature for HCC patients, risk scores for patients in the test group were calculated according to the constructed formula based on the expression of the 5 lncRNAs. The test group was also divided into high-risk ($n = 97$) and low-risk ($n = 89$) groups using the same cutoff as for the training group. Kaplan-Meier analysis revealed that the survival rate of the high-risk subgroup was much lower than that of the low-risk subgroup in the test set (median OS: 2.293 years vs 8.562 years; log-rank $P = 1.64\text{e-}05$) (Figure 3A). For the entire set, a similar result was obtained by Kaplan-Meier analysis. Among the entire cohort, the median survival of the high-risk group ($n = 189$) was 2.197 years, which was significantly lower than the median OS of 6.937 years for the low-risk group ($n = 181$) ($P = 2.69\text{e-}13$, Figure 4A). The AUC for the 5-lncRNA-based risk score of overall survival was 0.709 and 0.769 for

the test group (Figure 3B) and the entire group (Figure 4B), respectively, with both showing robust utility. In addition, ranked risk scores and survival status for each subject were plotted for the test group (Figure 3C) and the entire set (Figure 4C). Heatmaps display the expression profiles of the five lncRNAs for each subject in the test group (Figure 3D) and the entire cohort (Figure 4D), which were ranked according to risk score.

The prognostic value of the 5-lncRNA signature was independent of clinical characteristics

Univariate and multivariate Cox regression analyses were performed with the 5-lncRNA-based risk score and clinicopathological factors, including age, gender, weight, Child-Pugh grading, fibrosis extent, vascular tumor invasion, serum FAP levels, tumor grade and pathological stage as explanatory variables and overall survival as the dependent variable. The univariate Cox regression demonstrated that the 5-lncRNA signature-based risk score and pathologic stage were able to effectively predict the prognosis of HCC patients. In addition, in the training set and the entire set, patient age seemed to be related to survival, although this did not reach significance (Table 3). In contrast, none of the other clinicopathological parameters were associated with prognosis in either set. Multivariate Cox regression analysis revealed that after adjusting for other factors, age (only for the entire set), pathologic stage and the 5-lncRNA signature were the only factors significantly associated with overall survival (Table 2). Patients from the entire cohort were then stratified by age (Figure 5A) and pathological stage (Figure 5B). Each subgroup was then divided into a high-risk and low-risk group based on the 5-lncRNA risk score median derived from the training group. Kaplan-Meier analysis revealed that for all of the subgroups, the high-risk group had significantly poorer survival than the low-risk group. All of these results strongly suggest that the prognostic value of the 5-lncRNA-based risk score is independent of clinicopathological factors.

Comparison of the 5-lncRNA signature with existing prognostic signatures for HCC

Two HCC-related prognostic signatures have recently been developed and reported, including a 3-gene signature by Binghua Li and a 4-lncRNA signature by Zhonghao Wang that were both derived from the TCGA dataset^[25,26]. To compare the prognostic value of the 5-lncRNA signature developed in our present study




Figure 2 The 5-long non-coding RNA signature-based risk score predicted the overall survival of patients with hepatocellular carcinoma in the training set ($n = 184$). A: Kaplan-Meier analysis of patients' overall survival in the high-risk ($n = 92$) and low-risk ($n = 92$) subgroups of the training set; B: The receiver operating characteristic (ROC) analysis of the risk score for prediction the overall survival of the training set. The area under the curve was calculated for ROC curves; C: The 5-lncRNA-based risk score distribution, patient survival status; D: Heatmap of the 5-lncRNA expression profiles in the high-risk and low-risk subgroups for the training set.

(hereafter referred to as 5LncSig) with the existing 3-gene signature by Binghua Li (hereafter referred to as 3GeneSig) and the 4-lncRNA signature by Zhonghao Wang (hereafter referred to as ZhongSig), we calculated the risk scores of each patient in the entire cohort based on formulae derived from each of these signatures.

The 3GeneSig and ZhongSig both successfully and significantly predicted prognosis in the entire TCGA LIHC cohort (Figure 6A and B). Furthermore, comparison of the Kaplan-Meier curves revealed that patients in the high-risk group predicted by 5LncSig showed a dramatically poorer prognosis than those in the low-

Table 3 Univariate and multivariate Cox regression analysis of overall survival

Variables	Univariate analysis			Multivariate analysis		
	HR	95%CI of HR	P value	HR	95%CI of HR	P value
Training set (<i>n</i> = 184)						
Risk score	2.718	2.093-3.530	< 0.0001	2.830	2.091-3.831	< 0.0001
Age	1.018	0.998-1.038	0.073			
Sex (Male/Female)	1.022	0.619-1.690	0.931			
Weight	1.006	0.993-1.108	0.397			
Child-Pugh grade	0.845	0.253-2.827	0.195			
Fibrosis ishak score	0.685	0.756-1.283	0.910			
Vascular invasion (yes/no)	0.834	0.480-1.449	0.519			
FAP	0.805	0.932-1.056	0.805			
Tumor grade (G1 + G2/G3 + G4)	1.133	0.680-1.887	0.632			
Pathologic stage				1.900	1.098-3.288	0.022
I / II	0.604	0.303-1.203	0.152			
I / III + IV	0.298	0.163-0.543	< 0.0001			
Test set (<i>n</i> = 186)						
Risk score	1.603	1.270-2.024	< 0.0001	1.568	1.196-2.055	0.001
Age	1.006	0.987-1.026	0.522			
Sex (Male/Female)	1.420	0.856-2.356	0.174			
Weight	1.000	0.984-1.105	0.960			
Child-Pugh grade	0.475	0.195-1.157	0.101			
Fibrosis ishak score	1.640	0.706-3.813	0.250			
Vascular invasion (yes/no)	1.069	0.613-1.866	0.614			
FAP	1.035	0.981-1.093	0.210			
Tumor grade (G1 + G2/G3 + G4)	1.092	0.655-1.818	0.736			
Pathologic stage				2.103	1.193-3.707	0.010
I / II	0.865	0.426-1.757	0.688			
I / III + IV	0.445	0.250-0.793	0.006			
Entire set (<i>n</i> = 370)						
Risk score	1.957	1.646-2.327	< 0.0001	2.011	1.638-2.469	< 0.0001
Age	1.013	0.999-1.027	0.068	1.016	1.000-1.032	0.048
Sex (Male/Female)	1.166	0.817-1.664	0.396			
Weight	0.998	0.988-1.007	0.615			
Child-Pugh grade (A/B + C)	0.620	0.306-1.256	0.184			
Fibrosis ishak score	1.232	0.742-2.045	0.365			
Vascular invasion (yes/no)	0.962	0.650-1.424	0.846			
Serum AFP level	1.023	0.980-1.068	0.306			
Tumor grade (G1 + G2/G3 + G4)	1.119	0.780-1.604	0.542			
Pathologic stage				2.017	1.359-2.993	0.027
I / II	0.648	0.398-1.056	0.082			
I / III + IV	0.351	0.236-0.524	< 0.0001			

risk groups predicted by the 3GeneSig and ZhongSig (Figure 6A and B), and patients in the low-risk group predicted by 5LncSig had a much better prognosis than those in the high-risk group predicted by the other two signatures (Figure 6A and B). To compare the sensitivity and specificity of the 5LncSig for prognosis prediction with the other two existing signatures, we performed time-dependent ROC analysis. The AUC of overall survival for the 3GeneSig and the ZhongSig was 0.701 and 0.721, respectively (Figure 6C), both lower than that of the 5LncSig (0.769). Thus, the prognostic power of 5LncSig, developed in the present study, was superior to that of the previously developed 3-gene and 4-lncRNA signatures.

Functional characteristics of the five prognostic lncRNAs

To explore the functional implications of these 5 lncRNAs,

we performed Pearson correlation analyses between the 5 lncRNAs and protein-coding genes based on their expression levels in the TCGA LIHC cohort. The protein-coding genes that correlated with at least 1 of the 5 lncRNAs (Pearson coefficient > 0.5, $P < 0.01$) were considered to be correlated genes. We chose the 200 correlated genes with the highest Pearson coefficients for further analysis. Functional enrichment analysis revealed that these genes were primarily enriched in 32 GO terms (Benjamin P value < 0.1, Figure 7A) and 23 KEGG pathways ($P < 0.001$, Figure 7B). Further analysis revealed that these enriched GO functional terms are mostly involved in metabolic processes, fibrinolysis and complement activation (Figure 7A).

DISCUSSION

HCC is a heterogeneous disease with differential




Figure 3 Validation of the prognostic value of the 5-long non-coding RNA signature for the hepatocellular carcinoma patients in the test group. A: Kaplan-Meier analysis indicated that patients in the high-risk ($n = 97$) subgroup exhibited significantly poorer survival than the low-risk subgroup ($n = 89$) in the test group; B: The receiver operating characteristic analysis of the risk score for predicting the overall survival of the test set; C: The 5-long non-coding RNA (lncRNA)-based risk score distribution, patient survival status; D: Heatmap of the 5-lncRNA expression profiles in the high-risk and low-risk subgroups of the test set.

prognoses and a high mortality. Until now, no biomarkers have been shown to effectively predict the survival of HCC patients, and thus, finding effective biomarkers for HCC is crucial.

Previous investigations of gene regulation and disease pathogenesis have mainly focused on protein-coding genes, which account for only a very small proportion

(2%) of transcribed genes in eukaryotic species^[13]. Recent developments in genome and transcriptome sequencing technologies have profoundly expanded our knowledge of non-coding RNAs, which are much more abundant than canonical protein-coding mRNAs^[30,31]. Multiple studies indicate that lncRNAs act not only as intermediaries between DNA and protein but also




Figure 4 The prognostic value of the 5-long non-coding RNA signature for the hepatocellular carcinoma patients in the entire set ($n = 370$). A: Kaplan-Meier analysis of patients' overall survival in the high-risk ($n = 189$) and low-risk subgroups ($n = 181$) for the entire set; B: The receiver operating characteristic analysis of the risk score for predicting the overall survival of the test set; C: The 5-long non-coding RNA (5-lncRNA)-based risk score distribution, patient survival status; D: Heatmap of the 5-lncRNA expression profiles in the high-risk and low-risk subgroups for the entire set.

as important regulators of diverse cellular functions. lncRNAs have been shown to regulate the expression and function of protein-coding genes at the chromatin, transcriptional and post-transcriptional levels^[31]. Many studies have revealed the contribution of lncRNAs in cancer development, indicating their potential as novel biomarkers for cancer diagnosis and prognosis^[32-35].

lncRNA signatures for prognostic prediction prognoses have been developed for many cancers including renal cancer, glioblastoma, and colorectal cancer, among others^[18-21]. Regarding HCC, the existing gene signatures for survival prediction have focused mostly on mRNAs and microRNAs. Several potential lncRNA biomarkers associated with the progression and




Figure 5 Stratification analyses of the prognostic value of the 5-long non-coding RNA signature-based risk score for all 370 hepatocellular carcinoma patients in the entire set. A: Kaplan-Meier analysis of the overall survival of hepatocellular carcinoma patients < 60 or ≥ 60 years old; B: Kaplan-Meier analysis of the overall survival of patients with different pathological stages.

prognosis of HCC have been identified, such as TSLNC8, HOXD-AS1 and CACS2^[36-38]. These lncRNAs are thought to impact HCC progression through their regulation of tumor cell proliferation, EMT, apoptosis and migration. Although many of these lncRNAs are closely associated

with the prognosis and survival of HCC patients, their prognostic value has been tested only in small-scale studies; they have not yet been validated in a large clinical cohort. Until now, relatively few comprehensive lncRNA signatures for the prediction of HCC survival




Figure 6 Comparison of the sensitivity and specificity for hepatocellular carcinoma survival prediction by the 5-long non-coding RNA signature and two existing signatures in the entire set. A: Kaplan-Meier analysis of patients' overall survival in the high-risk ($n = 185$) and low-risk ($n = 185$) subgroups based on the 3GeneSig in the entire set (left panel) and comparison of the survival difference based on the 5 long non-coding RNA (lncRNA) signature and the 3GeneSig by Kaplan-Meier analysis (right panel); B: Kaplan-Meier analysis of patients' overall survival in the high-risk ($n = 186$) and low-risk ($n = 184$) subgroups based on the ZhongSig in the entire set (left panel) and comparison of the survival difference based on the 5 lncRNA signature and the ZhongSig (right panel); C: The ROC analysis of overall survival for the 5-lncRNA signature, ZhongSig and 3GeneSig.

have been constructed^[26]. TCGA is an open-access database including samples from hundreds of patients with various malignancies. In the present study, we downloaded the RNA sequencing data of the TCGA LIHC

cohort and acquired lncRNA expression profiles for HCC patients in the dataset. Using univariate and stepwise multivariate Cox regression analyses, we developed a prognostic formula for HCC based on the expression




Figure 7 Functional enrichment analysis of the 5-long non-coding RNA based on their correlated protein-coding genes. A: Functional enrichment map of significantly enriched gene ontology terms; B: KEGG pathways significantly associated with the correlated protein-coding genes.

of 5 lncRNAs including AC015908.3, AC091057.3, TMCC1-AS1, DCST1-AS1 and FOXD2-AS1. In the

training set, HCC patients with high-risk scores based on the 5-lncRNA signature had a significantly reduced

survival time compared to those with low-risk scores. The prognostic value of the 5-lncRNA signature for HCC patients was further validated in the test group and the entire group, with robust and reproducible predictive indices. The results of these analyses suggest that the prognostic value of the 5-lncRNA-based risk model is robust and reliable for predicting survival in HCC patients.

In the present study, when adjusted using multivariate Cox regression analyses, age (only for the entire set), pathological stage and the 5-lncRNA signature were shown to independently predict the survival of HCC patients. The results of stratification analyses demonstrated that the prognostic value of the 5-lncRNA signature remained significant and robust in HCC subgroups stratified by age and pathological stage. In an attempt to further validate its prognostic value in other HCC cohorts, we downloaded data from several GEO datasets. Unfortunately, most of the 5 lncRNA probes could not be found. There are many existing prognostic signatures for HCC, and we therefore compared our 5-lncRNA signature with two recently developed signatures: a 3-gene signature and a 4-lncRNA signature. The results indicate that the predictive performance of the 5-lncRNA signature was superior to that of the other two signatures for HCC overall survival.

To the best of our knowledge, the functions of these 5 lncRNAs have not been reported. Functional enrichment analysis revealed that the protein-coding genes that were significantly correlated with these 5lncRNAs are enriched for metabolic processes, fibrinolysis and complement activation. KEGG pathway analysis revealed that these genes are enriched in pathways related to metabolism. These results suggest that the 5 lncRNAs may participate in the initiation and progression of HCC through these pathways. However, further studies are needed to investigate and validate the functions of these 5 lncRNAs.

In conclusion, our present study developed a 5-lncRNA signature for predicting the prognosis of HCC patients. The signature was reproducible and robust in a second independent large-scale HCC cohort, supporting its value and effectiveness. In addition, the prognostic value of the 5-lncRNA signature was independent of clinicopathological variables. Our study indicates that the 5-lncRNA signature could improve survival prediction and could be used as a prognostic biomarker for HCC patients.

ARTICLE HIGHLIGHTS

Research background

Hepatocellular carcinoma (HCC) is the sixth most commonly diagnosed cancer in the world. Although treatment for HCC, including surgical resection, has improved over the past decades, its overall survival rate remains devastatingly high due to its high rate of recurrence. Because HCC is a heterogeneous disease with substantially variable clinical outcomes, the search for effective biomarkers to predict recurrence and prognosis is crucial.

Research motivation

Recent studies have demonstrated the importance of long non-coding RNAs (lncRNAs) in physiological and pathological cellular processes. Increasing evidence suggests that lncRNA dysregulation is associated with various human diseases, particularly the initiation and progression of various human cancers. For patients with HCC, most of the existing prognostic signatures have focused on mRNAs or microRNAs, and only a few lncRNA signatures have been developed. In the present study, we aimed to construct a lncRNA signature for the prediction of HCC prognosis with high efficiency.

Research objectives

To construct a lncRNA signature for the prediction of HCC prognosis with high efficiency.

Research methods

Differentially expressed lncRNAs (DELs) between HCC specimens and peritumor liver specimens were acquired from the The Cancer Genome Atlas (TCGA) LIHC dataset using the edgeR package. Univariate Cox proportional hazards regression was performed to identify the DELs that were significantly associated with overall survival for the training set. The stepwise multivariate Cox regression model was applied. Those lncRNAs fitted in the multivariate Cox regression model and independently associated with overall survival were chosen to build a prognostic risk formula. The prognostic value of this formula was validated in the test group and the full cohort and further compared with two previously developed prognostic signatures for HCC.

Research results

We identified a five-lncRNA prognostic signature from the TCGA dataset and determined that its prognostic value was independent from clinicopathological factors. The signature was reproducible and robust in another independent large-scale HCC cohort, supporting its utility and effectiveness.

Research conclusions

This study constructed a 5-lncRNA signature that improves survival prediction, and can be used as a prognostic biomarker for HCC patients.

REFERENCES

- 1 Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015; **136**: E359-E386 [PMID: 25220842 DOI: 10.1002/ijc.29210]
- 2 Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin* 2015; **65**: 87-108 [PMID: 25651787 DOI: 10.3322/caac.21262]
- 3 Venook AP, Papandreou C, Furuse J, de Guevara LL. The incidence and epidemiology of hepatocellular carcinoma: a global and regional perspective. *Oncologist* 2010; **15** Suppl 4: 5-13 [PMID: 21115576 DOI: 10.1634/theoncologist.2010-S4-05]
- 4 Allemani C, Weir HK, Carreira H, Harewood R, Spika D, Wang XS, Bannon F, Ahn JV, Johnson CJ, Bonaventure A, Marcos-Gragera R, Stiller C, Azevedo e Silva G, Chen WQ, Ogunbiyi OJ, Rachet B, Soeberg MJ, You H, Matsuda T, Bielska-Lasota M, Storm H, Tucker TC, Coleman MP; CONCORD Working Group. Global surveillance of cancer survival 1995-2009: analysis of individual data for 25,676,887 patients from 279 population-based registries in 67 countries (CONCORD-2). *Lancet* 2015; **385**: 977-1010 [PMID: 25467588 DOI: 10.1016/S0140-6736(14)62038-9]
- 5 Mittal S, El-Serag HB. Epidemiology of hepatocellular carcinoma: consider the population. *J Clin Gastroenterol* 2013; **47** Suppl: S2-S6 [PMID: 23632345 DOI: 10.1097/MCG.0b013e3182872f29]
- 6 El-Serag HB, Rudolph KL. Hepatocellular carcinoma: epidemiology and molecular carcinogenesis. *Gastroenterology* 2007; **132**: 2557-2576 [PMID: 17570226 DOI: 10.1053/j.gastro.2007.04.061]
- 7 Joliat GR, Allemann P, Labgaa I, Demartines N, Halkic N.

- Treatment and outcomes of recurrent hepatocellular carcinomas. *Langenbecks Arch Surg* 2017; **402**: 737-744 [PMID: 28497194 DOI: 10.1007/s00423-017-1582-9]
- 8 **Bruix J**, Gores GJ, Mazzaferro V. Hepatocellular carcinoma: clinical frontiers and perspectives. *Gut* 2014; **63**: 844-855 [PMID: 24531850 DOI: 10.1136/gutjnl-2013-306627]
 - 9 **Zheng J**, Kuk D, Gönen M, Balachandran VP, Kingham TP, Allen PJ, D'Angelica MI, Jarnagin WR, DeMatteo RP. Actual 10-Year Survivors After Resection of Hepatocellular Carcinoma. *Ann Surg Oncol* 2017; **24**: 1358-1366 [PMID: 27921192 DOI: 10.1245/s10434-016-5713-2]
 - 10 **Kudo M**. Surveillance, diagnosis, treatment, and outcome of liver cancer in Japan. *Liver Cancer* 2015; **4**: 39-50 [PMID: 26020028 DOI: 10.1159/000367727]
 - 11 **Bruix J**, Sherman M; American Association for the Study of Liver Diseases. Management of hepatocellular carcinoma: an update. *Hepatology* 2011; **53**: 1020-1022 [PMID: 21374666 DOI: 10.1002/hep.24199]
 - 12 **Schmitt AM**, Chang HY. Long Noncoding RNAs: At the Intersection of Cancer and Chromatin Biology. *Cold Spring Harb Perspect Med* 2017; **7**: [PMID: 28193769 DOI: 10.1101/cshperspect.a026492]
 - 13 **Hangauer MJ**, Vaughn IW, McManus MT. Pervasive transcription of the human genome produces thousands of previously unidentified long intergenic noncoding RNAs. *PLoS Genet* 2013; **9**: e1003569 [PMID: 23818866 DOI: 10.1371/journal.pgen.1003569]
 - 14 **Mercer TR**, Dinger ME, Mattick JS. Long non-coding RNAs: insights into functions. *Nat Rev Genet* 2009; **10**: 155-159 [PMID: 19188922 DOI: 10.1038/nrg2521]
 - 15 **Moran VA**, Perera RJ, Khalil AM. Emerging functional and mechanistic paradigms of mammalian long non-coding RNAs. *Nucleic Acids Res* 2012; **40**: 6391-6400 [PMID: 22492512 DOI: 10.1093/nar/gks296]
 - 16 **Hauptman N**, Glavač D. Long non-coding RNA in cancer. *Int J Mol Sci* 2013; **14**: 4655-4669 [PMID: 23443164 DOI: 10.3390/ijms14034655]
 - 17 **Gibb EA**, Brown CJ, Lam WL. The functional role of long non-coding RNA in human carcinomas. *Mol Cancer* 2011; **10**: 38 [PMID: 21489289 DOI: 10.1186/1476-4598-10-38]
 - 18 **Shi D**, Qu Q, Chang Q, Wang Y, Gui Y, Dong D. A five-long non-coding RNA signature to improve prognosis prediction of clear cell renal cell carcinoma. *Oncotarget* 2017; **8**: 58699-58708 [PMID: 28938589 DOI: 10.18632/oncotarget.17506]
 - 19 **Zhou M**, Zhang Z, Zhao H, Bao S, Cheng L, Sun J. An Immune-Related Six-lncRNA Signature to Improve Prognosis Prediction of Glioblastoma Multiforme. *Mol Neurobiol* 2018; **55**: 3684-3697 [PMID: 28527107 DOI: 10.1007/s12035-017-0572-9]
 - 20 **Zeng JH**, Liang L, He RQ, Tang RX, Cai XY, Chen JQ, Luo DZ, Chen G. Comprehensive investigation of a novel differentially expressed lncRNA expression profile signature to assess the survival of patients with colorectal adenocarcinoma. *Oncotarget* 2017; **8**: 16811-16828 [PMID: 28187432 DOI: 10.18632/oncotarget.15161]
 - 21 **Sun J**, Cheng L, Shi H, Zhang Z, Zhao H, Wang Z, Zhou M. A potential panel of six-long non-coding RNA signature to improve survival prediction of diffuse large-B-cell lymphoma. *Sci Rep* 2016; **6**: 27842 [PMID: 27292966 DOI: 10.1038/srep27842]
 - 22 **Lin Z**, Xu SH, Wang HQ, Cai YJ, Ying L, Song M, Wang YQ, Du SJ, Shi KQ, Zhou MT. Prognostic value of DNA repair based stratification of hepatocellular carcinoma. *Sci Rep* 2016; **6**: 25999 [PMID: 27174663 DOI: 10.1038/srep25999]
 - 23 **Lu M**, Kong X, Wang H, Huang G, Ye C, He Z. A novel microRNAs expression signature for hepatocellular carcinoma diagnosis and prognosis. *Oncotarget* 2017; **8**: 8775-8784 [PMID: 28060739 DOI: 10.18632/oncotarget.14452]
 - 24 **Borel F**, Konstantinova P, Jansen PL. Diagnostic and therapeutic potential of miRNA signatures in patients with hepatocellular carcinoma. *J Hepatol* 2012; **56**: 1371-1383 [PMID: 22314424 DOI: 10.1016/j.jhep.2011.11.026]
 - 25 **Li B**, Feng W, Luo O, Xu T, Cao Y, Wu H, Yu D, Ding Y. Development and Validation of a Three-gene Prognostic Signature for Patients with Hepatocellular Carcinoma. *Sci Rep* 2017; **7**: 5517 [PMID: 28717245 DOI: 10.1038/s41598-017-04811-5]
 - 26 **Wang Z**, Wu Q, Feng S, Zhao Y, Tao C. Identification of four prognostic lncRNAs for survival prediction of patients with hepatocellular carcinoma. *PeerJ* 2017; **5**: e3575 [PMID: 28729955 DOI: 10.7717/peerj.3575]
 - 27 **Huang da W**, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 2009; **37**: 1-13 [PMID: 19033363 DOI: 10.1093/nar/gkn923]
 - 28 **Huang da W**, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 2009; **4**: 44-57 [PMID: 19131956 DOI: 10.1038/nprot.2008.211]
 - 29 **Merico D**, Isserlin R, Stueker O, Emili A, Bader GD. Enrichment map: a network-based method for gene-set enrichment visualization and interpretation. *PLoS One* 2010; **5**: e13984 [PMID: 21085593 DOI: 10.1371/journal.pone.0013984]
 - 30 **Chen JA**, Conn S. Canonical mRNA is the exception, rather than the rule. *Genome Biol* 2017; **18**: 133 [PMID: 28687088 DOI: 10.1186/s13059-017-1268-1]
 - 31 **Shi X**, Sun M, Liu H, Yao Y, Song Y. Long non-coding RNAs: a new frontier in the study of human diseases. *Cancer Lett* 2013; **339**: 159-166 [PMID: 23791884 DOI: 10.1016/j.canlet.2013.06.013]
 - 32 **Bi M**, Yu H, Huang B, Tang C. Long non-coding RNA PCAT-1 over-expression promotes proliferation and metastasis in gastric cancer cells through regulating CDKN1A. *Gene* 2017; **626**: 337-343 [PMID: 28571676 DOI: 10.1016/j.gene.2017.05.049]
 - 33 **Hua F**, Liu S, Zhu L, Ma N, Jiang S, Yang J. Highly expressed long non-coding RNA NNT-AS1 promotes cell proliferation and invasion through Wnt/ β -catenin signaling pathway in cervical cancer. *Biomed Pharmacother* 2017; **92**: 1128-1134 [PMID: 28628975 DOI: 10.1016/j.biopha.2017.03.057]
 - 34 **Xu S**, Kong D, Chen Q, Ping Y, Pang D. Oncogenic long noncoding RNA landscape in breast cancer. *Mol Cancer* 2017; **16**: 129 [PMID: 28738804 DOI: 10.1186/s12943-017-0696-6]
 - 35 **Spurlock CF 3rd**, Shaginurova G, Tossberg JT, Hester JD, Chapman N, Guo Y, Crooke PS 3rd, Aune TM. Profiles of Long Noncoding RNAs in Human Naïve and Memory T Cells. *J Immunol* 2017; **199**: 547-558 [PMID: 28600289 DOI: 10.4049/jimmunol.1700232]
 - 36 **Zhang J**, Li Z, Liu L, Wang Q, Li S, Chen D, Hu Z, Yu T, Ding J, Li J, Yao M, Huang S, Zhao Y, He X. Long noncoding RNA TSLNC8 is a tumor suppressor that inactivates the interleukin-6/STAT3 signaling pathway. *Hepatology* 2018; **67**: 171-187 [PMID: 28746790 DOI: 10.1002/hep.29405]
 - 37 **Lu S**, Zhou J, Sun Y, Li N, Miao M, Jiao B, Chen H. The noncoding RNA HOXD-AS1 is a critical regulator of the metastasis and apoptosis phenotype in human hepatocellular carcinoma. *Mol Cancer* 2017; **16**: 125 [PMID: 28724429 DOI: 10.1186/s12943-017-0676-x]
 - 38 **Wang Y**, Liu Z, Yao B, Li Q, Wang L, Wang C, Dou C, Xu M, Liu Q, Tu K. Long non-coding RNA CASC2 suppresses epithelial-mesenchymal transition of hepatocellular carcinoma cells through CASC2/miR-367/FBXW7 axis. *Mol Cancer* 2017; **16**: 123 [PMID: 28716020 DOI: 10.1186/s12943-017-0702-z]

P- Reviewer: Namisaki T, Tsui SK S- Editor: Wang JL

L- Editor: Filipodia E- Editor: Huang Y





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327

