



1

**Favorable Clinical Outcome of Nonalcoholic Liver Cirrhosis Patients with
Coronary Artery Disease: A Population-based Study**

Tsai MC *et al.* Better survival in non-alcoholic LC with CAD

Ming-Chang Tsai, Tzu-Wei Yang, Chi-Chih Wang, Yao-Tung Wang,
Wen-Wei Sung, Ming-Hseng Tseng, Chun-Che Lin

Match Overview



- 1** **Crossref** 17 words
Ta-Liang Chen, Chao-Shun Lin, Chun-Chuan Shih, Yu-Feng Huang, Chun-Chieh Yeh, Chih-Hsing Wu, Yih-Giun Cherng, **<1%**
- 2** **Crossref** 13 words
"13th European Congress of Clinical Microbiology and Infectious Diseases", Clinical Microbiology and Infection, 2003 **<1%**
- 3** **Internet** 12 words
crawled on 25-Aug-2014 www.biomedcentral.com **<1%**



Favorable Clinical Outcome of Non-Alcoholic Liver Cirrhosis Patients wit



All

News

Images

Shopping

Videos

More

Settings

Tools

About 2,690,000 results (0.62 seconds)

Scholarly articles for **Favorable Clinical Outcome of Non-Alcoholic Liver Cirrhosis Patients with Coronary Artery Disease: A Population-Based Study**

... fatty liver disease: a **population-based** cohort **study** - Adams - Cited by 2320

... **disease and** increased risk of **cardiovascular disease** - Targher - Cited by 597

... **and** management of **non-alcoholic** fatty liver disease: ... - Chalasani - Cited by 2383

Non-Alcoholic Fatty Liver Disease and Cardiovascular Risk - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2732016/> ▼

by VL Misra - 2009 - Cited by 115 - Related articles

The latter presentation can progress to **cirrhosis** in 15–20% of subjects over the course of ...

Cardiovascular disease risk factors and **nonalcoholic** fatty liver disease ... Various surrogates used in **clinical** studies include Framingham Risk Score ... [22] A **population-based study** of 980 NAFLD **patients** and 6,594 controls ...

Fatty liver is associated with an increased risk of diabetes and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5124973/> ▼

by A Lonardo - 2016 - Cited by 30 - Related articles

Nov 28, 2016 - In this multidisciplinary **clinical** review, we aim to discuss the ... For example, **nonalcoholic** fatty liver disease (NAFLD) is defined by fatty changes which artery disease (**CAD**) as a pre-specified **outcome**; it reported that **patients** A **population-based** cohort **study** showed that HCV-infected **patients** have a ...

全部

新闻

图片

购物

地图

更多

设置

工具

找到约 257,000 条结果 (用时 0.66 秒)

Fatty liver is associated with an increased risk of diabetes and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5124973/> ▼ 翻译此页

作者: A Lonardo - 2016 - 被引用次数: 30 - 相关文章

2016年11月28日 - This review may also open new avenues in the clinical and research arenas ... For example, nonalcoholic fatty liver disease (NAFLD) is defined by fatty changes population-based cohort studies, have reported a strong and ... artery disease (CAD) as a pre-specified outcome; it reported that patients with ...

Non-alcoholic fatty liver disease and non-alcoholic steatohepatitis

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2766096/> ▼ 翻译此页

作者: R Vuppalanchi - 2009 - 被引用次数: 648 - 相关文章

This review is tailored towards clinicians caring for patients with NAFLD and it ... This notion is supported by a population-based study in which Ruhl and Undoubtedly, NASH is histologically progressive and can lead to cirrhosis and All these data provide unequivocal evidence that coronary artery disease is a serious ...

Nonalcoholic fatty liver disease and chronic vascular complications of ...

<https://www.nature.com/nature-reviews-endocrinology/review/article> - 翻译此页

作者: G Targher - 2018 - 被引用次数: 9 - 相关文章

2017年12月29日 - Nonalcoholic fatty liver disease (NAFLD) and diabetes mellitus are common ... act synergistically to increase the risk of hepatic and extra-hepatic clinical outcomes. ... risk of developing cardiovascular disease and other cardiac and ... of the population studied (for example, studies in patients with different ...

Prevalence and Prediction of Coronary Artery Disease in Patients With ...

circ.ahajournals.org/content/130/16/1353 ▼ 翻译此页

作者: J An - 2014 - 相关文章

2014年10月14日 - This study aimed to investigate the prevalence of silent CAD in comparison ... consecutive patients with liver cirrhosis without any history of chest pain or CAD, ... the result with matched individuals in the nonhepatic population. ... LC was diagnosed based on clinical features of portal hypertension together ...