

7 Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 39338

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Long-term outcomes of **3** endoscopic resection for small (≤ 4.0 cm) gastric gastrointestinal stromal tumors originating from the muscularis propria layer

Yu Zhang, Xin-Li Mao, Xian-Bin Zhou, Hai Yang, Lin-Hong Zhu; Guang Chen, Li-Ping Ye

Abstract

AIM

Match Overview

1	Crossref 163 words Ye, Li-Ping, Yu Zhang, Ding-Hai Luo, Xin-Li Mao, Hai-Hong Zheng, Xian-Bin Zhou, and Lin-Hong Zhu. "Safety of	4%
2	Crossref 26 words Ye, Li-Ping, Hai-Hong Zheng, Xin-Li Mao, Yu Zhang, Xian-Bin Zhou, and Lin-Hong Zhu. "Complete circular endos	1%
3	Crossref 26 words "Gastrointestinal Stromal Tumors", Springer Nature, 2017	1%
4	Crossref 24 words Qiang Zhang, Liang-Qing Gao, Ze-Long Han, Xiao-Feng Li, Li-Hui Wang, Si-De Liu. "Effectiveness and safety of e	1%
5	Crossref 15 words Joo, Moon Kyung, Jong-Jae Park, Ho Kim, Jin Sung Koh, Beom Jae Lee, Hoon Jai Chun, Sang Woo Lee, You-Jin	<1%
6	Crossref 15 words Shen, Chaoyong, Haining Chen, Yuan Yin, Jiaju Chen, Luyin Han, Bo Zhang, Zhixin Chen, and Jiaping Chen. "En	<1%
7	Internet 15 words crawled on 05-Oct-2017 real.mtak.hu	<1%

[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 5,030 条结果 (用时 0.60 秒)

Endoscopic Versus Open Resection for Small Gastric Gastrointestinal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4602836/> ▼ [翻译此页](#)

作者: C Shen - 2015 - 被引用次数: 14 - [相关文章](#)

2015年1月9日 - The medical records of 54 consecutive **gastric** GISTs patients with tumor size of ≤ 2 **cm**, who were surgically treated with **endoscopic resection** ... **Gastrointestinal stromal tumors** (GISTs), which **originate** from the interstitial cells of Cajal or its precursor, may be asymptomatic and nonmalignant when ...

Endoscopic treatment for gastrointestinal stromal tumor: Advantages ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4360437/> ▼ [翻译此页](#)

作者: HH Kim - 2015 - 被引用次数: 31 - [相关文章](#)

2015年3月16日 - Short- and **long-term outcomes** of laparoscopic wedge **resection** have been shown to be equivalent to the open surgical approach. Guidelines suggest that laparoscopic wedge **resection** can be used for **tumors ≤ 5 cm**[22]. Laparoscopic approaches to **GIST** management continue to expand and should ...

Comparison of Endoscopic and Open Resection for Small Gastric ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4700288/> ▼ [翻译此页](#)

作者: F Feng - 2015 - 被引用次数: 11 - [相关文章](#)

2015年12月12日 - Abstract. The National Comprehensive Cancer Network recommends conservative follow-up for **gastric gastrointestinal stromal tumors** (GISTs) less than 2 **cm**. We have ... The aim of the present study is to compare the safety and **outcomes** of **endoscopic** and open **resection** of **small**



Long-term outcomes of endoscopic resection for small (≤ 4.0 cm) gastric



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 21,000 条结果 (用时 0.50 秒)

Endoscopic resection of gastric gastrointestinal stromal tumors ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/28916847> - 翻译此页

作者: I Andalib - 2017 - 被引用次数: 2 - 相关文章

2017年9月15日 - Endoscopic resection of gastric gastrointestinal stromal tumors originating from the muscularis propria layer in North America: ... The main outcome measures were complete resection rate, operative time, postoperative ...

Endoscopic resection of gastric gastrointestinal stromal tumors

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5763013/> - 翻译此页

作者: Y Tan - 2017 - 被引用次数: 1 - 相关文章

2017年12月19日 - Gastrointestinal stromal tumors (GISTs) are the most common mesenchymal ... is recommended for small (<2 cm) asymptomatic gastric GISTs. ... And EBL is feasible only for GISTs originating from superficial muscularis propria layer. Long-term and surgical outcomes of laparoscopic surgery for gastric ...

Comparison of Endoscopic and Open Resection for Small Gastric ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4700288/> - 翻译此页

作者: F Feng - 2015 - 被引用次数: 13 - 相关文章

2015年12月12日 - Compared with the surgical resection group, the operation time was shorter ($P = .000$), ... Endoscopic resection of small gastric GISTs is safe and feasible ... Gastrointestinal stromal tumors (GISTs) are the most common ≤ 60 , 18, 28 .300 subepithelial tumors originating in the muscularis propria layer.

Long-term Outcomes of Submucosal Tunneling Endoscopic Resection ...

https://journals.lww.com/annalsofsurgery/.../Long_term_Outcomes_of_Sub... - 翻译此页

作者: T Chen - 2017 - 被引用次数: 24 - 相关文章

Objective: The aim of this study is to evaluate the long-term outcomes of a large series of patient ... submucosal tumors originating from the muscularis propria layer ... a transverse diameter <2.5 cm



Long-term outcomes of endoscopic resection for small (≤ 4.0 cm) gastric



全部

图片

视频

新闻

购物

更多

设置

工具

找到约 21,000 条结果 (用时 0.51 秒)

Endoscopic resection of gastric gastrointestinal stromal tumors ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/28916847> - 翻译此页

作者: I Andalib - 2017 - 被引用次数: 2 - 相关文章

2017年9月15日 - Endoscopic resection of gastric gastrointestinal stromal tumors originating from the muscularis propria layer in North America: ... The main outcome measures were complete resection rate, operative time, postoperative ...

Endoscopic resection of gastric gastrointestinal stromal tumors

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5763013/> - 翻译此页

作者: Y Tan - 2017 - 被引用次数: 1 - 相关文章

2017年12月19日 - Gastrointestinal stromal tumors (GISTs) are the most common mesenchymal ... is recommended for small (<2 cm) asymptomatic gastric GISTs. ... And EBL is feasible only for GISTs originating from superficial muscularis propria layer. Long-term and surgical outcomes of laparoscopic surgery for gastric ...

Comparison of Endoscopic and Open Resection for Small Gastric ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4700288/> - 翻译此页

作者: F Feng - 2015 - 被引用次数: 13 - 相关文章

2015年12月12日 - Compared with the surgical resection group, the operation time was shorter ($P = .000$), ... Endoscopic resection of small gastric GISTs is safe and feasible ... Gastrointestinal stromal tumors (GISTs) are the most common $\leq 60, 18, 28, 300$ subepithelial tumors originating in the muscularis propria layer.

Long-term Outcomes of Submucosal Tunneling Endoscopic Resection ...

https://journals.lww.com/annalsofsurgery/.../Long_term_Outcomes_of_Sub... - 翻译此页

作者: T Chen - 2017 - 被引用次数: 24 - 相关文章

Objective: The aim of this study is to evaluate the long-term outcomes of a large series of patient ... submucosal tumors originating from the muscularis propria layer. a transverse diameter ≤ 3.5 cm could facilitate a high en bloc resection rate. relatively small (2–5 cm) gastrointestinal stromal tumor of the stomach based ...