

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

获得 10 条结果 (用时 0.48 秒)

### Prognostic factors in neuroendocrine carcinoma ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5225444/> [▼ 翻译此页](#)

作者: P Freis - 2017 - 被引用次数: 3 - 相关文章

2017年1月11日 - Gastroenteropancreatic neuroendocrine carcinomas (GEP-NEC) are a very aggressive type of cancer, for which **prognostic factors** are lacking. We analysed .... The other factors investigated, such as inflammation markers (low albumin, high CRP, and high NLR ratio) were not **prognostic factors**. Patients ...

### C-reactive protein level is a prognostic indicator for survival and ...

<https://www.nature.com › british journal of cancer › clinical study> [- 翻译此页](#)

作者: KT Troppan - 2014 - 被引用次数: 22 - 相关文章

2014年5月29日 - In the present study, we demonstrated high CRP levels at diagnosis of DLBCL as an independent poor **prognostic factor** for clinical outcome. Adding CRP to ... GEP has revealed three subtypes of aggressive lymphoma, discerning in developmental stage and usage of different oncogenic pathways. Patients ...

### C-Reactive Protein as a Prognostic Factor for Human Osteosarcoma ...

<journals.plos.org/plosone/article?id=10.1371/journal.pone.0094632> [▼ 翻译此页](#)

作者: JH Yi - 2014 - 被引用次数: 26 - 相关文章

2014年5月6日 - Citation: Yi J-H, Wang D, Li Z-Y, Hu J, Niu X-F, Liu X-L (2014) **C-Reactive Protein** as a **Prognostic Factor** for Human Osteosarcoma: A Meta-Analysis and ..... of nonmetastatic osteosarcoma of the extremity with preoperative and postoperative chemotherapy: a report from the Children's Cancer Group.

缺少字词: gep

### Prognostic factors in neuroendocrine carcinoma ... - ResearchGate

[https://www.researchgate.net/.../312442720\\_Prognostic\\_factors\\_in\\_neuroen...](https://www.researchgate.net/.../312442720_Prognostic_factors_in_neuroen...) [- 翻译此页](#)

2017年12月21日 - Full-text (PDF) | Gastroenteropancreatic neuroendocrine carcinomas (GEP-NEC) are a very aggressive type of cancer, for which **prognostic factors** are lacking. We analysed clinical ... Very recently, one study performed on 149 pancreatic neuroendocrine neoplasms demonstrated that CRP level is a new ...

**Name of Journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 39357

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Study*

**Is C-reactive protein a prognostic factor for the whole GEP-NET group?**

**Ömer Komaç, Göksel Bengi, Özgül Sağol, Mesut Akarsu**

**Abstract**

**BACKGROUND:** Gastroenteropancreatic neuroendocrine tumors (GEP-NETs) constitute a heterogeneous group of tumors with variable clinical presentations, different growth rates and unpredictable prognoses. In our study, we aimed to identify the independent prognostic markers for the patients in the whole GEP-NET

### Match Overview

There are no matching sources for this report.

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 221,000 条结果 (用时 0.46 秒)

### Is C-reactive protein a prognostic factor of colorectal cancer? - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/18172726> - 翻译此页

作者: YC Shiu - 2008 - 被引用次数: 60 - 相关文章

PURPOSE: **C-reactive protein**, a commonly used inflammation marker, has been reported to be a **prognostic factor** of colorectal cancer. This prospective ... The quiescent inflammation-response **group** (< or =0.1 mg/dl) had excellent outcomes.

缺少字词: whole gep net

### Prognostic factors in neuroendocrine carcinoma: biological markers ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5225444/> - 翻译此页

作者: P Freis - 2017 - 被引用次数: 3 - 相关文章

2017年1月11日 - Gastroenteropancreatic neuroendocrine carcinomas (GEP-NEC) are a very ... and 8 months in the **group** with low LDH and AST levels ( $p < 0.001$ ). ... Nine patients were excluded because of well-differentiated NET (n ... The **CRP**/albumin ratio is also a **prognostic factor** for survival in small cell lung cancer.

### C-reactive protein level is a prognostic indicator for survival and ...

<https://www.nature.com/british journal of cancer/clinical study> - 翻译此页

作者: KT Troppan - 2014 - 被引用次数: 24 - 相关文章

2014年5月29日 - The **prognostic** influence of **CRP** and other **factors**, including age, tumour ... GEP has revealed three subtypes of aggressive lymphoma, ... whereas a third, smaller **group** lies in between (Rosenwald et al, 2002). ... Using this cutoff value of  $15 \text{ mg l}^{-1}$  for the **whole** cohort, the association between **CRP** levels ...

### C-Reactive Protein as a Prognostic Factor for Human Osteosarcoma ...

[journals.plos.org/plosone/article?id=10.1371/journal.pone.0094632](https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0094632) - 翻译此页

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 35,600 条结果 (用时 0.43 秒)

### Prognostic factors in neuroendocrine carcinoma: biological markers ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5225444/> ▼ [翻译此页](#)作者: P Freis - 2017 - 被引用次数: 3 - [相关文章](#)

2017年1月11日 - **Gastroenteropancreatic neuroendocrine** carcinomas (GEP-NEC) are a very ... and 8 months in the **group** with low LDH and AST levels ( $p < 0.001$ ). .... The median values for **complete** blood cell count and coagulation .... The **CRP**/albumin ratio is also a **prognostic factor** for survival in small cell lung cancer.

### Is C-reactive protein a prognostic factor of colorectal cancer? - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/18172726> - [翻译此页](#)作者: YC Shiu - 2008 - 被引用次数: 63 - [相关文章](#)

2018年1月3日 - **PURPOSE: C-reactive protein**, a commonly used inflammation marker, has been reported to be a **prognostic factor** of colorectal cancer. ... The quiescent inflammation-response **group** ( $< \text{or} = 0.1 \text{ mg/dl}$ ) had excellent outcomes.

缺少字词: whole gastroenteropancreatic neuroendocrine

### Gastroenteropancreatic neuroendocrine tumors: diagnosis and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3959515/> ▼ [翻译此页](#)作者: M Díez - 2013 - 被引用次数: 25 - [相关文章](#)

**Gastroenteropancreatic neuroendocrine tumors** (GEP-NETs) are relatively rare ... Chromogranin A determination is also useful for staging, **prognosis** and .... A small **group** of patients with bilobar liver metastases without extrahepatic .... **growth factor** receptors (PDGFRs), c-kit and epidermal growth factor receptor (EGFR).

### Prognostic factors in neuroendocrine carcinoma - ResearchGate

[https://www.researchgate.net/.../312442720\\_Prognostic\\_factors\\_in\\_neuroen...](https://www.researchgate.net/.../312442720_Prognostic_factors_in_neuroen...) - [翻译此页](#)2018年8月1日 - Full-Text Paper (PDF): **Prognostic factors in neuroendocrine carcinoma: ...**

**Gastroenteropancreatic neuroendocrine** carcinomas (GEP-NEC) are a very ... 13 months and 8 months in the **group** with low LDH and AST levels ( $p < 0.001$ ). ..... albumin, high **CRP**, and high NLR ratio) were not **prognostic factors**.

### Epigenetics of gastroenteropancreatic neuroendocrine tumors: A ...

<https://www.wjnet.com/1948-5204/full/v9/i9/341.htm> ▼ [翻译此页](#)作者: BM Finnerty - 2017 - 被引用次数: 3 - [相关文章](#)

2017年9月15日 - **Gastroenteropancreatic neuroendocrine tumors** (GEP-NETs) are a heterogeneous ... data identifying potential **prognostic** biomarkers, and some of these have ... lack of a **complete** response