

World Journal of *Nephrology*

World J Nephrol 2018 October 10; 7(6): 117-128



**MINIREVIEWS**

- 117 Oral alkali therapy and the management of metabolic acidosis of chronic kidney disease: A narrative literature review

Ahmed AR, Lappin D

ORIGINAL ARTICLE**Observational Study**

- 123 Bicarbonate levels in hemodialysis patients switching from lanthanum carbonate to sucroferric oxyhydroxide

Stavroulopoulos A, Aresti V, Papadopoulos C, Nennes P, Metaxaki P, Galinas A

ABOUT COVER

Editorial Board Member of *World Journal of Nephrology*, Sebastian Doff, MD, PhD, Doctor, Department of Nephrology, University Hospital Essen, Essen 45122, Netherlands

AIM AND SCOPE

World Journal of Nephrology (*World J Nephrol*, *WJN*, online ISSN 2220-6124, DOI: 10.5527) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJN covers topics concerning kidney development, renal regeneration, kidney tumors, therapy of renal disease, hemodialysis, peritoneal dialysis, kidney transplantation, diagnostic imaging, evidence-based medicine, epidemiology and nursing. Priority publication will be given to articles concerning diagnosis and treatment of nephrology diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJN*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Nephrology (*WJN*) is now abstracted and indexed in PubMed, PubMed Central, China National Knowledge Infrastructure (CNKI), and Superstar Journals Database.

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Yan Huang*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fan-Fan Ji*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Nephrology

ISSN
ISSN 2220-6124 (online)

LAUNCH DATE
February 6, 2012

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjnet.com/2220-6124/editorialboard.htm>

EDITORIAL OFFICE
Jin-Lei Wang, Director
World Journal of Nephrology
Baishideng Publishing Group Inc
7901 Stoneridge Drive,
Suite 501, Pleasanton, CA 94588, USA

Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive,
Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjnet.com>

PUBLICATION DATE
October 10, 2018

COPYRIGHT

© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Observational Study

Bicarbonate levels in hemodialysis patients switching from lanthanum carbonate to sucroferric oxyhydroxide

Aristeidis Stavroulopoulos, Vasiliki Aresti, Christoforos Papadopoulos, Panagiotis Nennes, Polixeni Metaxaki, Anastasios Galinas

Aristeidis Stavroulopoulos, Vasiliki Aresti, Christoforos Papadopoulos, Panagiotis Nennes, Polixeni Metaxaki, Anastasios Galinas, Department of Nephrology, IASIO Hospital, General Clinic of Kallithea, Athens 17676, Greece

ORCID number: Aristeidis Stavroulopoulos (0000-0002-1368-5892); Vasiliki Aresti (0000-0002-9741-9005); Christoforos Papadopoulos (0000-0002-4370-6084); Panagiotis Nennes (0000-0003-1760-3302); Polixeni Metaxaki (0000-0003-0107-3431); Anastasios Galinas (0000-0002-0566-8558).

Author contributions: All of the authors contributed to the study design; Stavroulopoulos A, Aresti V and Papadopoulos C collected the data; Stavroulopoulos A analyzed the data and drafted the manuscript; all authors contributed to the conception of the paper, interpretation of data, and subsequent revisions of the manuscript.

Institutional review board statement: The protocol for the research project has been approved by the IASIO Hospital's Ethics Committee and conforms to the provisions of the Declaration of Helsinki.

Informed consent statement: Informed written consent was obtained from all patients before they entered the study.

Conflict-of-interest statement: None to declare.

STROBE statement: The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Aristeidis Stavroulopoulos, BM BCh, MD, Chief Doctor, Doctor, Department of Nephrology, IASIO Hospital, General Clinic of Kallithea, Sivitanidou 36, Athens 17676, Greece. stavoulopoulos@yahoo.co.uk

Telephone: +30-69-47824197

Fax: +30-21-09583722

Received: June 4, 2018

Peer-review started: June 4, 2018

First decision: July 11, 2018

Revised: August 2, 2018

Accepted: August 30, 2018

Article in press: August 30, 2018

Published online: October 10, 2018

Abstract

AIM

To examine possible alterations in acid-base parameters in patients switching from lanthanum carbonate (LanC) to sucroferric oxyhydroxide (SFOH).

METHODS

Fifteen stable hemodialysis patients were switched from LanC to SFOH. Only nine continued on SFOH, three returned to LanC and the other three switched to sevelamer carbonate. The later six patients served as a control group to the SFOH group of nine patients. Blood was sampled on the 3-d and the last 2-d interval of the week prior to switching and six weeks after. Bicarbonate levels (HCO_3^-), pH, pO_2 , pCO_2 were measured, and the mean of the two measurements (3-d and 2-d interval) was calculated.

RESULTS

Comparing pre-switching to post-switching measurements

in the SFOH group, no statistically significant differences were found in any of the parameters studied. The mean pre-switching HCO_3^- was 22.41 ± 1.66 mmol/L and the mean post-switching was 22.62 ± 2.25 mmol/L ($P = 0.889$). Respectively, the mean pH= 7.38 ± 0.03 vs 7.39 ± 0.03 ($P = 0.635$), mean pCO_2 = 38.41 ± 3.29 vs 38.37 ± 3.62 mmHg ($P = 0.767$), and Phosphate = 1.57 ± 0.27 vs 1.36 ± 0.38 mmol/L ($P = 0.214$). There were not any significant differences when we performed the same analyses in the control group or between the SFOH group and control group. No correlations were found, either between pre-switching LanC daily dose or between post-switching daily dose of the new binder and the measured parameters.

CONCLUSION

In our small study, switching from LanC to SFOH did not have any significant effect on blood bicarbonate levels and gas analysis, indicating that there is no need to change hemodialysis prescription regarding these parameters.

Key words: Gas analysis; Hemodialysis; Lanthanum carbonate; Acidosis; Bicarbonate; Phosphate binder; SuCroferric oxyhydroxide

© The Author(s) 2018. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Phosphate binders used for the control of hyperphosphatemia contribute to acid-base balance through their effect on serum phosphate and through the effect of the binders' constituents that have alkaline or acidic properties. This is the first study showing that switching from Lanthanum Carbonate to the novel phosphate binder sucroferric oxyhydroxide did not have any significant effect on blood bicarbonate levels and gas analysis. Thus, there is no need to change hemodialysis prescription regarding these parameters.

Stavroulopoulos A, Aresti V, Papadopoulos C, Nennes P, Metaxaki P, Galinas A. Bicarbonate levels in hemodialysis patients switching from lanthanum carbonate to sucroferric oxyhydroxide. *World J Nephrol* 2018; 7(6): 123-128 Available from: URL: <http://www.wjgnet.com/2220-6124/full/v7/i6/123.htm> DOI: <http://dx.doi.org/10.5527/wjn.v7.i6.123>

INTRODUCTION

Metabolic acidosis is a characteristic complication of end stage kidney disease (ESKD) and is associated with major systemic effects and increased mortality^[1,2]. Hyperphosphatemia, another common complication of ESKD, partially contributes to metabolic acidosis and affects the acid-base status of the patients^[1]. Phosphate binders used for control of hyperphosphatemia contribute to acid-base balance through their effect on serum phosphate and through the effect of the binders'

constituents that have alkaline or acidic properties^[3]. The most characteristic is the case of sevelamer hydrochloride (SevH), which is associated with dose-dependent aggravation of metabolic acidosis in ESKD patients due to its hydrochloride content^[4,5]. This acidosis is ameliorated after switching to sevelamer carbonate (SevC) or other phosphate binders with alkaline content, such as calcium carbonate (CaC) or lanthanum carbonate (LanC)^[6-9]. However, despite the above mentioned studies, a recent meta-analysis suggests that the effect of phosphate binders in clinically important outcomes, such as metabolic acidosis, remains understudied^[10].

SuCroferric oxyhydroxide (SFOH) is a polynuclear iron(III)-oxyhydroxide-based phosphate binder recently approved for the treatment of hyperphosphatemia in patients with ESKD. It is a safe and potent phosphate binder with increasing use among hemodialysis patients, both as an initial treatment choice or as an alternative when other binders fail^[11,12]. However, little is known regarding the influence of this novel binder on the acid-base status of hemodialysis patients. Aim of our study was to examine possible alterations in acid-base parameters in hemodialysis patients switching from LanC to SFOH.

MATERIALS AND METHODS

Study population

We prospectively evaluated 15 clinically stable, Caucasian, anuric patients from the same unit who switched from LanC to SFOH (Table 1). The patients were taking LanC in the form of 750 mg chewable pills (Fosrenol, Shire Pharmaceuticals Cont. Ltd, United Kingdom) for at least 6 mo, but they had to change phosphate binder due to logistic reasons (disposal and access to the drug limitations). The new binder SFOH (Velphoro, Vifor Fresenius Medical Care Renal Pharma Ltd) was chosen as it is chewable with low pill burden, like LanC. SFOH pills of 500 mg were gradually introduced and titrated within two weeks to a maintenance dose. However, only nine continued on the new binder. The main reasons for discontinuation were gastrointestinal symptoms and taste of the product; thus, three patients returned to LanC and the rest switched to SevC. Therefore, we decided to use these six patients as a control group to the SFOH group of nine patients. No significant differences in the basic characteristics (age, gender, ESKD cause, dialysis vintage, Kt/V, dialysis modality) existed between the two groups. All patients were dialyzed using polysulphone membranes with 1.8-2 m² surface area, blood flow rate of 300-350 mL/min, and dialysate flow rate of 600 mL/min. Fifty-percent of them were on online hemodiafiltration. All patients had a well-functioning vascular access and dialyzed adequately within the recommended targets (12 h weekly, spKt/V = 1.32-2.14). Dialysis fluid had the following composition:

Table 1 Baseline characteristics of the study population (*n* = 15)

	SFOH group, <i>n</i> = 9	Control group, <i>n</i> = 6	<i>P</i> value
Age (yr)	78 (35-90)	73 (49-91)	NS
Male/Female	9/0	4/2	NS
Cause of ESKD	4 DM, 2 GN, 2 UN, 1 ADPKD	2 DM, 2 HN, 1 GN, 1 UN	NS
Dialysis vintage (mo)	53 (19-131)	49 (21-67)	NS
Arteriovenous fistula/Graft	6/3	4/2	NS
Single pool Kt/V	1.56 ± 0.16	1.57 ± 0.29	NS
Dose of LanC pre switching (mg/d)	2250 (750-3000)	3000 (1500-3750)	NS
Number of LanC pills pre switching	3 (1-4)	4 (2-5)	NS
Calcium carbonate: <i>n</i> , dose (mg/d)	1, 750	1, 1250	NS
Cinacalcet: <i>n</i> , dose (mg/d)	1, 60	1, 30	NS
Paricalcitol: <i>n</i> , dose (mcg/wk)	6, 10 (7.5-15)	4, 5 (5-10)	NS

ADPKD: Adult dominant polycystic kidney disease; DM: Diabetes mellitus; ESKD: End stage kidney disease; GN: Glomerulonephritis; HN: Hypertensive nephrosclerosis; LanC: Lanthanum carbonate; NS: Non-significant; SFOH: Sucroferric oxyhydroxide.

Sodium: 138 mmol/L, Potassium: 2.0 mmol/L, Calcium: 1.25-1.75 mmol/L, Magnesium: 0.5 mmol/L, Chloride: 108.5-109.5 mmol/L, Acetate: 3 mmol/L, Bicarbonate: 32 mmol/L, and Glucose: 5.55 mmol/L. Dialysate composition and doses of vitamin D analogs, cinacalcet and CalC, as well as hemodialysis prescription (including bicarbonate conductivity), remained unchanged throughout the study for each patient. Ultrafiltration rates were adjusted to the individual needs of the patients. Dietary consultation regarding phosphorus, potassium and protein intake was the same before and during the study. The study was performed in accordance with the Declaration of Helsinki and with the approval of the hospital ethics committee. Informed written consent was obtained from all patients before they entered the study.

Blood analysis

Blood was sampled from the "arterial needle" of the vascular access in all studied patients before dialysis sessions. Samples were collected on the 3-d and the last 2-d interdialytic interval of the week prior to switching (Baseline) and six weeks after at the same intervals. Whole blood pH, pO₂, pCO₂, bicarbonate levels (HCO₃⁻), base excess (BE) and potassium (K⁺) were measured using a Cobas B221 blood gas analyzer (Roche Diagnostics Limited, CH-6343 Rotkreuz, Switzerland). Serum phosphate levels were measured the same days using standard laboratory methods. The arithmetic mean (average) of the two pre-switching (baseline) measurements (2 and 3-d interdialytic interval) and the respective mean of the two post-switching measurements (6 wk) were also calculated for each of the above mentioned parameters.

Statistical analysis

All data are presented as mean ± SD and median with (minimum-maximum), according to the distribution. Due to the small sample size, non-parametric tests were used. Differences in parameters between studied groups of patients were analyzed using the Wilcoxon signed-rank test for paired samples and the Mann-

Whitney *U* test for independent samples. Bivariate correlations were performed using Spearman's rank correlation analysis. Statistical significance was set at *P* < 0.05.

RESULTS

Patients who switched to SFOH (SFOH group) had an effective phosphate control (Table 2), without increasing the pill burden, [number of LanC pills: 3 (1-4) vs SFOH pills: 2 (1-6), *P* = 0.849]. Comparing the mean values of the measured acid-base balance parameters, pre- and post-switching, no difference was found in any of them (Table 2). No statistically significant differences were found even when we analyzed the data for the 3-d and the 2-d interdialytic intervals separately (Supplementary Tables 1 and 2), or when we performed the same analyses in the control group or between the two groups (data not shown).

No correlations were found, either between pre-switching daily LanC dose and the measured parameters, or between post-switching daily dose of the new binder and the measured parameters (data not shown).

The only significant differences that we found were between the 3-d and 2-d measurements. HCO₃⁻, BE, and pH were significantly lower and K⁺ was higher at the 3-d vs 2-d interdialytic interval, as expected. For example, in the whole group of patients, baseline 3-d HCO₃⁻ were 21.2 ± 2.17 mmol/L, BE: -3.5 (-4.8 to -2.3) mmol/L, pH: 7.373 ± 0.031, and K⁺: 4.69 ± 0.46 mmol/L. Respective values for the baseline 2-d interval were: HCO₃⁻: 22.53 ± 2.68 mmol/L (*P* = 0.005, comparing to 3-d), BE: -2.2 (-3.3 to -0.4) mmol/L (*P* = 0.001), pH 7.389 ± 0.029 (*P* = 0.003) and K⁺ 4.29 ± 0.42 mmol/L (*P* = 0.04). Accordingly, 3-d pCO₂ was lower (37.2 ± 3.34 mmHg) vs 2-d (38.19 ± 4.51 mmHg), however the difference did not reach statistical significance (*P* = 0.061). Phosphate levels were the same, 1.42 ± 0.4 mmol/L vs 1.46 ± 0.26 mmol/L (*P* = 0.801), indicating that phosphate binders are effective in maintaining stable phosphate levels all days. When we performed the same analyses in the post-switching

Table 2 Paired comparisons, pre- and post-switching in the two groups

	SFOH group, <i>n</i> = 9			Control group, <i>n</i> = 6		
	Baseline	6 wk	<i>P</i> value	Baseline	6 wk	<i>P</i> value
HCO ₃ ⁻ (mmol/L)	22.41 ± 1.66	22.62 ± 2.25	0.889	21.05 ± 3.14	21.12 ± 2.27	0.917
pH	7.38 ± 0.03	7.39 ± 0.03	0.635	7.37 ± 0.033	7.36 ± 0.019	0.917
pCO ₂ (mmHg)	38.41 ± 3.29	38.37 ± 3.62	0.767	36.62 ± 4.55	37.82 ± 4.97	0.173
pO ₂ (mmHg)	93.1 ± 12.04	97.4 ± 10.08	0.953	82.8 ± 11.47	79.9 ± 14.35	0.463
BE (mmol/L)	-2.55 (-3.8 to -0.98)	-2.9 (-3.92 to -0.22)	0.722	-3.27 (-5.6 to -2.1)	-3.85 (-5.5 to -2)	0.917
K ⁺ (mmol/L)	4.47 ± 0.32	4.38 ± 0.64	0.678	4.52 ± 0.23	4.6 ± 0.23	0.600
Phosphate (mmol/L)	1.57 ± 0.27	1.36 ± 0.38	0.214	1.34 ± 0.33	1.57 ± 0.48	0.249

The laboratory values refer to the average of the respective 2- and 3-d interdialytic interval and are expressed as mean ± SD or as median (min.-max.), *P* value denotes in group comparison (Wilcoxon signed rank test). SFOH: Suroferric oxyhydroxide; BE: Base excess.

measurements or in the different groups, the findings were similar (data not shown).

DISCUSSION

In our study, we found that when hemodialysis patients are switching from LanC to SFOH, despite the loss of carbonate provided by LanC, there is no significant consequence on acid-base balance. Several explanations may exist for this finding. First, the iron oxide hydroxide binds phosphate in the gastrointestinal tract through a direct ionic interaction between the negatively charged oxygen ions on the phosphate and the ferric ions in the ferric oxide, forming FePO₄^[13]. During this interaction, hydroxyl groups are released^[13], indicating that SFOH acts as a base. An additional explanation could be that the effect of the alkali contained in LanC is small and its loss is compensated by endogenous adaption from blood and bone buffering and residual renal function, when present. This may be particularly true when doses of LanC are not very high or when there is no severe acidosis, like in our patients. There was only one prospective study found where LanC was introduced in phosphate binder-naïve patients. There, serum bicarbonate levels remained stable after 9 mo of LanC in 28 patients with moderate CKD (21.9 ± 2.9 mmol/L at baseline vs 21.8 ± 2.4 mmol/L at 9 mo), indicating that the effect of the carbonate is small^[14]. Furthermore, it is well known that switching to LanC can ameliorate the acidosis caused by SevH^[7,8]. However, this may not be the result of the alkalinizing agent of LanC, but rather the consequence of the withdrawal of the hydrochloride of SevH. Treatment with SevH results in a significant increase in dietary acid load, and this is the main cause of the observed acidosis^[4]. In favor of this is the observation that patients who switch from SevH to bicalomer, a non-hydrochloride and non-carbonate phosphate binder, improve their metabolic acidosis. This indicates that the withdrawal of chloride without the additional effect of carbonate is sufficient to ameliorate the acidosis^[15]. Lastly, in our study, no correlation was found between LanC dose and acid-base parameters. The above support the hypothesis that the effect of the carbonate content of LanC on

acid-base status is rather small. Moreover, concomitant use of CalC could compensate for the effects of loss of carbonate provided by the LanC on acid-base status. However, only two of our patients were on CalC, and the doses remained stable throughout the study. Finally, due to the small sample size of our study, a type II error cannot be excluded. Nevertheless, the harmful effect of SevH on acid-base balance was shown even in studies with 8 to 16 patients^[7,15,16]. Thus, the effects of SFOH switching would probably be detected in our study if they were significant. In addition, finding a significant difference between the acid-base status of 2-d and 3-d interdialytic interval could be considered a supporting finding that despite the small sample size, differences were detected when they were significant.

In our study, SFOH was effective in controlling phosphate without increasing the pill burden, in accordance with previous studies^[17,18]. However, some of our patients could not tolerate the new binder, as treatment with SFOH appears to be predominantly complicated by gastrointestinal disturbances^[17,19].

Finally, we observed that patients were more acidotic in the long interdialytic interval. This has been already shown in previous studies^[20], and this is why we decided to perform measurements both in the long and the short interdialytic interval in an effort to increase the possibility of detecting potential effects of SFOH switching on acid-base status.

The main limitation of the present study is the small sample size, as possible effects of switching on acid-base parameters may not be detected, even if present. Yet, the small number of patients allowed for stable conditions and better compliance, and this, together with the prospective nature, can be considered a strength of the study. Another limitation is that we did not measure the protein dietary intake that can affect acid-base balance. However, nutritional instructions remained the same during the study, and albumin and cholesterol levels did not change, supporting that there were no major changes in dietary habits. Finally, we cannot generalize our findings in patients with severe acidosis or in patients with ESKD on peritoneal dialysis.

To the best of our knowledge, this is the first study that evaluates acid-base status in patients receiving the

new phosphate binder SFOH. Therefore, it can serve as a pilot for further studies with a larger number of patients, different binders and extended durations to better understand their effect in clinically important outcomes, such as metabolic acidosis.

In conclusion, switching from LanC to SFOH did not have any significant effect on blood bicarbonate levels and gas analysis, indicating that there is no need to change hemodialysis prescription regarding these parameters. However, when selecting a phosphate binder, potential consequences on acid-base balance should be considered, and monitoring of serum bicarbonate levels is part of good clinical practice.

ARTICLE HIGHLIGHTS

Research background

The effect of phosphate binders in clinically important outcomes, such as metabolic acidosis, remains understudied.

Research motivation

There are no studies examining the effect of the novel phosphate binder sucroferric oxyhydroxide (SFOH) on acid-base status.

Research objectives

Examine possible alterations in acid-base parameters in hemodialysis patients switching from lanthanum carbonate (LanC) to SFOH.

Research methods

Fifteen stable hemodialysis patients switched from LanC to SFOH. We compared pre- and post-switching blood gas analyses, whilst hemodialysis conditions and medications remained stable.

Research results

Switching from LanC to the novel phosphate binder SFOH did not have any significant effect on blood bicarbonate levels and gas analysis. No correlations were found, either between pre-switching LanC daily dose or between post-switching daily dose of the new binder and the measured parameters.

Research conclusions

This is the first study that evaluates acid-base status in patients switching from LanC to the new phosphate binder SFOH, showing that there is no need to change hemodialysis prescription regarding these parameters.

Research perspectives

Our study can serve as a pilot for further studies with a larger number of patients, different binders and extended durations, to better understand their effect in clinically important outcomes, such as metabolic acidosis.

REFERENCES

- Kraut JA, Madias NE. Metabolic Acidosis of CKD: An Update. *Am J Kidney Dis* 2016; **67**: 307-317 [PMID: 26477665 DOI: 10.1053/j.ajkd.2015.08.028]
- Bommer J, Locatelli F, Satayathum S, Keen ML, Goodkin DA, Saito A, Akiba T, Port FK, Young EW. Association of predialysis serum bicarbonate levels with risk of mortality and hospitalization in the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Am J Kidney Dis* 2004; **44**: 661-671 [PMID: 15384017]
- Thet Z, Win AK, Pedagogos E, Beavis J, Crikis S, Nelson C. Differential effects of phosphate binders on pre-dialysis serum bicarbonate in end-stage kidney disease patients on maintenance haemodialysis. *BMC Nephrol* 2013; **14**: 205 [PMID: 24079654 DOI: 10.1186/1471-2369-14-205]
- Brezina B, Qunibi WY, Nolan CR. Acid loading during treatment with sevelamer hydrochloride: mechanisms and clinical implications. *Kidney Int Suppl* 2004; S39-S45 [PMID: 15296506 DOI: 10.1111/j.1523-1755.2004.09007.x]
- Oka Y, Miyazaki M, Matsuda H, Takatsu S, Katsube R, Mori T, Takehara K, Umeda Y, Uno F. Sevelamer hydrochloride dose-dependent increase in prevalence of severe acidosis in hemodialysis patients: analysis of nationwide statistical survey in Japan. *Ther Apher Dial* 2014; **18**: 37-43 [PMID: 24499082 DOI: 10.1111/1744-9987.12052]
- Matsushita Y, Yamnouchi E, Matsuoka K, Arizono K. Effect of calcium carbonate administration on metabolic acidosis induced by sevelamer hydrochloride in chronic hemodialysis patients. *J Jpn Soc Dial Ther* 2006; **39**: 1245-1250 [DOI: 10.4009/jstd.39.1245]
- Filiopoulos V, Koutis I, Trompouki S, Hadjiyannakos D, Lazarou D, Vlassopoulos D. Lanthanum carbonate versus sevelamer hydrochloride: improvement of metabolic acidosis and hyperkalemia in hemodialysis patients. *Ther Apher Dial* 2011; **15**: 20-27 [PMID: 21272248 DOI: 10.1111/j.1744-9987.2010.00868.x]
- Pai AB, Shepler BM. Comparison of sevelamer hydrochloride and sevelamer carbonate: risk of metabolic acidosis and clinical implications. *Pharmacotherapy* 2009; **29**: 554-561 [PMID: 19397463 DOI: 10.1592/phco.29.5.554]
- Fan S, Ross C, Mitra S, Kalra P, Heaton J, Hunter J, Plone M, Pritchard N. A randomized, crossover design study of sevelamer carbonate powder and sevelamer hydrochloride tablets in chronic kidney disease patients on haemodialysis. *Nephrol Dial Transplant* 2009; **24**: 3794-3799 [PMID: 19666658 DOI: 10.1093/ndt/gfp372]
- Habbous S, Przech S, Acedillo R, Sarma S, Garg AX, Martin J. The efficacy and safety of sevelamer and lanthanum versus calcium-containing and iron-based binders in treating hyperphosphatemia in patients with chronic kidney disease: a systematic review and meta-analysis. *Nephrol Dial Transplant* 2017; **32**: 111-125 [PMID: 27651467 DOI: 10.1093/ndt/gfw312]
- Cozzolino M, Funk F, Rakov V, Phan O, Teitelbaum I. Preclinical Pharmacokinetics, Pharmacodynamics and Safety of Sucroferric Oxyhydroxide. *Curr Drug Metab* 2014; **15**: 953-965 [PMID: 25658128]
- Cernaro V, Santoro D, Lacquaniti A, Costantino G, Visconti L, Buemi A, Buemi M. Phosphate binders for the treatment of chronic kidney disease: role of iron oxyhydroxide. *Int J Nephrol Renovasc Dis* 2016; **9**: 11-19 [PMID: 26893577 DOI: 10.2147/IJNRD.S78040]
- Velphoro Drug Approval Package FDA. [Accessed 13 Nov 2017]. Available from: URL: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2013/205109Orig1s000ChemR.pdf
- Block GA, Wheeler DC, Persky MS, Kestenbaum B, Ketteler M, Spiegel DM, Allison MA, Asplin J, Smits G, Hoofnagle AN, Kooienga L, Thadhani R, Mannstadt M, Wolf M, Chertow GM. Effects of phosphate binders in moderate CKD. *J Am Soc Nephrol* 2012; **23**: 1407-1415 [PMID: 22822075 DOI: 10.1681/ASN.2012030223]
- Sanai T, Tada H, Ono T, Fukumitsu T. Phosphate binders and metabolic acidosis in patients undergoing maintenance hemodialysis—sevelamer hydrochloride, calcium carbonate, and bicalomer. *Hemodial Int* 2015; **19**: 54-59 [PMID: 24980286 DOI: 10.1111/hdi.12188]
- Gallieni M, Cozzolino M, Brancaccio D. Transient decrease of serum bicarbonate levels with Sevelamer hydrochloride as the phosphate binder. *Kidney Int* 2000; **57**: 1776-1777 [PMID: 10809607 DOI: 10.1046/j.1523-1755.2000.00025.x]
- Floege J, Covic AC, Ketteler M, Rastogi A, Chong EM, Gaillard S, Lisk LJ, Sprague SM; PA21 Study Group. A phase III study of the efficacy and safety of a novel iron-based phosphate binder in dialysis patients. *Kidney Int* 2014; **86**: 638-647 [PMID: 24646861 DOI: 10.1038/ki.2014.58]
- Coyne DW, Ficociello LH, Parameswaran V, Anderson L, Vemula S, Ofsthun NJ, Mullon C, Maddux FW, Kossmann RJ, Sprague SM. Real-world effectiveness of sucroferric oxyhydroxide in

- patients on chronic hemodialysis: A retrospective analysis of pharmacy data. *Clin Nephrol* 2017; **88**: 59-67 [PMID: 28587714 DOI: 10.5414/CN109021]
- 19 **Wüthrich RP**, Chonchol M, Covic A, Gaillard S, Chong E, Tumlin JA. Randomized clinical trial of the iron-based phosphate binder PA21 in hemodialysis patients. *Clin J Am Soc Nephrol* 2013; **8**: 280-289 [PMID: 23124782 DOI: 10.2215/CJN.08230811]
- 20 **Graham KA**, Hoenich NA, Goodship TH. Pre and interdialytic acid-base balance in hemodialysis patients. *Int J Artif Organs* 2001; **24**: 192-196 [PMID: 11394698]
- P- Reviewer:** Choi MR, Ekpenyong CEE, Kilinc M, Nechifor G, Tiwari S
S- Editor: Cui LJ **L- Editor:** Filipodia **E- Editor:** Huang Y





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

