

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 4,610 条结果 (用时 0.73 秒)

Phase II study of adjuvant chemotherapy of S-1 plus oxaliplatin for ...

<https://www.ncbi.nlm.nih.gov/pubmed/26626800> - 翻译此页

作者: K Shitara - 2017 - 被引用次数: 19 - 相关文章

Phase II study of adjuvant chemotherapy of S-1 plus oxaliplatin for patients with stage III gastric cancer after D2 gastrectomy. ... for Gastric Cancer (ACTS-GC) demonstrated a survival benefit by adjuvant S-1 ... resection of stage II/III gastric cancer, but there is still a need to improve the efficacy of treatment of stage III disease.

Effect of Early Adjuvant Chemotherapy on Survival of Advanced ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5881011/> - 翻译此页

作者: Y Lee - 2018

2018年3月20日 - In this study, we investigated the effect of early AC on the survival of stage ... After propensity score matching, RFS was similar between early and late ... Patients who undergo curative D2 gastrectomy for pathologic stage II or III gastric cancer are ... (S-1) for 1 year or capecitabine plus intravenous oxaliplatin ...

Three-year outcomes of two phase II studies of adjuvant ...

ascopubs.org/doi/abs/10.1200/JCO.2018.36.4_suppl.94 - 翻译此页

作者: Y Nakamura - 2018 - 被引用次数: 1

Three-year outcomes of two phase II studies of adjuvant chemotherapy with S-1 plus oxaliplatin or capecitabine plus oxaliplatin in patients with stage III gastric ...

Efficacy of Adjuvant S-1 Versus XELOX Chemotherapy for Patients ...

https://www.researchgate.net/.../323205198_Efficacy_of_Adjuvant_S-1_Ver... - 翻译此页

Background: After curative resection of gastric cancer with D2 lymph node dissection, ... adjuvant chemotherapy with S-1 or capecitabine plus oxaliplatin (XELOX) is ... This study aimed to compare the efficacies of adjuvant S-1 and XELOX ... After propensity score-matching, the 3-year disease-free survival rate (DFS) was ...

Gastrointestinal (Noncolorectal) Cancer : Esophageal or Gastric Cancer

abstracts.asco.org/214/CatAbstView_214_553_AT.html - 翻译此页

A phase II study of perioperative intraperitoneal oxaliplatin plus S-1/oxaliplatin for ... in advanced gastric

Match Overview

1	Internet 67 words crawled on 22-May-2014 www.isesnet.org	1%
2	Crossref 67 words Qu, Jing-lei, Xin Li, Xiu-juan Qu, Zhi-tu Zhu, Li-zhong Zhou, Yue-e Teng, Jing-dong Zhang, Bo Jin, Ming-fang Zhao,	1%
3	Crossref 57 words Shitara, Kohei, Keisho Chin, Takaki Yoshikawa, Hitoshi Katai, Masanori Terashima, Seiji Ito, Motohiro Hirao, Kazuhiro	1%
4	Internet 45 words crawled on 18-Mar-2016 www.sabcs.org	1%
5	Crossref 40 words "Saturday 27 August 2016", European Heart Journal, 2016	1%
6	Crossref 28 words Lordick, Florian, and Masanori Terashima. "Gastric Cancer Adjuvant Therapy", Best Practice & Research Clinical Gast	<1%
7	Crossref 27 words Guo-Shuang Shen, Jiu-Da Zhao, Jun-Hui Zhao, Xin-Fu Ma et al. "Association of HER2 status with prognosis in gastric	<1%
8	Internet 16 words crawled on 11-May-2016 eest.com	<1%

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 40026

Manuscript Type: BASIC ARTICLE

Retrospective Study

³ Adjuvant chemotherapy with S-1 plus oxaliplatin improves survival of patients with gastric cancer after D2 gastrectomy: A multicenter propensity score-matched study

Ren DF *et al.* Adjuvant S-1 plus Oxaliplatin for GC

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 5,420 条结果 (用时 0.70 秒)

Phase II study of adjuvant chemotherapy of S-1 plus oxaliplatin for ...

<https://www.ncbi.nlm.nih.gov/pubmed/26626800> ▼ [翻译此页](#)

作者 : K Shitara - 2017 - 被引用次数 : 20 - 相关文章

2017年12月1日 - 2017 Jan;20(1):175-181. doi: 10.1007/s10120-015-0581-1. ... S-1 plus oxaliplatin for patients with stage III gastric cancer after D2 ... a survival benefit by adjuvant S-1 monotherapy in patients who had undergone curative resection of stage II/III gastric cancer, but there is still a need to improve the efficacy of ...

Adjuvant capecitabine plus oxaliplatin after D2 gastrectomy in ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/26956689> ▼ [翻译此页](#)

作者 : N Fuse - 2017 - 被引用次数 : 17 - 相关文章

1. Gastric Cancer. 2017 Mar;20(2):332-340. doi: ... after D2 gastrectomy in Japanese patients with gastric cancer: a phase II study. Fuse N(1), Bando H(2), Chin K(3), Ito S(4), Yoshikawa T(5), Tsuburaya A(5), ... Adjuvant chemotherapy with XELOX (capecitabine plus oxaliplatin) has ... Clinical Trial, Phase II; Multicenter Study ...

Chemoradiotherapy versus chemotherapy as adjuvant treatment for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5883367/> - [翻译此页](#)

作者 : DM Girardi - 2018 - 被引用次数 : 1 - 相关文章

2018年4月3日 - In this study, we have used a propensity score matched analysis to retrospectively evaluate The pivotal Japanese ACTS-GC study evaluated post-operative S-1, an oral Adjuvant capecitabine plus oxaliplatin for gastric cancer after D2 ... Adjuvant radiotherapy improves overall survival in patients with ...

Gastrointestinal (Noncolorectal) Cancer : Esophageal or Gastric Cancer

abstracts.asco.org/214/CatAbstView_214_553_AT.html ▼ [翻译此页](#)

A phase II study of perioperative intraperitoneal paclitaxel plus S-1/paclitaxel for ... in advanced gastric cancer patients treated with S-1 adjuvant chemotherapy. ... Can chemoradiotherapy improve survival after an R1 resection for gastric cancer: A ... A multicenter case-control study with propensity score matching analysis.

找到约 5,150 条结果 (用时 0.62 秒)

Adjuvant capecitabine plus oxaliplatin after D2 gastrectomy in ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/26956689> ▼ 翻译此页

作者: N Fuse - 2017 - 被引用次数: 17 - 相关文章

2018年3月8日 - 1. *Gastric Cancer*. 2017 Mar;20(2):332-340. doi: ... after D2 gastrectomy in Japanese patients with gastric cancer: a phase II study. Fuse N(1), Bando H(2), Chin K(3), Ito S(4), Yoshikawa T(5), ... Adjuvant chemotherapy with XELOX (capecitabine plus oxaliplatin) ... Clinical Trial, Phase II; Multicenter Study ...

Phase II study of adjuvant chemotherapy of S-1 plus oxaliplatin for ...

<https://www.ncbi.nlm.nih.gov/pubmed/26626800> ▼ 翻译此页

作者: K Shitara - 2017 - 被引用次数: 20 - 相关文章

2017年12月1日 - 1. *Gastric Cancer*. 2017 Jan;20(1):175-181. doi: ... of S-1 plus oxaliplatin for patients with stage III gastric cancer after D2 ... a survival benefit by adjuvant S-1 monotherapy in patients who had ... II/III gastric cancer, but there is still a need to improve the efficacy of ... Clinical Trial, Phase II; Multicenter Study ...

Selective Gastric Cancer Patients with Peritoneal Seeding Benefit from ...

www.jcancer.org/v08p2231.htm ▼ 翻译此页

作者: SQ Yuan - 被引用次数: 1 - 相关文章

2017年7月20日 - Propensity score matching (PSM) was performed to balance the selection bias. Results: After PSM, compared with non-gastrectomy group, ... standardized D2 lymphadenectomy and adjuvant chemotherapy [2-4], the overall ... to improve the prognosis of gastric cancer patients with peritoneal seeding [6-8].

Association of Adjuvant Chemotherapy With Survival in Patients With ...

<https://jamanetwork.com/journals/jamasurgery/fullarticle/2627289> - 翻译此页

作者: Y Jiang - 2017 - 被引用次数: 4 - 相关文章

2017年5月24日 - This study analyzes patient and tumor characteristics to create a model that ... patients with stage II or stage III gastric cancer, 2 nomograms and 1 Propensity score matching analysis was performed for receiving the plus oxaliplatin with observation after D2 gastrectomy for patients with stage II or III GC.

[PDF] Effect of Early Adjuvant Chemotherapy on Survival of Advanced ...

<https://jgc-online.org/Synapse/Data/PDFData/1100JGC/jgc-18-58.pdf> ▼ 翻译此页

作者: Y Lee - 2018 - 相关文章

2018年3月13日 - in the early AC group (n=174) were matched 1:1 with patients in the ... Patients who undergo curative D2 gastrectomy for pathologic stage II or III gastric ... of AC were used in this study: