



Predictive factors for lymph node metastasis and defining a subgroup treatable



全部

新闻

图片

购物

地图

更多

设置

工具

找到约 7,490 条结果 (用时 0.44 秒)

Defining a Subgroup Treatable for Laparoscopic Surgery in Poorly ...

https://www.researchgate.net/.../236925627_Defining_a_Subgroup_Treatabl... - 翻译此页

2018年4月9日 - The present study aims to identify the clinicopathologic factors predictive of lymph node metastasis (LNM) in poorly differentiated early gastric ...

Predictive factors of lymph node metastasis in intramucosal poorly ...

https://www.researchgate.net/.../257495018_Predictive_factors_of_lymph_n... - 翻译此页

2018年2月12日 - ... clinicopathological factors predictive of lymph node metastasis (LNM) in intramucosal poorly differentiated early gastric cancer (EGC), and further to expand the possibility of using laparoscopic surgery for the treatment of. ... Defining a Subgroup Treatable for Laparoscopic Surgery in Poorly Differentiated ...

Predictive factors for lymph node metastasis in early gastric cancer

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2975097/> - 翻译此页

作者: CM Sung - 2010 - 被引用次数: 39 - 相关文章

2010年11月7日 - Predictive factors for lymph node metastasis in early gastric cancer which subgroup of EGC patients could be treated with EMR or ESD instead of radical surgery. ... Poorly differentiated adenocarcinomas, signet ring cell carcinomas, Y. Survival after surgical treatment of early gastric cancer: surgical ...

Predictive Factors for Lymph Node Metastasis in Signet Ring Cell ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3705138/> - 翻译此页

作者: JY Kim - 2013 - 被引用次数: 14 - 相关文章

2013年6月25日 - The lymph node metastasis rate for signet ring cell carcinoma was 4.3% (n=8). ... undifferentiated gastric carcinoma includes poorly differentiated adenocarcinoma ... application of ESD in a suitable subgroup of patients with early SRC. ... SRC was defined pathologically as an adenocarcinoma in which the ...

A Risk Prediction Model Based on Lymph-Node Metastasis in Poorly ...

journals.plos.org/plosone/article?id=10.1371/journal.pone.0156207 - 翻译此页

作者: JH Pyo - 2016 - 被引用次数: 5 - 相关文章

2016年5月26日 - A risk model based on independent predicting factors of LNM was ... characteristic curve for predicting LNM after internal validation was 0.69 ... lymph-node metastasis (LNM) in intramucosal poorly differentiated adenocarcinomas (PDC). Incidence of lymph node metastasis from early gastric cancer: ...

缺少字词: subgroup lled

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 40553

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Predictive factors for lymph node metastasis and defining a subgroup treatable for laparoscopic lymph node dissection in poorly differentiated early gastric cancer

Hua Li, Zhi-Bin Huo, Qing-Qiang He, Fan-Ting Kong, Yun-He Gao, Wen-Quan Liang, Deng-Xiang Liu

Abstract

Match Overview

| | | |
|---|---|-----|
| 1 | Crossref 51 words Hua Li, Ping Lu, Yang Lu, Caigang Liu, Huimian Xu, Shubo Wang, Junqing Chen. "Predictive factors of lymph node m | 2% |
| 2 | Crossref 26 words Li, Hua. "Predictive factors for lymph node metastasis in poorly differentiated early gastric cancer and their impact on ... | 1% |
| 3 | Crossref 11 words Chen Li. "Risk Factors for Lymph Node Metastasis in Unifocally Differentiated Early Gastric Cancer", Annals of Surgical Oncol | 1% |
| 4 | Crossref 10 words Hua Li. "Lymph node metastasis in early signet ring cell gastric carcinoma: Endoscopic mucosal resection", Chinese Journal | <1% |

找到约 87,000 条结果 (用时 0.54 秒)

Defining a Subgroup Treatable for Laparoscopic Surgery in Poorly ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3643644/> - 翻译此页

作者: Z Huo - 2012 - 被引用次数: 4 - 相关文章

The present study aims to identify the clinicopathologic factors predictive of lymph node metastasis (LNM) in poorly differentiated early gastric cancer (EGC) and ...

Defining a Subgroup Treatable for Laparoscopic Surgery in Poorly ...

https://www.researchgate.net/.../236925627_Defining_a_Subgroup_Treatabl... - 翻译此页

2018年4月9日 - differentiated early gastric cancer (EGC) and to expand the possibility of using laparoscopic surgery for the treatment of poorly ... gastric cancer, lymph nodes, metastasis, laparoscopy ... clinicopathologic factors that are predictive of LNM in poorly ... lymph node dissection beyond limited (D1) dissection was.

Predictive Factors for Lymph Node Metastasis in Undifferentiated Early ...

https://www.researchgate.net/.../312889215_Predictive_Factors_for_Lymph... - 翻译此页

Predictive Factors for Lymph Node Metastasis in Undifferentiated Early ... Objectives: Less invasive surgery is gaining popularity for the treatment of early gastric cancer (EGC). ... Even for undifferentiated EGC, such as poorly differentiated adenocarcinoma ... Defining a Subgroup Treatable for Laparoscopic and Endoscopic ...

Defining a Subgroup Treatable for Laparoscopic Surgery in Poorly ...

www.cancerbiomed.org > Home > Vol 9, No 1 ▼ 翻译此页

作者: Z Huo - 2012 - 被引用次数: 4 - 相关文章

Defining a Subgroup Treatable for Laparoscopic Surgery in Poorly Differentiated ... factors predictive of lymph node metastasis (LNM) in poorly differentiated early gastric ... Keywords: gastric cancer; lymph nodes; metastasis; laparoscopy ... Early gastric cancer (EGC) is defined as gastric carcinoma in which the invasion is ...

A nomogram for predicting the likelihood of lymph node metastasis in ...

<https://bmccancer.biomedcentral.com/articles/10.1186/s12885-016-2132-5> ▼ 翻译此页

作者: Z Zheng - 2016 - 被引用次数: 19 - 相关文章

2016年2月12日 - Early gastric cancer is defined as a lesion confined to the mucosa or ... Early gastric cancer; Lymph node metastasis; Nomogram ... Currently, although gastrectomy plus lymph node dissection is still the gold standard of treatment for ... two groups: the differentiated group, which included well or moderately ...

A Risk Prediction Model Based on Lymph-Node Metastasis in Poorly ...