

Retrospective Study

Effect of primary tumor side on survival outcomes in metastatic colorectal cancer patients after hepatic artery infusion chemotherapy

Zhang HY *et al.* mCRC treated by HAIC

Hang-Yu Zhang, Jian-Hai Guo, Song Gao, Hui Chen, Xiao-Dong Wang,

Peng-Jun Zhang, Peng Liu, Guang Cao, Hai-Feng Xu, Lin-Zhong Zhu, Ren-Jie

Yang, Jian Li, Xu Zhu

Match Overview

1	Internet 22 words crawled on 20-Sep-2009 www.biomedcentral.com	1%
2	Internet 19 words crawled on 09-Apr-2016 www.karger.com	1%
3	Crossref 19 words Annie Lim, Samuel Le Sourd, H�el�ene Senellart, Dominique Luet et al. "Hepatic Arterial Infusion Chemotherapy for	1%
4	Internet 17 words crawled on 19-Aug-2016 www.intrinsiccro.com	1%
5	Internet 15 words crawled on 27-Jun-2018 www.cancer.gov	1%
6	Crossref 14 words Groote, Kerlijne De. "Intrahepatic therapy for liver-dominant metastatic colorectal cancer", World Journal of Gastr	<1%
7	Internet 13 words crawled on 16-Jun-2018 springerplus.springeropen.com	<1%
8	Internet 13 words crawled on 03-Dec-2016 atvb.ahajournals.org	<1%
9	Crossref 13 words Boisen, M. K., J. S. Johansen, C. Dehlendorff, J. S. Larsen, K. Osterlind, J. Hansen, S. E. Nielsen, P. Pfeiffer, L. S. T	<1%
10	Crossref 12 words Yoshikazu Kawazuma. "Improvement of Survival Outcome for Colon Cancer Patients by Anatomical Sub-sites",	<1%



Effect of Primary Tumor Side on Survival Outcomes in Metastatic Colorec



全部

图片

新闻

购物

地图

更多

设置

工具

找到约 246,000 条结果 (用时 0.56 秒)

Effect of Primary Tumor Side on Survival Outcomes in Untreated ...

<https://www.ncbi.nlm.nih.gov/pubmed/30033117> - 翻译此页

作者 : P Gibbs - 2018 - 相关文章

Effect of Primary Tumor Side on Survival Outcomes in Untreated Patients With Metastatic Colorectal Cancer When Selective Internal Radiation Therapy Is Added to Chemotherapy: Combined Analysis of Two Randomized Controlled Studies. ... primary (RSP) tumors and those with left-sided primary (LSP) tumors after ...

缺少字词 : artery infusion

An Update on Hepatic Arterial Infusion Chemotherapy for Colorectal ...

theoncologist.alphamedpress.org/content/8/6/553.full ▼ 翻译此页

作者 : AD Cohen - 2003 - 被引用次数 : 138 - 相关文章

Hepatic metastases are a frequent complication of colorectal cancer (CRC), affecting ... Prognosis is dismal for untreated patients, with median survival times of 6 to 12 ... intraarterially via the side port of the hepatic arterial infusion pump, and both ... adjuvant chemotherapy following resection of their primary CRC, and 15% ...

Hepatic artery infusion therapy is effective for chemotherapy-resistant ...

<https://wjso.biomedcentral.com/articles/10.1186/s12957-015-0704-5> ▼ 翻译此页

作者 : T Goi - 2015 - 被引用次数 : 3 - 相关文章

2015年10月9日 - The subjects had liver metastatic colorectal cancers that were unresponsive to ... tomography imaging findings obtained after hepatic artery infusion. ... decades, the survival prognosis of unresectable or recurrent colorectal cancer ... However, in cases of solitary hepatic tumors, hepatic artery infusion can ...

Hepatic Artery Infusion Pump Chemotherapy With Floxuridine and ...

<https://clinicaltrials.gov/ct2/show/NCT03366155> ▼ 翻译此页

2017年12月8日 - Of patients with metastatic colorectal cancer, the liver will be the sole site ... Hepatic artery infusion of floxuridine (FUDR) via an implantable ... Primary Outcome Measures : ... The fraction of patients whose tumors in liver shrunk after therapy ... for teratogenic or abortifacient effects of the HAIP chemotherapy.

缺少字词 : side

找到约 471,000 条结果 (用时 0.55 秒)

Effect of Primary Tumor Side on Survival Outcomes in Untreated ...

<https://www.ncbi.nlm.nih.gov/pubmed/30033117> - 翻译此页

作者 : P Gibbs - 2018 - 相关文章

Effect of Primary Tumor Side on Survival Outcomes in Untreated Patients With Metastatic Colorectal Cancer When Selective Internal Radiation Therapy Is Added to Chemotherapy: Combined Analysis of Two Randomized Controlled Studies. ... primary (RSP) tumors and those with left-sided primary (LSP) tumors after ...

缺少字词 : artery infusion

An Update on Hepatic Arterial Infusion Chemotherapy for Colorectal ...

theoncologist.alphamedpress.org/content/8/6/553.full ▼ 翻译此页

作者 : AD Cohen - 2003 - 被引用次数 : 138 - 相关文章

Hepatic metastases are a frequent complication of colorectal cancer (CRC), affecting ... Prognosis is dismal for untreated patients, with median survival times of 6 to 12 ... intraarterially via the side port of the hepatic arterial infusion pump, and both ... adjuvant chemotherapy following resection of their primary CRC, and 15% ...

Hepatic Artery Infusion Pump Chemotherapy With Floxuridine and ...

<https://clinicaltrials.gov/ct2/show/NCT03366155> ▼ 翻译此页

2017年12月8日 - Of patients with metastatic colorectal cancer, the liver will be the sole site ... Hepatic artery infusion of floxuridine (FUDR) via an implantable ... Primary Outcome Measures : ... The fraction of patients whose tumors in liver shrunk after therapy ... for teratogenic or abortifacient effects of the HAIP chemotherapy.

缺少字词 : side

Hepatic artery infusion therapy is effective for chemotherapy-resistant ...

<https://wjso.biomedcentral.com/articles/10.1186/s12957-015-0704-5> ▼ 翻译此页

作者 : T Goi - 2015 - 被引用次数 : 3 - 相关文章

2015年10月9日 - The subjects had liver metastatic colorectal cancers that were unresponsive to ... tomography imaging findings obtained after hepatic artery infusion. ... decades, the survival prognosis of unresectable or recurrent colorectal cancer ... However, in cases of solitary hepatic tumors, hepatic artery infusion can ...

Hepatic arterial infusion in the management of colorectal cancer liver ...

<https://www.sciencedirect.com/science/article/pii/S159086581731321X> - 翻译此页

作者 : N Chapelle - 2017 - 被引用次数 : 1 - 相关文章

Overall, 20-60% of patients with colorectal cancer (CRC) will develop ... Five years overall survival (OS) in