

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2018 December 14; 24(46): 5189-5296



**GUIDELINES**

- 5189** Chinese consensus on management of tyrosine kinase inhibitor-associated side effects in gastrointestinal stromal tumors

*Li J, Wang M, Zhang B, Wu X, Lin TL, Liu XF, Zhou Y, Zhang XH, Xu H, Shen LJ, Zou J, Lu P, Zhang D, Gu WJ, Zhang MX, Pan J, Cao H; Chinese Society of Surgeons for Gastrointestinal Stromal Tumor of the Chinese Medical Doctor Association*

**MINIREVIEWS**

- 5203** Liver transplantation for critically ill cirrhotic patients: Overview and pragmatic proposals

*Artzner T, Michard B, Besch C, Levesque E, Faitot F*

- 5215** Management of sub-centimeter recurrent hepatocellular carcinoma after curative treatment: Current status and future

*Lee MW, Lim HK*

**ORIGINAL ARTICLE****Basic Study**

- 5223** Modulation of faecal metagenome in Crohn's disease: Role of microRNAs as biomarkers

*Rojas-Feria M, Romero-García T, Fernández Caballero-Rico JÁ, Pastor Ramírez H, Avilés-Recio M, Castro-Fernandez M, Chueca Porcuna N, Romero-Gómez M, García F, Grande L, Del Campo JA*

- 5234** MicroRNA-15a - cell division cycle 42 signaling pathway in pathogenesis of pediatric inflammatory bowel disease

*Tang WJ, Peng KY, Tang ZF, Wang YH, Xue AJ, Huang Y*

- 5246** Effect of photodynamic therapy with (17R,18R)-2-(1-hexyloxyethyl)-2-devinyl chlorine E6 trisodium salt on pancreatic cancer cells *in vitro* and *in vivo*

*Shen YJ, Cao J, Sun F, Cai XL, Li MM, Zheng NN, Qu CY, Zhang Y, Shen F, Zhou M, Chen YW, Xu LM*

- 5259** Identification and prediction of novel non-coding and coding RNA-associated competing endogenous RNA networks in colorectal cancer

*Liang Y, Zhang C, Ma MH, Dai DQ*

**Retrospective Study**

- 5271** Relationship between response to lusutrombopag and splenic volume

*Uojima H, Arase Y, Itokawa N, Atsukawa M, Satoh T, Miyazaki K, Hidaka H, Sung JH, Kako M, Tsuruya K, Kagawa T, Iwakiri K, Horie R, Koizumi W*

- 5280** Preliminary application of 3D-printed coplanar template for iodine-125 seed implantation therapy in patients with advanced pancreatic cancer

*Huang W, Lu J, Chen KM, Wu ZY, Wang QB, Liu JJ, Gong J, Chen ZJ, Ding XY, Wang ZM*

**Observational Study**

- 5288** Loss of efficacy and safety of the switch from infliximab original to infliximab biosimilar (CT-P13) in patients with inflammatory bowel disease

*Guerra Veloz MF, Argüelles-Arias F, Castro Laria L, Maldonado Pérez B, Benítez Roldán A, Perea Amarillo R, Merino Bohórquez V, Calleja MA, Caunedo Álvarez Á, Vilches Arenas Á*

**ABOUT COVER**

Editorial board member of *World Journal of Gastroenterology*, İlhami Yuksel, MD, Professor, Department of Gastroenterology, Yildirim Beyazıt University School of Medicine, Ankara 06100, Turkey

**AIMS AND SCOPE**

*World Journal of Gastroenterology* (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, and 28<sup>th</sup> each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

**INDEXING/ABSTRACTING**

*World Journal of Gastroenterology* (*WJG*) is now indexed in Current Contents<sup>®</sup>/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch<sup>®</sup>), Journal Citation Reports<sup>®</sup>, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2018 edition of Journal Citation Reports<sup>®</sup> cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35<sup>th</sup> among 80 journals in gastroenterology and hepatology (quartile in category Q2).

**EDITORS FOR THIS ISSUE**

**Responsible Assistant Editor:** *Xiang Li*  
**Responsible Electronic Editor:** *Shu-Yu Yin*  
**Proofing Editor-in-Chief:** *Lian-Sheng Ma*

**Responsible Science Editor:** *Rao-Yu Ma*  
**Proofing Editorial Office Director:** *Ze-Mao Gong*

**NAME OF JOURNAL**

*World Journal of Gastroenterology*

**ISSN**

ISSN 1007-9327 (print)  
ISSN 2219-2840 (online)

**LAUNCH DATE**

October 1, 1995

**FREQUENCY**

Weekly

**EDITORS-IN-CHIEF**

**Andrzej S Tarnawski, MD, PhD, DSc (Med),**  
**Professor of Medicine, Chief Gastroenterology, VA**  
Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach, CA 90822, United States

**EDITORIAL BOARD MEMBERS**

All editorial board members resources online at <https://www.wjgnet.com/1007-9327/editorialboard.htm>

**EDITORIAL OFFICE**

Ze-Mao Gong, Director  
*World Journal of Gastroenterology*  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
Help Desk: <https://www.fjgpublishing.com/helpdesk>  
<https://www.wjgnet.com>

**PUBLISHER**

Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <https://www.fjgpublishing.com/helpdesk>  
<https://www.wjgnet.com>

**PUBLICATION DATE**

December 14, 2018

**COPYRIGHT**

© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

**SPECIAL STATEMENT**

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

**INSTRUCTIONS TO AUTHORS**

Full instructions are available online at <https://www.wjgnet.com/bpg/gerinfo/204>

**ONLINE SUBMISSION**

<https://www.fjgpublishing.com>

## Observational Study

# Loss of efficacy and safety of the switch from infliximab original to infliximab biosimilar (CT-P13) in patients with inflammatory bowel disease

María Fernanda Guerra Veloz, Federico Argüelles-Arias, Luisa Castro Laria, Belén Maldonado Pérez, Antonio Benítez Roldan, Raúl Perea Amarillo, Vicente Merino Bohórquez, Miguel Angel Calleja, Ángel Caunedo Álvarez, Ángel Vilches Arenas

María Fernanda Guerra Veloz, Federico Argüelles-Arias, Luisa Castro Laria, Belén Maldonado Pérez, Antonio Benítez Roldan, Raúl Perea Amarillo, Ángel Caunedo Álvarez, Department of Gastroenterology, University Hospital Virgen Macarena, Seville 41007, Spain

Vicente Merino Bohórquez, Miguel Angel Calleja, Pharmacy Unit, University Hospital Virgen Macarena, Seville 41007, Spain

Ángel Vilches Arenas, Preventive Medicine and Public Health, University Hospital Virgen Macarena, Seville 41007, Spain

**ORCID number:** María Fernanda Guerra Veloz (0000-0001-9708-8245); Federico Argüelles-Arias (0000-0003-3646-8919); Luisa Castro Laria (0000-0002-7971-9303); Belén Maldonado Pérez (0000-0001-8025-1472); Antonio Benítez Roldan (0000-0001-5662-2544); Raúl Perea Amarillo (0000-0003-1447-9457); Vicente Merino Bohórquez (0000-0002-0315-3411); Miguel Angel Calleja (0000-0001-8449-5490); Ángel Caunedo Álvarez (0000-0003-4444-233X); Ángel Vilches Arenas (0000-0003-2197-3760).

**Author contributions:** Guerra Veloz MF did the study design, data analysis and writing up of the first draft of the paper; Argüelles-Arias F did the data analysis and patient recruitment; Castro Laria L and Maldonado Pérez B did the patient recruitment; Perea Amarillo R and Merino Bohórquez V did the Data collection; Caunedo Álvarez Á did the study design; Vilches Arenas Á did the data analysis.

**Institutional review board statement:** The study was approved by the Research Ethics Committee of the Virgen Macarena Hospital, Seville, Spain.

**Informed consent statement:** All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

**Conflict-of-interest statement:** Guerra Veloz MF, Castro Laria L, Maldonado Pérez B, Perea Amarillo R and Argüelles-Arias F have received financial support to attend scientific meetings from Kern Pharma. Benítez Roldán A, Merino Bohórquez V, Calleja MA, Caunedo Álvarez Á, and Vilches Arenas Á do not have conflict of interest.

**Data sharing statement:** No additional data are available.

**STROBE statement:** the guidelines of the STROBE Statement have been adopted

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Corresponding author to:** Federico Argüelles-Arias, PhD, Doctor, Professor, Department of Gastroenterology, University Hospital Virgen Macarena, Dr. Fedriani 3, Seville 41007, Spain. [farguelles@telefonica.net](mailto:farguelles@telefonica.net)  
Telephone: +34-61-7348501

**Received:** August 29, 2018

**Peer-review started:** August 29, 2018

**First decision:** October 14, 2018

**Revised:** November 18, 2018

**Accepted:** December 7, 2018

**Article in press:** December 7, 2018

**Published online:** December 14, 2018



## Abstract

### BACKGROUND

Infliximab original has changed the natural history of inflammatory bowel diseases (IBD) over the past two decades. However, the recent expiration of its patent has allowed the entry of the first Infliximab biosimilar into the European and Spanish markets. Currently switching drugs data in IBD are limited.

### AIM

To compare the efficacy of infliximab biosimilar, CT-P13, against infliximab original, analyzing the loss of response of both at the 12 mo follow-up in patients with IBD.

### METHODS

An observational study of two cohorts has been conducted. One retrospective cohort that included patients with IBD treated with Infliximab original, and a prospective cohort of patients who were switching from infliximab original to infliximab biosimilar (CT-P13). We had analyzed the overall efficacy and loss of efficacy in patients in remission at the end of one year after treatment with the original drug compared to the results of the year of treatment with the biosimilar.

### RESULTS

98 patients (CD 67, CU 31) were included in both cohorts. The overall efficacy for infliximab original per year of treatment was 71% *vs* 68.2% for infliximab biosimilar ( $P = 0.80$ ). The loss of overall efficacy at 12 mo for infliximab original was 6.6% *vs* 14.5% for infliximab biosimilar ( $P = 0.806$ ). The loss of efficacy in patients who were in basal remission was 16.3% for infliximab original *vs* 27.1% for infliximab biosimilar. Adverse events were 9.2% for infliximab original *vs* 11.2% for infliximab biosimilar.

### CONCLUSION

The overall efficacy and loss of treatment response with infliximab biosimilar (CT-P13) is similar to that observed with infliximab original in patients who were switching at the 12 mo follow-up. There is no difference in the rate of adverse events.

**Key words:** Crohn's disease; Ulcerative colitis; CT-P13; Inflammatory bowel disease; Biosimilar agent; Infliximab original; Efficacy

© The Author(s) 2018. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Although not strictly necessary, there are few studies comparing efficacy and safety of the switch from infliximab original (Remicade®) to infliximab biosimilar CT-P13 *vs* the maintenance of the original infliximab. For this reason, we presented a comparative study with the original infliximab. Our observational study demonstrates the real-life clinical results of efficacy and safety of infliximab original and the efficacy and safety after switching from original infliximab to infliximab biosimilar CT-P13 at the 12 mo follow-up. Our results demonstrate there is no

statistical difference in remission rate, secondary loss of response or adverse events between both therapies.

Guerra Veloz MF, Argüelles-Arias F, Castro Laria L, Maldonado Pérez B, Benítez Roldán A, Perea Amarillo R, Merino Bohórquez V, Calleja MA, Caunedo Álvarez Á, Vilches Arenas Á. Loss of efficacy and safety of the switch from infliximab original to infliximab biosimilar (CT-P13) in patients with inflammatory bowel disease. *World J Gastroenterol* 2018; 24(46): 5288-5296  
URL: <https://www.wjgnet.com/1007-9327/full/v24/i46/5288.htm>  
DOI: <https://dx.doi.org/10.3748/wjg.v24.i46.5288>

## INTRODUCTION

Inflammatory bowel disease (IBD), which includes Crohn's disease (CD) and ulcerative colitis (UC), is a chronic gastrointestinal disorder of unknown etiology which causes deterioration in the quality of life of the patient<sup>[1]</sup>. The introduction of biological therapies two decades ago, particularly anti-tumor necrosis factor (TNF)-alpha drugs<sup>[2,3]</sup>, has revolutionized the therapeutic approach of IBD, especially in those patients with severe or refractory disease<sup>[2]</sup>. Despite the undoubted efficacy of biological therapy, these biological agents are much more expensive than traditional treatments, and so impose a considerable burden on the national healthcare system<sup>[4]</sup>. However, many biological products have reached or are close to patent expiration. This has led to the development of biosimilar drugs<sup>[5-7]</sup>.

CT-P13 (Remsima® and Inflectra®) was the first biosimilar of infliximab approved by the European Medicines Agency<sup>[8]</sup> in September 2013 and by the United States Food and Drug Administration (United States FDA)<sup>[9]</sup> in April 2016 for all indications of the originator product including IBD. The extrapolation of use of biosimilar infliximab was based on two phase III clinical trials in patients with rheumatoid arthritis (RA) (PLANETRA)<sup>[10]</sup> and ankylosing spondylitis (AS) (PLANETAS)<sup>[11]</sup>. Both studies at 30 wk proved the similarity of infliximab biosimilar against the reference product (Remicade®) in terms of efficacy, safety and immunogenicity. Furthermore, the results of the 102 wk open label extensions of PLANETAS<sup>[11]</sup> and PLANETRA<sup>[10]</sup> trials have been recently published, providing important information about the safety, efficacy and immunogenicity of switching from infliximab original to infliximab biosimilar in both AS and RA patients<sup>[12,13]</sup>.

During these years, some observational studies and real-life cohorts have been published on the efficacy and safety of the infliximab biosimilar in IBD<sup>[14-25]</sup>. Although not strictly necessary, there are few studies comparing efficacy and safety of the switch from infliximab original (Remicade®) to infliximab biosimilar CT-P13 *vs* the maintenance of the infliximab original<sup>[26,27]</sup>. One of the few studies that provide relevant evidence is the results of the phase IV, double-blind, parallel-group NOR-SWITCH

study<sup>[27]</sup>. This trial proved that switching from infliximab RP to CT-P13 was not inferior to continued treatment with infliximab RP. However, this study has received much criticism because of its methodological limitations and is not powered to perform a subgroup analysis, especially IBD patients<sup>[28]</sup>.

Despite initial concerns about infliximab biosimilar not having enough data about safety, security and immunogenicity, guidance from inflammatory bowel disease societies has gradually moved towards a positive and confident position on CT-P13. The European Crohn's Colitis Organization (ECCO) published its position statement on the use of biosimilars for IBD in December 2016 which states that "when a biosimilar product is registered in the European Union, it is considered to be as efficacious as the reference product when used in accordance with the information provided in the Summary of Product Characteristics"<sup>[29]</sup>.

For all the above, this study is designed to compare the efficacy of infliximab biosimilar CT-P13 against infliximab original, analyzing the loss of response of both at the 12 mo follow-up in patients with IBD.

## MATERIALS AND METHODS

### Study design

This was an observational study of two cohorts conducted at the Virgen Macarena Hospital (Seville, Spain). One retrospective cohort that included patients with IBD treated with infliximab original from January 2014 to December 2014, and one prospective cohort with patients who were switching from infliximab original to infliximab biosimilar (CT-P13) from March 2015 to February 2016. We had analyzed the overall efficacy and loss of efficacy in patients in remission at the end of one year after treatment with the original product compared to the results of the year of treatment with infliximab biosimilar. The study was approved by the Research Ethics Committee of the Virgen Macarena Hospital. Good clinical practice guidelines were followed and written informed consent was obtained from all patients.

### Patients

In the retrospective cohort we included all the patients with CD or UC who were treated with infliximab original at least once in 2014. In the prospective cohort we included all the patients with CD or UC who had been switching from infliximab original to infliximab biosimilar and had completed a 12 mo follow-up. Montreal classification status was recorded in all patients before enrolment.

### Study endpoints and assessments

The efficacy endpoint was the change in clinical remission in patients with infliximab original, and in patients switched from infliximab RP to CT-P13 assessed at 12 mo.

In patients with CD and UC remission was considered when: (1) Harvey-Bradshaw score (HB)  $\leq 4$  in patients

with CD or partial Mayo score  $\leq 2$  in patients with UC; (2) C-reactive protein (CRP)  $\leq 5$  mg/dL; (3) no use of steroids; (4) no surgery related to the disease activity; and (5) no increased dosage at the established follow-up time.

Adverse events (AE) were monitored from the first infusion of CT-P13 until the end of the study and they were recorded according to the Office of Human Research Protection.

### Statistical analysis

Demographic and nominal results were reported in percentages and frequencies. Numerical results were reported as average and standard deviation in cases of normal distribution and as median and interquartile range (IQR) in cases of asymmetrical distribution. The Wilcoxon test was used to compare the clinical scores (HB Score and partial Mayo Score) and the CRP values of patients. McNemar's test and the Cochran Q test were used to compare both groups' remission. 95% confidence intervals (CIs) were calculated and  $\alpha$  was set at 0.05 for the determination of statistical significance. Analysis was performed using SPSS 23 (IBM Corporation).

## RESULTS

A total of 98 patients were included in each cohort. 68.4% ( $n = 67$ ) were CD, and 31.6% ( $n = 31$ ) were UC.

### Retrospective cohort

The median age of patients was 39.9 (standard deviation SD 12.5 years old), (95%CI: 37.4-42.4). Over half of the patient population (55.4%) were men and 70.4% were non-smokers. Median time of the disease before starting the follow-up was 44 [Interquartile range (IQR) = 18; 100 mo]. Median duration of ongoing infliximab original treatment at the start of the study was 55 (IQR = 28.7; 72 mo). 40.8% used concomitant thiopurines before starting the follow-up. The baseline demographics and phenotypic characteristics of patients with CD and UC according to the Montreal classification are shown in Table 1.

At the start of the study the global remission was 77.6% (76/98) (95%CI: 68.8-86.3) and 71% (66/93) (95%CI: 61.2-80.7) at the 12 mo follow-up. In total, 76 patients were in remission at the time of switch, 83.7% (62/74) (95%CI: 74.7-92.9) had remained in remission at 12 mo of follow-up.

Five patients discontinued treatment (two endoscopic mucosa healing, three adverse effects including development of cervical neoplasia).

### CD

In the CD patient group, the median age was 37 (IQR = 30; 47 years). The median time of treatment with infliximab original before starting the follow-up was 59 (IQR = 30; 78 mo). The locations at diagnostic were colonic in 37.3%, with non-stricturing, non-penetrating

**Table 1** Comparative baseline demographics and phenotypic characteristics of patients with Crohn's disease and ulcerative colitis in both groups *n* (%)

		Prospective cohort	95%CI	Retrospective cohort	95%CI	P value
Characteristics (n = 98)						
Sex	men	51 (52)	41.6-62.4	56 (57.1)	46.8-67.4	0.280
	women	47 (48)	37.6-58.4	42 (42.9)	32.5-53.2	
Smoking status	Never	67 (68.3)	58.7-78.1	69 (70.4)	60.9-79.9	0.929
	Previous	18 (18.4)	10.2-26.5	16 (16.3)	8.5-24.1	
	Current	13 (13.3)	6.0-20.5	13 (13.3)	6.1-20.5	
Crohn's disease (n = 67)						
Age at diagnosis	A1 (< 17)	7 (10.5)	2.4-18.5	10 (15)	5.6-24.2	0.691
	A2 (17-40)	49 (73.1)	61.8-84.5	48 (71.6)	60.1; 83.2	
	A3 (> 40)	11 (16.4)	6.8-26.0	9 (13.4)	4.5; 22.3	
Location at diagnosis	L1 (ileal)	18 (26.9)	15.5-38.2	15 (22.4)	11.7-33.1	0.887
	L2 (colonic)	26 (38.8)	26.4-51.2	25 (37.3)	25.0-49.6	
	L3 (ileocolonic)	21 (31.3)	19.5-43.2	25 (37.3)	25.0-49.6	
	L4 (upper gastrointestinal tract)	2 (3.0)	0.3-10.4	2 (3.0)	0.4-10.4	
Disease behavior	B1 (non-stricturing, non-penetrating)	38 (56.7)	44.169.3	39 (58.2)	45.7-70.8	0.860
	B2 (stricturing)	14 (20.9)	10.4-31.4	13 (19.4)	9.2-29.6	
	B3 (penetrating)	15 (22.4)	11.7-33.1	15 (22.4)	11.7-33.1	
Perianal disease	Yes	37 (55.2)	42.6-67.9	39 (58.2)	45.7-70.8	0.727
Extraintestinal manifestations	No	42 (62.7)	50.4-75.0	46 (68.7)	56.8-80.5	0.466
Ulcerative colitis (n = 31)						
Extension	E1 (proctitis)	13 (41.9)	22.9-60.9	12 (38.7)	19.9-57.5	0.957
	E2 (left colitis)	10 (32.3)	14.2-50.3	11 (35.5)	17.0-53.9	
	E3 (pancolitis)	8 (25.8)	8.8-42.9	8 (25.8)	8.8-42.9	
	S1 (mild)	11 (35.5)	17.0-53.9	10 (32.3)	14.2-50.3	
Severity	S2 (moderate)	16 (51.6)	32.4-70.8	17 (54.8)	35.7-74.0	0.962
	S3 (severe)	4 (12.9)	3.6-29.8	4 (12.9)	3.6-29.8	
	Extraintestinal manifestations	No	26 (83.9)	66.3-94.5	27 (87.1)	
Prior treatment		75 (76.5)	67.6-85.4	85 (86.7)	79.5-94	0.097
Thiopurines		25 (25.5)	16.4-34.7	25 (25.5)	16.4-34.7	0.970
Methotrexate						
Concomitant treatment		50 (51)	40.6-61.4	40 (40.8)	30.6-51.1	0.197
Thiopurines		8 (8)	2.2-14.1	7 (7,1)	1.1-10,1	0.999
Methotrexate						

behavior in 58.2%. 46.3% used concomitant thiopurines before starting the follow-up.

Of the 67 patients analyzed, the basal remission was 76.1% (51/67) (95%CI: 65.2-87.1) and 69.2% (45/65) (95%CI: 57.2-81.2) at the 12 mo follow-up. Of the 51 patients who were in initial remission this was maintained in 82.3% (42/51) (95%CI: 70.9-93.8) of patients at the 12 mo follow-up. Two patients discontinued treatment: Two adverse effects including development of a cervix neoplasia.

The HB score showed significant changes over the 12 mo period [median HB score 95%CI: 1 (0-2) vs 1 (1-3) at month 0 and 12,  $P = 0.005$ ]. No significant changes were observed in median of CRP levels [1.0 (0-5) vs 0.4 (0.2-2) at 0 and 12 mo,  $P = 0.464$ ].

### Ulcerative colitis

In the UC patient, the median time of treatment with infliximab original before starting the follow-up was 42 (IQR = 20; 60 mo). The locations at diagnostic were proctitis in 38.7% with a moderate severity in 54.8%.

Of the 31 patients analyzed, the basal remission was 80.6% (25/31) (95%CI: 62.5-92.5), and 75% (21/28) (95%CI: 57.2-92.8) at the 12 mo follow-up. Of the 25

patients who were in initial remission this was maintained in 87% (20/23) (95%CI: 70.9-93.8) of patients at the 12 mo follow-up. Three patients discontinued treatment: (two endoscopic mucosa healing and one adverse effect).

No significant changes in the median (95%CI) partial Mayo score were observed in switched patients over the 12 mo [2 (1-3); vs 1 (1-3) at 0;  $P = 0.067$ ]. No significant changes in the median CRP level (95%CI) were observed over the same period [2 (1-10) vs 0.6 (0.2-5.5) at months 0 and 12,  $P = 0.654$ ].

### Prospective cohort

The median age of patients was 41 (SD 12.7 years). Over half of the patient population (56.1%) were men and 68.3% were non-smokers. Median time of the disease before starting the follow-up was 100 (IQR = 77; 151 mo). Median duration of ongoing infliximab original treatment at the start of the study was 60.7 (IQR = 10.5; 73.5 mo). 51% used concomitant thiopurines before starting the follow-up. The baseline demographics and phenotypic characteristics of patients with CD and UC according to the Montreal classification are shown in Table 1.

Of the 98 patients analyzed, 82.7% (81/98) (95%CI:



**Table 2** Comparison of efficacy and loss of efficacy in both groups

	Infliximab original Group	95%CI	CT-P13 Group	95%CI	P value	Rate difference (95%CI)
Basal global remission	76/98 (77.6%)	66.8-86.3	81/98 (82.7%)	74.6-90.7	0.474	-0.173-0.071
Global remission 12 mo	66/93 (71%)	61.2-80.7	60/88 (68.2%)	51.1-71.4	0.806	-0.117-0.173
Basal remission CD	51/67 (76.1%)	65.2-87.1	56/67 (83.6%)	73.9-93.2	0.389	-0.225-0.076
Remission 12 mo CD	45/65 (69.2%)	57.2-81.2	42/62 (67.7%)	55.3-80.2	0.992	-0.163-0.192
Basal UC remission	25/31 (80.6%)	62.5-92.5	25/31 (80.6%)	62.5-92.5	0.748	-0.29-0.229
Remission 12 mo UC	21/28 (75%)	57.2-92.8	18/26 (69.2%)	49.6-88.9	0.866	-0.219-0.334

UC: Ulcerative colitis. CD: Crohn's disease.

74.6-90.7) were in remission at the beginning of the study, and 68.2% (60/88) (95%CI: 57.9-78.5) at the 12 mo follow-up. Of the 81 patients who were in initial remission this was maintained in 72.9% (54/74) (95%CI: 62.2-83.8) of patients at 12 mo.

Ten patients discontinued treatment (3 endoscopic mucosal remission, 6 adverse effects, 1 patient did not attend follow-up visits).

### Crohn's disease

In the CD patient, median time of treatment with infliximab original before starting the follow-up was 74.3 (IQR = 41; 108 mo). The location at diagnostic was colonic in 38.3%, with non-stricturing, non-penetrating behavior in 56.7%, and 55% had perineal disease. 56.7% used concomitant thiopurines before starting the follow-up.

Of the 67 patients analyzed, the basal remission was 83.6% (56/67) (95%CI: 87.3-93.2) and 67.7% (42/62) (95%CI: 55.3-80.2) at the 12 mo follow-up. Of the 56 patients who were in initial remission this was maintained in 69.8% (37/53) (95%CI: 56.5-83.1) of patients at the 12 mo follow-up ( $P = 0.634$ ). 92.5% (62/67) patients completed the follow-up.

Five patients discontinued treatment: Four adverse effects, one patient did not attend follow-up visits.

The HB score showed significant changes over the 12 mo period [median HB score 95%CI: 1 (1-2) vs 1 (1-3) at months 0 and 12,  $P = 0.007$ ]. No significant changes were observed in median of CRP levels [1.0 (0-6) vs 0.36 (0.2-2) at 0 and 12 mo ( $P = 0.364$ )].

### Ulcerative colitis

In the UC patient, the median time of treatment with infliximab original before starting the follow-up was 52 (IQR = 20; 60 mo). The location at diagnostic was left in 41.9% with a moderate severity in 51.6%.

Of the 31 patients analyzed, the basal remission was 80.6% (25/31) (95%CI: 62.5-92.5), and 69.2% (18/26) (95%CI: 49.6-88.9) at the 12 mo follow-up. Of the 25 patients who were in initial remission this was maintained in 81% (17/21) (95%CI: 62.5-92.5) of patients at the 12 mo follow-up. Five patients discontinued treatment: Three endoscopic mucosa healing and two adverse effects.

No significant changes in the median (95%CI) partial Mayo score were observed in switched patients over the

12 mo [2 (1-3) vs 1 (0-3) at month 0 and 12,  $P = 0.058$ ]. No significant changes in the median CRP level (95%CI) were observed over the same period [2 (1-10) vs 0.6 (0.2-1.4) at month 0 and 12,  $P = 0.329$ ].

### Comparison analysis

The basal remission rate of the infliximab original group was 77.6% vs 82.7% of infliximab biosimilar ( $P = 0.474$ ). At 12 mo the remission rate was 71% in infliximab original vs 68.2% of biosimilar infliximab ( $P = 0.806$ ) without achieving statistical significance. This is showed in Table 2.

The loss of overall efficacy at 12 mo in the infliximab original group was 6.6% and 14.5% in the infliximab biosimilar group, without achieving statistical significance ( $P = 0.835$ ). There were not statistically significant differences observed when comparing the loss of efficacy by type of disease, CD ( $P = 0.992$ ) and UC ( $P = 0.866$ , Figure 1).

When we analyzed patients, who were in basal remission, the loss of efficacy was 16.3% in the infliximab original vs 27.1% in the infliximab biosimilar at the 12 mo follow-up, without statistical differences ( $P = 0.162$ ). There were also no statistically significant differences when we analyzed by type of disease CD ( $P = 0.205$ ) and UC ( $P = 0.890$ , Table 3).

No significant changes in disease activity, measured by HB in CD ( $P = 0.385$ ) and Partial Mayo score in UC ( $P = 0.349$ ) were observed between both groups. Significant changes in the median CRP level (95%CI) were observed between both groups ( $P = 0.014$ ), nevertheless, it was always in remission ( $\leq 5$  mg/dL).

### Safety


Retrospective cohort: 9 AEs occurred in 9/98 (9.2%) patients: One hypertensive crisis, three infusion reactions, one palpitations, one asthenia, one arthralgia, one HBV reactivation, one cervical carcinoma. Four patients discontinued treatment because of AEs. Severe adverse events were considered in a patient with UC for HBV reactivation that needed treatment and one CD patient who developed a cervical carcinoma.

Prospective cohort: 11 AEs occurred in 11/98 (11.2%) patients: One skin reaction, one case of abdominal pain, two cases of headache and two of paresthesia during infusion, one case of Sweet's syndrome, two of polyarthralgia, and two of palpitations. Six patients dis-

**Table 3** Comparative of maintenance of remission at 12 mo in patients with initial remission in both groups

	Infliximab original group	95%CI	CT-P13 group	95%CI	P value	Rate Difference (95%CI)
Maintained basal remission at 12 mo	62/74 (83.7%)	74.7-92.9	54/74 (72.9%)	62.2-83.5	0.162	-0.037-0.253
Maintained basal remission at 12 mo CD	42/51 (82.3%)	70.9-93.8	37/53 (69.8%)	56.5-83.1	0.205	-0.056-0.307
Maintained basal remission at 12 mo UC	20/23 (87%)	66.4-97.2	17/21 (81.0%)	62.5-92.5	0.890	-0.203-0.23

CD: Crohn's disease; UC: Ulcerative colitis.

**Figure 1** Loss of efficacy at 12 mo in both groups.

continued treatment because of AEs. Severe adverse event was considered in one patient with CD who had Sweet's syndrome needing hospitalization and discontinued treatment. In addition, one patient with UC discontinued because of paresthesia during the infusion.

There were not statistical differences between each group ( $P = 0.814$ , 95%CI: 0.115-0.075)

## DISCUSSION

Despite initial concerns about infliximab biosimilar, different articles have proved the efficacy and safety of CT-P13. Also, the European Crohn's Colitis Organization (ECCO) published its favorable position statement on the use of biosimilars for IBD in December 2016<sup>[29]</sup>. We recently have published three studies in this matter<sup>[15,24-25]</sup>. In these studies, we proved the efficacy of CT-P13, although our patients' cohorts (switch cohort) have not been compared to a non-switch cohort. Many studies and practical clinical trials have proved the efficacy of anti-TNF- $\alpha$  in IBD<sup>[2-4]</sup>. However, up to 30% of patients show no clinical benefit after induction therapy (primary non-responders), and another 30%-40% lose response during the first year of treatment (secondary non-responders)<sup>[30]</sup>, with an estimated annual risk of loss of efficacy between 13%-15% per patient/year<sup>[31]</sup>. Consequently, we wanted to explore whether our loss of efficacy in the switch cohort was similar to a non-switch cohort in our centre.

Our observational study proves the real-life clinical results of efficacy and safety of infliximab original and the efficacy and safety after switching from infliximab original to infliximab biosimilar CT-P13 at 12 mo follow-up. Our results prove there is no statistical difference in remission rate, secondary loss of response or adverse events between both therapies. Both groups had similar baseline characteristics. The basal remission in the infliximab original group was 77.6% vs 82.7% for infliximab biosimilar. At 12 mo, the infliximab original remission rate was 71% vs 68.2% for the infliximab biosimilar group. The loss of overall efficacy at 12 mo in the infliximab original group was 6.6% and 14.5% in the infliximab biosimilar one ( $P = 0.806$ ). The loss of efficacy in patients who were in basal remission was 16.3% in the infliximab original vs 27.1% in the infliximab biosimilar. It is important to note that, although the difference in the loss of efficacy between the two groups (10.8%) is high, no statistical differences were observed. We believe that this difference may be caused by the fact that an insufficient number of patients were included in the study and that the intervals of confidence were so high that it was impossible to obtain a statistical difference.

In other hand, nocebo is a concept that should be taken into account in our results. Many patients did not have enough confidence in the biosimilar, and as it has been proved in some studies<sup>[32,33]</sup>, nocebo can play a role in the efficacy of the switch cohort. Another important point to analyze is that, although present in both groups,

the CD patients' median of HB score showed significant changes over the 12 mo, but it never went over the definition of remission ( $\leq 4$ ). In both groups, in UC patients the median partial Mayo score remained without changes, CRP levels remained without clinically relevant changes in both groups ( $\leq 5$ ). We did not find changes in the clinical and analytical parameters, and so we believe that the strong nocebo effect in the biosimilar group explains this loss of efficacy.

In the NOR-SWITCH trial<sup>[27]</sup> the proportion of patients with disease worsening after six months of stable disease was 26.2% in the infliximab original arm and 29.6% in the CT-P13 arm, after 52 wk. The worsening of the disease happened more frequently in CD patients, however the authors concluded that switching from infliximab original to infliximab biosimilar (CT-P13) was not inferior to continued treatment with infliximab original regarding efficacy, safety and immunogenicity. Also, in the 26 wk open label NOR-SWITCH EXTENSION trial reported in the ECCO congress last February<sup>[34]</sup>, safety and immunogenicity were assessed regarding CT-P13 treatment throughout the 78 wk study period (maintenance group) compared to switching from infliximab original to CT-P13 at week 52 (switch group). Overall disease worsening occurred in 16.8% of patients in the maintenance group and in 11.6% in the switch group. In CD, disease worsening occurred in 20.6% and 13.1% and in UC in 15.4% and 2.9% of patients in the maintenance and switch group, respectively. These results were within the non-inferiority margin (15%).

These results are similar in other studies. In the SECURE study<sup>[35]</sup> infliximab serum concentrations in adults with ulcerative colitis ( $n = 59$ ) and Crohn's disease ( $n = 61$ ) were examined 16 wk after switching from infliximab original to CT-P13. Serum infliximab concentration with CT-P13 was not inferior to those with infliximab original. Furthermore, no significant changes were noted in clinical and biochemical variables, quality of life and tolerability<sup>[35]</sup>. In the PROSIT-BIO study<sup>[36]</sup> the efficacy estimations were 95.7%, 86.4%, and 73.7% for naive; 97.2%, 85.2%, and 62.2% for pre-exposed; and 94.5%, 90.8%, and 78.9% for switch at 8 wk, 16 wk, and 24 wk respectively. 12.1% of adverse events were reported, 38 (6.9%) of them being infusion-related reactions. In a prospective observational single center study<sup>[26]</sup>, that included 191 patients who were switched from original infliximab to CT-P13 and 19 patients who continued with the original. They showed no statistical difference in remission (58.1% vs 47.4%,  $P = 0.37$ ), response (12.6% vs 10.5%,  $P = 0.80$ ), secondary loss of response (24.6% vs 42.1%,  $P = 0.10$ ), or adverse events (4.7% vs 0,  $P = 1.0$ ) between those who switched to CT-P13 and those who continued infliximab original at the 12 mo follow-up<sup>[26]</sup>.

All data is in accordance with our results. All observational post-marketing studies published to date have reported positive outcomes for efficacy measures in patients with CD and UC treated with CT-P13, irrespective of prior anti-TNF- $\alpha$  treatment<sup>[14-26]</sup>

Finally, the recently published systematic review with meta-analysis<sup>[37]</sup>, in patients with CD switching from infliximab RP to CT-P13 demonstrated high rates of sustained clinical response at 30-32 wk (0.85, 95%CI: 0.71-0.93) and 48-63 wk (0.75, 95%CI: 0.44-0.92) and high rates for sustained clinical remission 0.74 (95%CI: 0.55-0.87) and 0.92 (95%CI: 0.38-0.99), at 16 wk and 51 wk respectively.

In our study, 9.2% (9/98) of patients had drug-related adverse events with infliximab original, four patients discontinued the treatments for this reason. In the infliximab biosimilar 11.2% (11/98) of patients had drug-related adverse events, six patients discontinued treatment because of AEs. These data are related to the results of several studies on the rate of adverse events between 10%-20%.

Our study has some limitations. First of all, and perhaps most importantly, we analyzed a retrospective cohort of patients with infliximab original, which could be a methodological deficit with an error of data interpretation. Similarly, we could not measure calprotectin, drug trough levels, or the presence of antidrug antibodies as has been done in other studies, because at the time of the study these were not available in our hospital. Therefore, it has not been possible to ascertain the cause of the loss of response observed in some patients. We were also unable to measure mucosal healing or perform an endoscopy on any of the patients at the time of the study.

Despite the limitations in our study, we believe that it shows real data in clinical practice in the one-year follow-up. The loss of efficacy in patients in clinical remission treated with biosimilars was 10.8%, close to the margin of non-inferiority of 15% as other previously published studies<sup>[25-32]</sup>. We conclude that the overall efficacy and loss of treatment response with Infliximab biosimilar (CT-P13) is similar to that observed with Infliximab original in patients who were switching at the 12 mo follow-up.

## ARTICLE HIGHLIGHTS

### Research background

Infliximab was the first monoclonal antibody against tumor necrosis factor alpha approved by the US Food and Drug Administration (United States FDA) and European Medicines Agency (EMA) in the treatment of inflammatory bowel disease (IBD). Despite the undoubted efficacy of this biological therapy, this biological agent is much more expensive than traditional treatments and, therefore, imposes a considerable burden on the national health system. However, with the recent expiration of its patent, biosimilar medicines have been developed. CT-P13 (Remsuma® and Inflectra®) was the first biosimilar of infliximab approved by EMA and United States FDA for all indications of the originator product including IBD. The effectiveness in IBD is being debated due to the approval of this biosimilar was based on clinical trials in rheumatological diseases.

### Research motivation

Some observational studies and real-life cohorts have been published on the efficacy and safety of the infliximab biosimilar in naive patients with IBD. Although not strictly necessary, there are few studies comparing efficacy and safety of the switch from infliximab original to infliximab biosimilar CT-P13 vs

the maintenance of the infliximab original.

### Research objectives

To compare the efficacy of infliximab biosimilar, against infliximab original, analyzing the loss of response of both at the 12 mo follow-up in patients with IBD

### Research methods

An observational study of two cohorts has been conducted. One retrospective cohort that included patients with IBD treated with Infliximab original, and a prospective cohort of patients who were switching from infliximab original to infliximab biosimilar. 98 patients were included in each cohort. We had analyzed the overall efficacy and loss of efficacy in patients in remission at the end of one year after treatment with the original drug compared to the results of the year of treatment with the biosimilar. The efficacy was reported based clinical scores [Harvey-Bradshaw score (HB)  $\leq 4$  in patients with Crohn's disease (CD) or partial Mayo score  $\leq 2$  in patients with ulcerative colitis (UC)] and biochemical test. Wilcoxon test, McNemar's test and the Cochran Q test were used to compare the variables in both periods.

### Research results

This study demonstrated the overall efficacy and the loss of overall efficacy for infliximab original and infliximab biosimilar were similar per year of treatment.

The loss of efficacy in patients who were in basal remission was 16.3% in the infliximab original vs 27.1% in the infliximab biosimilar. This 10.8% of difference in the loss of efficacy between the two groups did not show a statistical difference. No significant changes in disease activity, measured by HB in CD ( $P = 0.385$ ) and Partial Mayo score in UC ( $P = 0.349$ ) were observed between both groups. Significant changes in the median CRP level (95%CI) were observed between both groups ( $P = 0.014$ ), nevertheless, it was always in remission ( $\leq 5$  mg/dL). Adverse events were similar in both cohorts.

### Research conclusions

Our study suggests that the overall efficacy and loss of treatment response with infliximab biosimilar (CT-P13) is similar to that observed with infliximab original in patients who were switching at the 12 mo follow-up. The high difference in the loss of efficacy shown in patients who were in basal remission with infliximab original vs infliximab biosimilar could be explained by the following: in our data we found the intervals of confidence in both groups were so high that it was impossible to obtain a statistical difference, and in other hand, placebo is a concept that should be taken into account in our results. Many patients did not have enough confidence in the biosimilar the first year.

### Research perspectives

Future prospective comparative studies with long-term follow-up are necessary to discriminate the loss of efficacy in patient who were switched to infliximab biosimilar.

## ACKNOWLEDGMENTS

The authors would like to thank the Andalusian Group of Inflammatory Bowel Disease and Kern Pharma Biologics.

## REFERENCES

- 1 **Feuerstein JD**, Cheifetz AS. Crohn Disease: Epidemiology, Diagnosis, and Management. *Mayo Clin Proc* 2017; **92**: 1088-1103 [PMID: 28601423 DOI: 10.1016/j.mayocp.2017.04.010]
- 2 **Côté-Daigneault J**, Bouin M, Lahaie R, Colombel JF, Poitras P. Biologics in inflammatory bowel disease: what are the data? *United European Gastroenterol J* 2015; **3**: 419-428 [PMID: 26535119 DOI: 10.1177/2050640615590302]
- 3 **Tracey D**, Klareskog L, Sasso EH, Salfeld JG, Tak PP. Tumor necrosis factor antagonist mechanisms of action: a comprehensive review. *Pharmacol Ther* 2008; **117**: 244-279 [PMID: 18155297 DOI: 10.1016/j.pharmthera.2007.10.001]
- 4 **Bodger K**, Kikuchi T, Hughes D. Cost-effectiveness of biological therapy for Crohn's disease: Markov cohort analyses incorporating United Kingdom patient-level cost data. *Aliment Pharmacol Ther* 2009; **30**: 265-274 [PMID: 19438428 DOI: 10.1111/j.1365-2036.2009.04033.x]
- 5 European Medicines Agency- New guide on biosimilar medicines for healthcare professionals. Available from: URL: [http://www.ema.europa.eu/ema/index.jsp?curl=pages/news\\_and\\_events/news/2017/05/news\\_detail\\_002739.jsp&mid=WC0b01ac058004d5c1](http://www.ema.europa.eu/ema/index.jsp?curl=pages/news_and_events/news/2017/05/news_detail_002739.jsp&mid=WC0b01ac058004d5c1)
- 6 **European Medicines Agency**. Procedural advice for users of the centralized procedure for similar biological medicinal products applications. London, UK: European Medicines Agency; 2011
- 7 **European Medicines Agency**. Guideline on similar biological medicinal products. London 2005
- 8 **European Medicines Agency**. Committee for Medicinal Products for Human Use (CHMP). Assessment report: Remsima (infliximab). 2013. Available from: URL: [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/EPAR\\_-\\_Public\\_assessment\\_report/human/002576/WC500151486.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Public_assessment_report/human/002576/WC500151486.pdf)
- 9 **INFLECTRA prescribing information**. Available from: URL: [http://www.accessdata.fda.gov/drugsatfda\\_docs/label/2016/125544s0001bl.pdf](http://www.accessdata.fda.gov/drugsatfda_docs/label/2016/125544s0001bl.pdf)
- 10 **Yoo DH**, Hrycaj P, Miranda P, Ramitterre E, Piotrowski M, Shevchuk S, Kovalenko V, Prodanovic N, Abello-Bañfi M, Gutierrez-Ureña S, Morales-Olazabal L, Tee M, Jimenez R, Zamani O, Lee SJ, Kim H, Park W, Müller-Ladner U. A randomised, double-blind, parallel-group study to demonstrate equivalence in efficacy and safety of CT-P13 compared with innovator infliximab when coadministered with methotrexate in patients with active rheumatoid arthritis: the PLANETRA study. *Ann Rheum Dis* 2013; **72**: 1613-1620 [PMID: 23687260 DOI: 10.1136/annrheumdis-2012-203090]
- 11 **Park W**, Hrycaj P, Jeka S, Kovalenko V, Lysenko G, Miranda P, Mikazane H, Gutierrez-Ureña S, Lim M, Lee YA, Lee SJ, Kim H, Yoo DH, Braun J. A randomised, double-blind, multicentre, parallel-group, prospective study comparing the pharmacokinetics, safety, and efficacy of CT-P13 and innovator infliximab in patients with ankylosing spondylitis: the PLANETAS study. *Ann Rheum Dis* 2013; **72**: 1605-1612 [PMID: 23687259 DOI: 10.1136/annrheumdis-2012-203091]
- 12 **Yoo DH**, Prodanovic N, Jaworski J, Miranda P, Ramitterre E, Lanzon A, Baranaukaite A, Wiland P, Abud-Mendoza C, Oparanov B, Smiyan S, Kim H, Lee SJ, Kim S, Park W. Efficacy and safety of CT-P13 (biosimilar infliximab) in patients with rheumatoid arthritis: comparison between switching from reference infliximab to CT-P13 and continuing CT-P13 in the PLANETRA extension study. *Ann Rheum Dis* 2017; **76**: 355-363 [PMID: 27130908 DOI: 10.1136/annrheumdis-2015-208786]
- 13 **Park W**, Yoo DH, Miranda P, Brzosko M, Wiland P, Gutierrez-Ureña S, Mikazane H, Lee YA, Smiyan S, Lim MJ, Kadinov V, Abud-Mendoza C, Kim H, Lee SJ, Bae Y, Kim S, Braun J. Efficacy and safety of switching from reference infliximab to CT-P13 compared with maintenance of CT-P13 in ankylosing spondylitis: 102-week data from the PLANETAS extension study. *Ann Rheum Dis* 2017; **76**: 346-354 [PMID: 27117698 DOI: 10.1136/annrheumdis-2015-208783]
- 14 **Jung YS**, Park DI, Kim YH, Lee JH, Seo PJ, Cheon JH, Kang HW, Kim JW. Efficacy and safety of CT-P13, a biosimilar of infliximab, in patients with inflammatory bowel disease: A retrospective multicenter study. *J Gastroenterol Hepatol* 2015; **30**: 1705-1712 [PMID: 25974251 DOI: 10.1111/jgh.12997]
- 15 **Argüelles-Arias F**, Guerra Veloz MF, Perea Amarillo R, Vilches-Arenas A, Castro Laria L, Maldonado Pérez B, Chaaro D, Benítez Roldán A, Merino V, Ramírez G, Caunedo Álvarez A, Romero Gómez M. Effectiveness and Safety of CT-P13 (Biosimilar Infliximab) in Patients with Inflammatory Bowel Disease in Real Life at 6 Months. *Dig Dis Sci* 2017; **62**: 1305-1312 [PMID: 28281165 DOI: 10.1007/s10620-017-4511-4]
- 16 **Gecse KB**, Lovász BD, Farkas K, Banai J, Bene L, Gasztonyi B, Golovics PA, Kristóf T, Lakatos L, Csontos AA, Juhász M, Nagy F, Palatka K, Papp M, Patai Á, Lakner L, Salamon Á, Szamosi T, Szepes Z, Tóth GT, Vincze Á, Szalay B, Molnár T, Lakatos PL. Efficacy and



- Safety of the Biosimilar Infliximab CT-P13 Treatment in Inflammatory Bowel Diseases: A Prospective, Multicentre, Nationwide Cohort. *J Crohns Colitis* 2016; **10**: 133-140 [PMID: 26661272 DOI: 10.1093/ecco-jcc/jjv220]
- 17 **Park SH**, Kim YH, Lee JH, Kwon HJ, Lee SH, Park DI, Kim HK, Cheon JH, Im JP, Kim YS, Lee SY, Lee SJ. Post-marketing study of biosimilar infliximab (CT-P13) to evaluate its safety and efficacy in Korea. *Expert Rev Gastroenterol Hepatol* 2015; **9** Suppl 1: 35-44 [PMID: 26395533 DOI: 10.1586/17474124.2015.1091309]
- 18 **Farkas K**, Rutka M, Golovics PA, Végh Z, Lovász BD, Nyári T, Gecse KB, Kolar M, Bortlik M, Duricova D, Machkova N, Hruha V, Lukas M, Mitrova K, Malickova K, Bálint A, Nagy F, Bor R, Milassin Á, Szepes Z, Palatka K, Lakatos PL, Lukas M, Molnár T. Efficacy of Infliximab Biosimilar CT-P13 Induction Therapy on Mucosal Healing in Ulcerative Colitis. *J Crohns Colitis* 2016; **10**: 1273-1278 [PMID: 27106537 DOI: 10.1093/ecco-jcc/jjw085]
- 19 **Jahnsen J**. Clinical experience with infliximab biosimilar Remsimab (CT-P13) in inflammatory bowel disease patients. *Therap Adv Gastroenterol* 2016; **9**: 322-329 [PMID: 27134662 DOI: 10.1177/1756283X16636764]
- 20 **Smits LJ**, Derikx LA, de Jong DJ, Boshuizen RS, van Esch AA, Drenth JP, Hoentjen F. Clinical Outcomes Following a Switch from Remicade® to the Biosimilar CT-P13 in Inflammatory Bowel Disease Patients: A Prospective Observational Cohort Study. *J Crohns Colitis* 2016; **10**: 1287-1293 [PMID: 27095751 DOI: 10.1093/ecco-jcc/jjw087]
- 21 **Sieczkowska J**, Jarzębicka D, Banaszkiewicz A, Plocek A, Gawronska A, Toporowska-Kowalska E, Oracz G, Meglicka M, Kierkus J. Switching Between Infliximab Originator and Biosimilar in Paediatric Patients with Inflammatory Bowel Disease. Preliminary Observations. *J Crohns Colitis* 2016; **10**: 127-132 [PMID: 26721942 DOI: 10.1093/ecco-jcc/jjv233]
- 22 **Smits LJ**, Grelack A, Derikx LAAP, de Jong DJ, van Esch AAJ, Boshuizen RS, Drenth JPH, Hoentjen F. Long-Term Clinical Outcomes After Switching from Remicade® to Biosimilar CT-P13 in Inflammatory Bowel Disease. *Dig Dis Sci* 2017; **62**: 3117-3122 [PMID: 28667429 DOI: 10.1007/s10620-017-4661-4]
- 23 **Gonczi L**, Gecse KB, Vegh Z, Kurti Z, Rutka M, Farkas K, Golovics PA, Lovasz BD, Banai J, Bene L, Gasztonyi B, Kristof T, Lakatos L, Miheller P, Nagy F, Palatka K, Papp M, Patai A, Salamon A, Szamosi T, Szepes Z, Toth GT, Vincze A, Szalay B, Molnar T, Lakatos PL. Long-term Efficacy, Safety, and Immunogenicity of Biosimilar Infliximab After One Year in a Prospective Nationwide Cohort. *Inflamm Bowel Dis* 2017; **23**: 1908-1915 [PMID: 28922253 DOI: 10.1097/MIB.0000000000001237]
- 24 **Argüelles-Arias F**, Guerra Veloz MF, Perea Amarillo R, Vilches-Arenas A, Castro Laria L, Maldonado Pérez B, Chaaro Benallal D, Benítez Roldán A, Merino V, Ramirez G, Calleja-Hernández MA, Caunedo Álvarez A, Romero Gómez M. Switching from reference infliximab to CT-P13 in patients with inflammatory bowel disease: 12 months results. *Eur J Gastroenterol Hepatol* 2017; **29**: 1290-1295 [PMID: 28902041 DOI: 10.1097/MEG.0000000000000953]
- 25 **Guerra Veloz MF**, Vázquez Morón JM, Belvis Jiménez M, Pallarés Manrique H, Valdés Delgado T, Castro Laria L, Maldonado Pérez B, Benítez Roldán A, Perea Amarillo R, Merino V, Caunedo Álvarez A, Argüelles Arias F. Switching from reference infliximab to CT-P13 in patients with inflammatory bowel disease: results of a multicenter study after 12 months. *Rev Esp Enferm Dig* 2018; **110**: 564-570 [PMID: 29893581 DOI: 10.17235/reed.2018.5368/2017]
- 26 **Ratnakumaran R**, To N, Gracie DJ, Selinger CP, O'Connor A, Clark T, Carey N, Leigh K, Bournier L, Ford AC, Hamlin PJ. Efficacy and tolerability of initiating, or switching to, infliximab biosimilar CT-P13 in inflammatory bowel disease (IBD): a large single-centre experience. *Scand J Gastroenterol* 2018; **53**: 700-707 [PMID: 29687730 DOI: 10.1080/00365521.2018.1464203]
- 27 **Jørgensen KK**, Olsen IC, Goll GL, Lorentzen M, Bolstad N, Haavardsholm EA, Lundin KEA, Mørk C, Jahnsen J, Kvien TK; NOR-SWITCH study group. Switching from originator infliximab to biosimilar CT-P13 compared with maintained treatment with originator infliximab (NOR-SWITCH): a 52-week, randomised, double-blind, non-inferiority trial. *Lancet* 2017; **389**: 2304-2316 [PMID: 28502609 DOI: 10.1016/S0140-6736(17)30068-5]
- 28 **Ribaldone DG**, Saracco GM, Astegiano M, Pellicano R. Efficacy of infliximab biosimilars in patients with Crohn's disease. *Lancet* 2017; **390**: 2435-2436 [PMID: 29208303 DOI: 10.1016/S0140-6736(17)33047-7]
- 29 **Danese S**, Fiorino G, Raine T, Ferrante M, Kemp K, Kierkus J, Lakatos PL, Mantzaris G, van der Woude J, Panes J, Peyrin-Biroulet L. ECCO Position Statement on the Use of Biosimilars for Inflammatory Bowel Disease-An Update. *J Crohns Colitis* 2017; **11**: 26-34 [PMID: 27927718 DOI: 10.1093/ecco-jcc/jjw198]
- 30 **Papamichael K**, Gils A, Rutgeerts P, Levesque BG, Vermeire S, Sandborn WJ, Vande Casteele N. Role for therapeutic drug monitoring during induction therapy with TNF antagonists in IBD: evolution in the definition and management of primary nonresponse. *Inflamm Bowel Dis* 2015; **21**: 182-197 [PMID: 25222660 DOI: 10.1097/MIB.0000000000000202]
- 31 **Gisbert JP**, Panés J. Loss of response and requirement of infliximab dose intensification in Crohn's disease: a review. *Am J Gastroenterol* 2009; **104**: 760-767 [PMID: 19174781 DOI: 10.1038/ajg.2008.88]
- 32 **Rezk MF**, Pieper B. To See or NOsee: The Debate on the Nocebo Effect and Optimizing the Use of Biosimilars. *Adv Ther* 2018; **35**: 749-753 [PMID: 29873005 DOI: 10.1007/s12325-018-0719-8]
- 33 **Faasse K**, Petrie KJ. The nocebo effect: patient expectations and medication side effects. *Postgrad Med J* 2013; **89**: 540-546 [PMID: 23842213 DOI: 10.1136/postgradmedj-2012-131730]
- 34 **Jørgensen KK**, Goll GL, Sexton I, Olsen IC. Long-term efficacy and safety of biosimilar infliximab (CT-P13) after switching from originator infliximab: Explorative subgroup analyses in IBD from the NOR-SWITCH EXTENSION trial. *J Crohns Colitis* 2018; **S348-S349** [DOI: 10.1093/ecco-jcc/jjx180.610]
- 35 **Strik AS**, van de Vrie W, Bloemsaat-Minekus JPI, Nurmohamed M, Bossuyt PJJ, Bodelier A, Rispens T, van Megen YJB, D'Haens GR; SECURE study group. Serum concentrations after switching from originator infliximab to the biosimilar CT-P13 in patients with quiescent inflammatory bowel disease (SECURE): an open-label, multicentre, phase 4 non-inferiority trial. *Lancet Gastroenterol Hepatol* 2018; **3**: 404-412 [PMID: 29606564 DOI: 10.1016/S2468-1253(18)30082-7]
- 36 **Fiorino G**, Manetti N, Armuzzi A, Orlando A, Variola A, Bonovas S, Bossa F, Maconi G, D'Incà R, Lionetti P, Cantoro L, Fries W, Annunziata ML, Costa F, Terpin MM, Biancone L, Cortezzi CC, Amato A, Ardizzone S, Danese S, Guidi L, Rizzuto G, Massella A, Andriulli A, Massari A, Lorenzon G, Ghione S, Kohn A, Ventra A, Annese V; PROSIT-BIO Cohort. The PROSIT-BIO Cohort: A Prospective Observational Study of Patients with Inflammatory Bowel Disease Treated with Infliximab Biosimilar. *Inflamm Bowel Dis* 2017; **23**: 233-243 [PMID: 28092307 DOI: 10.1097/MIB.0000000000000955]
- 37 **Komaki Y**, Yamada A, Komaki F, Micic D, Ido A, Sakuraba A. Systematic review with meta-analysis: the efficacy and safety of CT-P13, a biosimilar of anti-tumour necrosis factor- $\alpha$  agent (infliximab), in inflammatory bowel diseases. *Aliment Pharmacol Ther* 2017; **45**: 1043-1057 [PMID: 28239873 DOI: 10.1111/apt.13990]

**P- Reviewer:** Matowicka-Karna J, Ribaldone DG, Vasudevan A

**S- Editor:** Wang XJ **L- Editor:** A **E- Editor:** Yin SY







Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>



ISSN 1007-9327

