

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 41459

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort study

End-stage renal disease is associated with increased post endoscopic retrograde cholangiopancreatography adverse events in hospitalized patients

Sawas T *et al.* ERCP outcomes in renal disease

Tarek Sawas, Fateh Bazerbachi, Samir Haffar, Won K Cho, Michael J Levy, John A Martin, Bret T Petersen, Mark D Topazian, Vinay Chandrasekhara, Barham Abu Dayyeh

Abstract

AIM

To determine if end-stage renal disease (ESRD) is a risk factor for post endoscopic retrograde cholangiopancreatography (ERCP) adverse events.

Match Overview

1	Internet 28 words crawled on 14-Nov-2017 flboardofmedicine.gov	1%
2	Crossref 19 words Tarek Sawas, Fateh Bazerbachi, Barham K. Abu Dayyeh. "Sa 1421 Endoscopic Retrograde Cholangiopancreatography S ...	<1%
3	Internet 14 words crawled on 12-Nov-2017 stroke.ahajournals.org	<1%
4	Publications 14 words "Findings from E. Noel-Savina and Co-Authors Reveals Ne ... Information on Venous Thromboembolism (Tinza", Biotech W	<1%
5	Internet 13 words crawled on 21-Jul-2016 www.globalhealthaction.net	<1%



End-Stage Renal Disease Is Associated with Increased Post Endoscopic



All

News

Images

Videos

Shopping

More

Settings

Tools

About 82,800 results (0.67 seconds)

Anxiety and depression in patients with end-stage renal disease ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5856029/>

by ZS Goh - 2018 - Cited by 1 - Related articles

Mar 12, 2018 - In the final stage of CKD (ie, stage 5 or **end-stage renal disease [ESRD]**), renal **associated** with **increased** mortality and **hospitalization**.⁴¹ Several other ...

Patients who had symptoms of depression **after** kidney transplant were ... have found a trend of anxiety symptoms leading to **adverse clinical events**.

Missing: endoscopic retrograde cholangiopancreatography

Fluid hydration to prevent post-ERCP pancreatitis in average- to high ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5879873/>

by XJNM Smeets - 2018 - Cited by 1 - Related articles

Apr 2, 2018 - The primary endpoint is the incidence of **post-ERCP** pancreatitis. active gastrointestinal bleeding, ulcer **disease**, **renal insufficiency** ... Longer monitoring, **hospitalization**, and treatment of **adverse events** is Protocol version 3 is being used and **patient** recruitment is expected to **last** until the **end** of 2019.

Chronic Kidney Disease Adversely Influences Patient Safety

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2588106/> ▼

by SL Seliger - 2008 - Cited by 61 - Related articles

Patients with **chronic kidney disease** (CKD) may be at **higher** risk for **adverse** ... (29%) **hospitalized** veterans with CKD had a **higher** risk for several PSI, even **after** ... Among surgical **hospitalizations**, CKD was **associated** with **increased** risk for Strengths of the study include the examination of **patient safety events** across ...

Missing: endoscopic retrograde cholangiopancreatography

[PDF] Evaluation of the Safety of ERCP in Older Patients ... - SciELO Colom...

www.scielo.org.co/pdf/rcg/v30n2/en_v30n2a03.pdf ▼

by V Parra - 2015 - Related articles

of choledocholithiasis in **patients** over 70 years valued at University **Hospital** Clínica ... **post-ERCP** pancreatitis and **post** papilotomy bleeding; however, were not **related** to **patient** age. ... Any **adverse event related** to **ERCP** occurring within 30 ... COPD, chronic

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 141,000 条结果 (用时 0.55 秒)

Risk Factors of Post Endoscopic Retrograde ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3607778/> - 翻译此页

作者: MS Kwak - 2013 - 被引用次数: 14 - 相关文章

2012年11月13日 - Pseudomonas infection was significantly **more** common in **patients** ... The actual incidence of **post-ERCP** bacteremia remains unknown; ... cerebrovascular accident, **chronic kidney disease**, liver cirrhosis, ... Procedure **related** complications such as pancreatitis, bleeding and perforation were also reviewed.

Increased risk and severity of ERCP-related complications associated ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5585073/> - 翻译此页

作者: H Saito - 2017 - 被引用次数: 3 - 相关文章

2017年9月5日 - Because **patients** are asymptomatic, the risk of **ERCP-related** ... Complications of **ERCP** were defined as any **adverse events** occurring **after** the **ERCP** procedure that required **more** than 1 night of **hospitalization**. the 82 cases that underwent single-**stage** stone removal with EST or EPBD were shown.

Analysis of the risk factors for severity in post endoscopic retrograde ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5394726/> - 翻译此页

作者: H Matsubara - 2017 - 被引用次数: 2 - 相关文章

2017年4月16日 - At our **hospital**, **endoscopic retrograde cholangiopancreatography** (**ERCP**) was ... (**PEP**) is a typical endoscopy-**related** accident in the biliopancreatic field. ... **adverse reactions** of preventive drug administration, and cost[13]. was heart disease in 21 **patients**, diabetes in 24, **chronic kidney disease** in 41, ...

National trends of endoscopic retrograde cholangiopancreatography ...

https://www.researchgate.net/.../325979763_National_trends_of_endoscopic... - 翻译此页

2018年8月17日 - Limited data exist regarding its national utilization in **patients** with DC.... ... cirrhosis (DC) due to **increased** risk of **adverse events related** to liver dysfunction. ... and outcomes of **ERCP** among **patients** with DC in US **hospitalizations**. in **patients** with **end-stage renal disease** (**ESRD**) on hemodialysis (HD).

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 105,000 条结果 (用时 0.61 秒)

Risk Factors of Post Endoscopic Retrograde ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3607778/> ▼ [翻译此页](#)

作者: MS Kwak - 2013 - 被引用次数: 14 - 相关文章

2012年11月13日 - Pseudomonas infection was significantly **more** common in **patients** ... The actual incidence of **post-ERCP** bacteremia remains unknown; ... cerebrovascular accident, **chronic kidney disease**, liver cirrhosis, ... Procedure **related** complications such as pancreatitis, bleeding and perforation were also reviewed.

Increased risk and severity of ERCP-related complications associated ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5585073/> - [翻译此页](#)

作者: H Saito - 2017 - 被引用次数: 3 - 相关文章

2017年9月5日 - Because **patients** are asymptomatic, the risk of **ERCP-related** ... Complications of **ERCP** were defined as any **adverse events** occurring **after** the **ERCP** procedure that required **more** than 1 night of **hospitalization**. the 82 cases that underwent single-stage stone removal with EST or EPBD were shown.

Sa1421 Endoscopic Retrograde Cholangiopancreatography Safety in ...

[www.giejournal.org/article/S0016-5107\(17\)30714-9/fulltext](http://www.giejournal.org/article/S0016-5107(17)30714-9/fulltext) - [翻译此页](#)

作者: T Sawas - 2017

Chronic kidney disease (CKD) is an **increasing**, high prevalence public health problem. ... 48 **patients** developed **post ERCP** pancreatitis in the **ESRD** group (9.13%) compared to 1,530 in ... **ESRD** is **associated** with a significant health care burden in **hospitalized patients** requiring ... **Adverse events associated** with **ERCP**.

[PDF]

Evaluation of the Safety of ERCP in Older Patients ... - SciELO Colombia

www.scielo.org.co/pdf/rcg/v30n2/en_v30n2a03.pdf ▼ [翻译此页](#)

作者: V Parra - 2015 - 相关文章

of choledocholithiasis in **patients** over 70 years valued at University Hospital Clínica ... **post-ERCP** pancreatitis and **post** papilotomy bleeding; however, were not **related** to **patient** age. ... Any **adverse event related** to **ERCP** occurring within 30 COPD, chronic obstructive pulmonary disease; CKD: **chronic kidney disease** ...