



Histological predictive factors in patients with colorectal liver metastases



全部

新闻

图片

购物

地图

更多

设置

工具

找到约 197,000 条结果 (用时 0.68 秒)

Google 学术: Histological predictive factors in patients with colorectal liver metastases after neoadjuvant chemotherapy.

... liver metastases downstaged by chemotherapy: a ... - Adam - 被引用次数: 1543

... the outcome in patients with colorectal liver metastases ... - Rubbia-Brandt - 被引用次数: 329

... after hepatic resection for metastatic colorectal cancer: ... - Fong - 被引用次数: 3167

Histopathologic factors affecting tumor recurrence after hepatic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4091444/> - 翻译此页

作者: MS Park - 2014 - 被引用次数: 10 - 相关文章

2014年6月24日 - Keywords: Colorectal neoplasm, Liver metastasis, Recurrence, Hepatic ...

Neoadjuvant chemotherapy was given in patients with high preoperative CEA The positive histopathologic markers predicting tumor recurrence, i.e., ...

Long-term postoperative survival prediction in patients with colorectal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5668107/> - 翻译此页

作者: K Wang - 2017 - 被引用次数: 2 - 相关文章

2017年8月18日 - Numerous factors affect the prognosis of colorectal liver metastasis (CRLM) patients after hepatic resection. Neoadjuvant chemotherapy has been also proposed for patients with resectable disease and B. Importance of histological tumor response assessment in predicting the outcome in patients with ...



Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 42297

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Predictive factors of histological response of colorectal liver metastases after neoadjuvant chemotherapy

Serayssol C *et al.* Histological TRG and liver metastases

Chloé Serayssol, Charlotte Maulat, Florence Breibach, Fatima-Zohra Mokrane, Janick Selves, Rosine Guimbaud, Philippe Otal, Bertrand Suc, Emilie Berard, Fabrice Muscari

Abstract

BACKGROUND

Match Overview

1	Crossref 47 words "Abstracts", Journal of Thoracic Oncology, 2009	1%
2	Crossref 42 words "Abstracts", HPB, 2007	1%
3	Crossref 38 words "Abstracts", HPB, 2008	1%
4	Internet 32 words crawled on 14-Dec-2018 www.wjgnet.com	1%
5	Internet 21 words crawled on 28-Nov-2018 publications.iarc.fr	<1%
6	Internet 20 words crawled on 28-Jan-2017 e-sciencecentral.org	<1%
7	Internet 11 words crawled on 22-Nov-2006 www.ewma.org	<1%
8	Internet 9 words crawled on 01-Aug-2018 insights.ovid.com	<1%

找到约 249,000 条结果 (用时 0.50 秒)

Google 学术 : Predictive factors of histological response of colorectal liver metastases after neoadjuvant chemotherapy

... liver metastases following infusional chemotherapy ... - Giacchetti - 被引用次数 : 767

... histological tumor response assessment in predicting ... - Rubbia-Brandt - 被引用次数 : 341

... liver metastases downstaged by chemotherapy: a ... - Adam - 被引用次数 : 1572

(PDF) Neoadjuvant Chemotherapy in Patients with Stage IV Colorectal ...

https://www.researchgate.net/.../43202510_Neoadjuvant_Chemotherapy_in_... - 翻译此页

2018年7月31日 - ... stage IV colorectal cancer, liver metastases exhibit a better histological response than ... pathologic response after neoadjuvant chemotherapy and, resection of hepatic metastases is an independent predictor of 5-year overall ... as a prognostic factor for disease-free and overall survival. 16. The high ...

(PDF) Retrospective analysis of pathological response in colorectal ...

https://www.researchgate.net/.../259336015_Retrospective_analysis_of_path... - 翻译此页

2018年7月31日 - PDF | Pathological response has been shown to be a predictor for ... neoadjuvant chemotherapy in patients with metastatic colorectal ... cancer liver metastases following treatment with bevacizumab Patients with a histologically confirmed diagnosis of stage Predicting factors in acute pancreatitis.

(DFS) of TRG of resected colorectal liver metastases (CRLM) after ...

[https://www.hpbonline.org/article/S1365-182X\(18\)34050-4/abstract](https://www.hpbonline.org/article/S1365-182X(18)34050-4/abstract) - 翻译此页

作者 : C Maulat - 2018

Predictive factors and impact on overall survival (OS) and disease-free survival ... colorectal liver metastases (CRLM) after neoadjuvant chemotherapy (NACT) ... the predictive factors of histological response and its impact on OS and DFS.

找到约 247,000 条结果 (用时 0.48 秒)

Google 学术 : Predictive factors of histological response of colorectal liver metastases after neoadjuvant chemotherapy

... liver metastases following infusional chemotherapy ... - Giacchetti - 被引用次数 : 765

... histological tumor response assessment in predicting ... - Rubbia-Brandt - 被引用次数 : 341

... liver metastases downstaged by chemotherapy: a ... - Adam - 被引用次数 : 1574

(PDF) Neoadjuvant Chemotherapy in Patients with Stage IV Colorectal ...

https://www.researchgate.net/.../43202510_Neoadjuvant_Chemotherapy_in_... - 翻译此页

2018年7月31日 - ... stage IV colorectal cancer, liver metastases exhibit a better histological response than ... pathologic response after neoadjuvant chemotherapy and, resection of hepatic metastases is an independent predictor of 5-year overall ... as a prognostic factor for disease-free and overall survival. 16. The high ...

Factors Predicting Response to Neoadjuvant Therapy of Hepatic ...

<https://clinmedjournals.org/.../international-journal-of-pathology-and-clinical-...> - 翻译此页

Factors Predicting Response to Neoadjuvant Therapy of Hepatic Metastasis ... patients should be operated on after chemotherapy regardless of response. ... Colorectal carcinoma, Neoadjuvant therapy, Prognosis, Histopathological response.

Proposal for novel histological findings of colorectal liver metastases ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4690511/> ▼ 翻译此页

作者 : K Ishida - 2015 - 被引用次数 : 3 - 相关文章

2015年5月4日 - Keywords: chemotherapy regimen, cholesterol clefts, colorectal liver ... neoadjuvant chemotherapy alters the natural history of unresectable CRLM by ... that the pathological tumor response to chemotherapy is an important factor in ... The following histological features were evaluated: usual necrosis (UN), ...

Recent Advances in the Curative Treatment of Colorectal Liver ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3283006/> ▼ 翻译此页

作者 : A Andreou - 2011 - 被引用次数 : 20 - 相关文章

Neoadjuvant chemotherapy resulted in a cumulative response rate of 66%, and 34% ... R0 resection, which was a major predictive factor for prolonged survival. Thus, new diagnostic tools, such as histologic analysis of the resected tumor, have liver metastases from colorectal cancer after neoadjuvant chemotherapy.