

World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2019 March 15; 11(3): 181-269



ORIGINAL ARTICLE**Basic Study**

- 181 Glycerophospholipids pathways and chromosomal instability in gastric cancer: Global lipidomics analysis
Hung CY, Yeh TS, Tsai CK, Wu RC, Lai YC, Chiang MH, Lu KY, Lin CN, Cheng ML, Lin G
- 195 Human colorectal cancer cells frequently express IgG and display unique Ig repertoire
Geng ZH, Ye CX, Huang Y, Jiang HP, Ye YJ, Wang S, Zhou Y, Shen ZL, Qiu XY

Retrospective Cohort Study

- 208 Post-operative computed tomography scan – reliable tool for quality assessment of complete mesocolic excision
Livadaru C, Morarasu S, Frunza TC, Ghitun FA, Paiu-Spiridon EF, Sava F, Terinte C, Ferariu D, Lunca S, Dimofte GM
- 227 Hepatic resection vs percutaneous radiofrequency ablation of hepatocellular carcinoma abutting right diaphragm
Song KD, Lim HK, Rhim H, Lee MW, Kang TW, Paik YH, Kim JM, Joh JW

Observational Study

- 238 Risk of cholangiocarcinoma in patients undergoing therapeutic endoscopic retrograde cholangiopancreatography or cholecystectomy: A population based study
Wang CC, Tsai MC, Sung WW, Yang TW, Chen HY, Wang YT, Su CC, Tseng MH, Lin CC

SYSTEMATIC REVIEWS

- 250 Near-infrared fluorescence guided esophageal reconstructive surgery: A systematic review
Van Daele E, Van Nieuwenhove Y, Ceelen W, Vanhove C, Braeckman BP, Hoorens A, Van Limmen J, Varin O, Van de Putte D, Willaert W, Pattyn P

CASE REPORT

- 264 Stent placement followed by preoperative chemotherapy and elective surgery for acute malignant colorectal obstruction: Six cases of report
Liu JJ, Ma TH, Qin QY, Wang L

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Oncology*, Marco E Allaix, MD, PhD, Assistant Professor, Department of Surgical Sciences, University of Torino, Torino 10126, Italy

AIMS AND SCOPE

World Journal of Gastrointestinal Oncology (*World J Gastrointest Oncol*, *WJGO*, online ISSN 1948-5204, DOI: 10.4251) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJGO covers topics concerning carcinogenesis, tumorigenesis, metastasis, diagnosis, prevention, prognosis, clinical manifestations, nutritional support, etc. The current columns of *WJGO* include editorial, frontier, field of vision, review, original articles, case report.

We encourage authors to submit their manuscripts to *WJGO*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great clinical significance.

INDEXING/ABSTRACTING

The *WJGO* is now indexed in Science Citation Index Expanded (also known as SciSearch®), PubMed, and PubMed Central. The 2018 edition of Journal Citation Reports® cites the 2017 impact factor for *WJGO* as 3.140 (5-year impact factor: 3.228), ranking *WJGO* as 39 among 80 journals in gastroenterology and hepatology (quartile in category Q2), and 114 among 222 journals in oncology (quartile in category Q3).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Han Song*

Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Monjur Ahmed, Rosa M Jimenez Rodriguez, Pashtoon Murtaza Kasi

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

EDITORIAL OFFICE

Jin-Lei Wang, Director

PUBLICATION DATE

March 15, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Retrospective Cohort Study

Hepatic resection vs percutaneous radiofrequency ablation of hepatocellular carcinoma abutting right diaphragm

Kyoung Doo Song, Hyo Keun Lim, Hyunchul Rhim, Min Woo Lee, Tae Wook Kang, Yong Han Paik, Jong Man Kim, Jae-Won Joh

ORCID number: Kyoung Doo Song (0000-0002-2767-3622); Hyo Keun Lim (0000-0003-3269-7503); Hyunchul Rhim (0000-0002-9737-0248); Min Woo Lee (0000-0001-9048-9011); Tae Wook Kang (0000-0002-8991-6407); Yong Han Paik (0000-0002-3076-2327); Jong Man Kim (0000-0002-1903-8354); Jae-Won Joh (0000-0003-1732-6210).

Author contributions: Song KD and Lim HK designed the study, analyzed data and wrote the manuscript; Lim HK, Rhim H, Lee MW, Paik YH, Joh JW and Kim JM were all responsible for the provision of study materials or patients; Kang TW and Kim JM contributed to collection and assembly of data; all authors provided critical review of the manuscript and approved the final version of the manuscript.

Institutional review board statement: This study was reviewed and approved by the Institutional Review Board of the Samsung Medical Center.

Informed consent statement: Informed consent was waived by the Institutional Review Board of the Samsung Medical Center because this study was a retrospective study.

Conflict-of-interest statement: All authors declare no conflicts of interest related to this article.

Open-Access: This article is an open-access article which was

Kyoung Doo Song, Hyunchul Rhim, Min Woo Lee, Tae Wook Kang, Department of Radiology, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul 06351, South Korea

Hyo Keun Lim, Department of Radiology, Samsung Medical Center, Sungkyunkwan University School of Medicine, Department of Health Sciences and Technology, SAIHST, Sungkyunkwan University, Seoul 06351, South Korea

Yong Han Paik, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul 06351, South Korea

Jong Man Kim, Jae-Won Joh, Department of Surgery, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul 06351, South Korea

Corresponding author: Hyo Keun Lim, MD, PhD, Professor, Department of Radiology and Department of Health Sciences and Technology, Samsung Medical Center, SAIHST, Sungkyunkwan University School of Medicine, 81 Irwon-ro, Gangnam-gu, Seoul 06351, South Korea. hyokeun.lim@samsung.com

Telephone: +82-2-34102505

Fax: +82-2-34102559

Abstract

BACKGROUND

It is usually difficult to adequately conduct percutaneous ultrasound-guided radiofrequency (RF) ablation for hepatocellular carcinomas (HCCs) abutting the diaphragm. Our hypothesis was that the subphrenic location of HCC could have an effect on the long-term therapeutic outcomes after hepatic resection and RF ablation.

AIM

To compare the long-term therapeutic outcomes of hepatic resection and percutaneous RF ablation for HCCs abutting the diaphragm.

METHODS

A total of 143 Child-Pugh class A patients who had undergone hepatic resection ($n = 80$) or percutaneous ultrasound-guided RF ablation ($n = 63$) for an HCC (≤ 3 cm) abutting the right diaphragm were included. Cumulative local tumor progression (LTP), cumulative intrahepatic distant recurrence (IDR), disease-free survival (DFS), and overall survival (OS) rates were estimated. Prognostic factors

selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: October 23, 2018

Peer-review started: October 23, 2018

First decision: November 29, 2018

Revised: December 12, 2018

Accepted: January 5, 2019

Article in press: January 6, 2019

Published online: March 15, 2019

for DFS and OS were analyzed. Complications were evaluated.

RESULTS

The cumulative IDR rate, DFS rate, and OS rate for the hepatic resection group and RF ablation group at 5 years were "35.9% vs 65.8%", "64.1% vs 18.3%", and "88.4% vs 68.7%", respectively. Hepatic resection was an independent prognostic factor for DFS ($P \leq 0.001$; hazard ratio, 0.352; 95% CI: 0.205, 0.605; with RF ablation as the reference category); however, treatment modality was not an independent prognostic factor for OS. The LTP rate was 46.6% at 5 years for the RF ablation group. The major complication rate was not significantly different between the groups ($P = 0.630$). The rate of occurrence of peritoneal seeding was higher in the RF ablation group (1.3% vs 9.5%, $P = 0.044$).

CONCLUSION

Although OS was not significantly different between patients who had gone hepatic resection or percutaneous RF ablation for HCCs abutting the diaphragm, DFS was better in the hepatic resection group.

Key words: Hepatic resection; Radiofrequency ablation; Hepatocellular carcinoma; Diaphragm; Treatment outcome

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The aim of this study was to compare the long-term therapeutic outcomes of hepatic resection and percutaneous radiofrequency (RF) ablation for hepatocellular carcinomas abutting the diaphragm. The disease-free survival (DFS) rate was 64.1% and 18.3% for the hepatic resection group and the RF ablation group, and overall survival (OS) rate was 88.4% and 68.7% for the hepatic resection group and the RF ablation group at 5 years. The local tumor progression rate was as high as 46.6% for the RF ablation group. Although OS was not significantly different between two groups, DFS was better in the hepatic resection group.

Citation: Song KD, Lim HK, Rhim H, Lee MW, Kang TW, Paik YH, Kim JM, Joh JW. Hepatic resection vs percutaneous radiofrequency ablation of hepatocellular carcinoma abutting right diaphragm. *World J Gastrointest Oncol* 2019; 11(3): 227-237

URL: <https://www.wjgnet.com/1948-5204/full/v11/i3/227.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v11.i3.227>

INTRODUCTION

Both hepatic resection and radiofrequency (RF) ablation are considered curative procedures for very early or early-stage hepatocellular carcinoma (HCC)^[1]. Many studies have revealed that RF ablation is comparable to hepatic resection in terms of long-term survival for patients with early-stage HCC^[2-4]. However, most studies have not taken into account the location of HCCs. Tumor location is an important factor affecting local tumor control especially for RF ablation due to its technical complexity^[5].

When an HCC is located in the liver abutting the right diaphragm, an adequate accomplishment of percutaneous ultrasound (US)-guided RF ablation is difficult due to the poor sonic window resulting from lung shadowing and the potential risk of collateral thermal injury to the diaphragm. According to a preliminary study, local tumor progression (LTP) after percutaneous RF ablation was more frequent in patients with subphrenic HCCs (29%) than in nonsubphrenic HCCs (6%)^[6]. To overcome this inherent limitation, many investigators have used the infusion of artificial ascites or pleural effusion. Several studies have reported that percutaneous RF ablation with infusion of artificial ascites or pleural effusion was safe and effective^[7-10]. However, the LTP rate after RF ablation for subphrenic HCCs remained high even with the application of these special techniques^[9]. The effect of the specific location of HCC on the long-term therapeutic outcomes after hepatic resection and RF ablation has not yet been investigated. Thus, the aim of this study was to compare the long-term therapeutic outcomes of hepatic resection vs percutaneous RF ablation for

the curative treatment of HCCs abutting the diaphragm.

MATERIALS AND METHODS

Our Institutional Review Board approved this retrospective study, and informed consent was waived.

Patients

Between January 2006 and October 2010, 5981 patients were diagnosed with HCC at our institution. This study included patients from the same population as in a previous study that was conducted at our institution; however, the study design and result analysis methods are different^[11]. Inclusion criteria for our study were as follows: (1) patients who had undergone percutaneous US-guided RF ablation or hepatic resection for HCC as a first-line treatment; (2) patients who had a single HCC ≤ 3 cm; (3) patients with HCC abutting the right diaphragm (subphrenic HCC); and (4) patients with Child-Pugh class A. A subphrenic HCC in our study was defined as a tumor that abutted the right diaphragm on axial or coronal images of computed tomography (CT) or magnetic resonance imaging. We excluded tumors that abutted the left diaphragm because they are different from the tumors abutting the right diaphragm in many ways, in terms of treatment. Most tumors abutting the left diaphragm are located under the heart and are, hence, considered more technically difficult to treat compared to those close to the right diaphragm. In addition, the use of artificial ascites or pleural effusion is usually ineffective for tumors abutting the left diaphragm. Instead, hepatic resection of tumors abutting the left diaphragm (especially in the left lateral segment) is easily performed either after laparotomy or with a laparoscopic approach. Finally, our study included 63 patients (49 men, 14 women; mean age, 60.3 years; range, 41–78 years) who had undergone percutaneous RF ablation and 80 patients (62 men, 18 women; mean age, 53.5 years; range, 30–78 years) who had been treated with hepatic resection. The patient inclusion flowchart is shown in [Figure 1](#).

In 3 (4.7%) patients in the RF ablation group, HCC was confirmed histologically via percutaneous US-guided biopsy. In the remainder of the patients in the RF ablation group, HCC was diagnosed based on one of two clinical guidelines from the American Association for the Study of Liver Diseases at the time of RF ablation^[1,12]. For all patients in the hepatic resection group, HCC was diagnosed histologically after hepatic resection.

Treatment of HCC and follow-up

The general inclusion criteria for hepatic resection at our institution were as follows: (1) a single tumor or oligonodular tumors within a monosegment of the liver; (2) an indocyanine green retention rate less than 20% at 15 min; (3) serum total bilirubin level less than 1.5 mg/dL; (4) no severe portal hypertension; and (5) no gross ascites. The inclusion criteria for percutaneous RF ablation at our institution were as follows: (1) a single tumor (≤ 5 cm in the greatest dimension) or multiple nodular tumors (three or fewer, each ≤ 3 cm in the greatest dimension); (2) Child-Pugh class A or B disease; (3) no evidence of portal vein thrombosis or extrahepatic metastasis; and (4) prothrombin time ratio $> 50\%$, and platelet count $> 50000/\text{mm}^3$ ($50 \times 10^9/\text{L}$). Treatment modality was decided based on age, liver function reserve, tumor location, surgical risk, and patient preference by a multidisciplinary tumor board composed of hepatologists, radiologists, surgeons, and medical and radiation oncologists.

Hepatic resection was performed by one of two surgeons (JHK and JWJ) with more than 10 years of experience in hepatobiliary surgery by the end of the study. The types of hepatic resection were as follows: subsegmentectomy in 58 patients, bisegmentectomy in five patients, posterior sectionectomy in 12 patients, right hemihepatectomy in two patients, anterior sectionectomy in one patient, central hepatectomy in one patient, and extended left hemihepatectomy in one patient. As a result, anatomical resection was performed in 17 (21.3%) patients and non-anatomical resection was performed in 63 (78.8%) patients^[13]. Hepatic resection was performed after laparotomy in 78 (97.5%) patients and with laparoscopy in two (2.5%) patients. RF ablation was performed by one of five interventional radiologists (MWL, DC, HR, HKL, and YK) with more than 6 years of experience in RF ablation by the end of the study. The process and method of RF ablation were the same as those described in a previous study^[14]. In brief, RF ablation was performed percutaneously under the guidance of real-time US. We used internally cooled electrode systems with generators (Cool-tip RF System, Covidien, Mansfield, MA, United States; or VIVA RFA System, STARmed, Goyang, South Korea). Sedation was performed via an intravenous injection of pethidine hydrochloride (Samsung Pharmaceuticals, Seoul,




Figure 1 Flowchart of patient inclusion. HCC: Hepatocellular carcinoma; RF: Radiofrequency.

South Korea) and fentanyl citrate (GUJU Pharma, Seoul, South Korea). To improve the sonic window and avoid thermal injury to the diaphragm, artificial ascites (5% dextrose in a water solution) was infused into the perihepatic space using a 5F angiosheath in 39 (61.9%) patients.

After RF ablation, immediate follow-up contrast agent-enhanced CT was performed to evaluate the therapeutic response and possible complications. Contrast agent-enhanced CT was performed at the 1 mo follow-up, every 3 mo during the first 2 years, followed by every 4-6 mo according to the risk of recurrence for both the hepatic resection group and RF ablation group.

Data acquisition

Baseline characteristics of patients and HCCs were obtained through review of their electronic medical record from our institution. To compare the therapeutic outcomes between the two groups, intrahepatic distant recurrence (IDR), disease-free survival (DFS), and overall survival (OS) were calculated. IDR was defined as a new tumor appearing in the liver separate from the treated area. DFS was defined as the time interval from the date of treatment to one of the following events: intrahepatic recurrence, extrahepatic recurrence, or death. OS was defined as the time interval from the date of treatment to death. If the patients had undergone liver transplantation, they were considered to have been censored at the time of liver transplantation. Complications were stratified according to the Clavien classification of postoperative complications, and complications of grade II or higher were considered major complications^[15]. Local tumor progression (LTP) was evaluated for the RF ablation group. LTP was defined as the appearance of new tumor foci at the margin of the ablation zone after at least one contrast-enhanced follow-up study had demonstrated an absence of viable tumors^[16].

Statistical analysis

Continuous data were compared using two-sample *t* tests, and categorical variables were compared using chi-squared tests between the two groups. Cumulative LTP, cumulative IDR, DFS, and OS rates were estimated using the Kaplan-Meier method. Prognostic factors for DFS and OS were assessed using Cox regression models. Proportional hazard (PH) assumption for the Cox proportional hazard model was tested using Schoenfeld's method. For the variables with violation of PH assumption, the time-dependent Cox regression was applied. When the time dependence was not significant, the Cox proportional hazard model was applied. Possible risk factors with *P* values of 0.1 or less at univariate analyses were entered into the multivariate Cox proportional hazard models. Subgroup analysis for patients with ≤ 2 cm HCCs was performed with Cox proportional hazard models. All statistical analyses were performed using a software (PASW statistical software, version 18.0; SPSS, Chicago, IL). For all tests, a *P* value < 0.05 was defined as a significant difference.

RESULTS

Baseline characteristics of patients and HCCs are shown in [Table 1](#). The median follow-up period was 74.9 mo (range, 10.3-117.8 mo) in the hepatic resection group and 65.3 mo (range, 4.1-113.9 mo) in the RF ablation group. The RF ablation group was significantly older, and they exhibited a lower α -fetoprotein level, platelet count, and serum albumin level, and a higher prothrombin time. In the RF ablation group, the proportion of patients with liver cirrhosis and hepatitis C virus was higher and the proportion of patients with hepatitis B virus was lower compared to that in the hepatic resection group. The mean size of HCCs was not significantly different between the two groups.

Therapeutic outcomes

The cumulative IDR rates at 1-, 3-, and 5-years were 15.0%, 29.1%, and 35.9%, respectively, for the hepatic resection group and 13.1%, 54.5%, and 65.8%, respectively, for the RF ablation group ([Figure 2A](#)). The estimated DFS rates at 1-, 3-, and 5-years were 85.0%, 70.9%, and 64.1%, respectively, for the hepatic resection group and 69.5%, 27.5%, and 18.3%, respectively, for the RF ablation group ([Figure 2B](#)). The estimated OS rates at 1-, 3-, and 5-years were 97.5%, 92.3%, and 88.4%, respectively, for the hepatic resection group and 100%, 81.4%, and 68.7%, respectively, for the RF ablation group ([Figure 2C](#)). For the RF ablation group, the cumulative LTP rates were 22.5%, 37.8%, and 46.6% at 1-, 3-, and 5-years, respectively ([Figure 3](#)).

Analysis of risk factors

Based on multivariate analysis, there was no independent prognostic factor for OS. Hepatic resection [$P \leq 0.001$; hazard ratio (HR), 0.352; 95% confidence interval (CI): 0.205, 0.605; with RFA as the reference category], alanine aminotransferase level ($P = 0.006$; HR, 1.011; 95%CI: 1.003, 1.020), and serum albumin level ($P = 0.014$; HR, 0.481; 95%CI: 0.269, 0.860) were independent prognostic factors for DFS ([Tables 2 and 3](#)).

Subgroup analysis for patients with ≤ 2 cm HCC

Thirty-seven patients in the hepatic resection group and 27 patients in the RF ablation group had ≤ 2 cm HCC. The cumulative IDR rates at 1-, 3-, and 5-years were 13.5%, 27.3%, and 33.1%, respectively, for the hepatic resection group and 15.3%, 60.3%, and 70.2%, respectively, for the RF ablation group. The estimated DFS rates at 1-, 3-, and 5-years were 86.5%, 72.7%, and 66.9%, respectively, for the hepatic resection group and 81.0%, 27.4%, and 18.3%, respectively, for the RF ablation group. The estimated OS rates at 1-, 3-, and 5-years were 100%, 94.5%, and 91.7%, respectively, for the hepatic resection group and 100%, 83.8%, and 65.4%, respectively, for the RF ablation group. In multivariate analysis, hepatic resection was an independent prognostic factor for DFS ($P = 0.018$; HR, 0.365; CI: 0.158-0.844), but was not an independent prognostic factor for OS.

Complications and treatment for recurrent HCC

There was no treatment-related mortality in either group. Major complications occurred in three patients (3.8%) in the hepatic resection group: Grade II, pneumonia ($n = 1$) and intraperitoneal hemorrhage ($n = 1$); and Grade III, wound infection requiring surgery ($n = 1$). In the RF ablation group, a major complication occurred in one patient (1.6%): Grade III, pleural effusion requiring drainage. The major complication rate was not significantly different between the two groups ($P = 0.060$). The posttreatment hospital stay was significantly longer in the hepatic resection group (median, 9 d; range, 5-23 d) than in the RF ablation group (median, 1.0 d; range, 1-4 d; $P < 0.001$).

During the follow-up period, peritoneal seeding occurred in one patient (1.3%) in the hepatic resection group and six patients (9.5%) in the RF ablation group, and the rate of peritoneal seeding was significantly different ($P = 0.044$).

During the follow-up period, LTP occurred in 29 (46.0%) of the 63 patients in the RF ablation group. The initial treatment modalities for LTP were as follows: transarterial chemoembolization (TACE) ($n = 14$), RF ablation ($n = 12$), hepatic resection ($n = 1$), combined TACE and RF ablation ($n = 1$), and combined TACE and radiation therapy ($n = 1$). In 26 of 29 patients, LTP was controlled with additional treatments, and the number of additional treatments was as follows: One ($n = 17$), two ($n = 3$), three ($n = 4$), and six ($n = 2$). For the remaining three patients, LTP was not controlled even though they received repeated treatments with TACE or RF ablation. In addition, multiple intra- and extrahepatic metastases occurred. Finally, sorafenib treatment was administered. IDR occurred in 31 (38.8%) of the 80 patients in the hepatic resection group, and treatment modalities were as follows: TACE ($n = 18$), RF ablation ($n = 11$), cryoablation ($n = 1$), and hepatic resection ($n = 1$). IDR occurred in 42 (66.7%) of the 63

Table 1 Baseline patient characteristics

Variable	Hepatic resection (n = 80)	RF ablation (n = 63)	P value
Mean age (yr)	53.5 ± 9.0	60.3 ± 8.7	< 0.001
Male sex	62 (78)	49 (78)	0.968
Etiology			0.042
HBV	68 (85)	43 (68)	
HCV	6 (8)	13 (20)	
NBNC	6 (8)	7 (11)	
Liver cirrhosis	50 (63)	50 (79)	0.029
Tumor size	2.1 ± 0.6	2.1 ± 0.5	0.906
α-fetoprotein (ng/mL)	200 ± 489	72 ± 165	0.031
Platelet count (× 10 ³ /mm ³)	148 ± 49	107 ± 49	< 0.001
Alanine aminotransferase (IU/L)	41 ± 21	40 ± 30	0.849
Total bilirubin (mg/dL)	0.7 ± 0.3	0.7 ± 0.4	0.99
Albumin (g/dL)	4.2 ± 0.4	3.9 ± 0.4	< 0.001
Prothrombin time (INR)	1.1 ± 0.1	1.2 ± 0.1	< 0.001

Continuous data were evaluated using two-sample *t* tests and categorical variables were analyzed using Chi-square tests. Data represent the number of patients with percentage in parentheses or the mean ± SD. RF: Radiofrequency; HBV: Hepatitis B virus; HCV: Hepatitis C virus; NBNC: Non-B non-C; INR: International normalized ratio.

patients in the RF ablation group, and treatment modalities were as follows: TACE (*n* = 16), RF ablation (*n* = 20), combined TACE and RF ablation (*n* = 3), hepatic resection (*n* = 1), liver transplantation (*n* = 1), and sorafenib treatment (*n* = 1).


DISCUSSION

In our study, we compared long-term therapeutic outcomes for treatments using hepatic resection and percutaneous RF ablation for HCCs (≤ 3 cm) abutting the right diaphragm; we found that the treatment modality was a significant prognostic factor for DFS, but was not an independent prognostic factor for OS. For the RF ablation group, the LTP rate was as high as 46.6% at 5 years. The location of tumors can affect the technical difficulty in local control of tumors, especially for RF ablation. Although there have been many studies that compared therapeutic outcomes between hepatic resection and RF ablation for HCC, most of them did not consider the location of tumors. In this way, the results of our study, which compares hepatic resection and percutaneous RF ablation for HCCs with consideration of the location of tumors, can provide important data for the proper management of HCCs abutting the diaphragm.


In our study, the LTP rate was 46.6% at 5 years for the RF ablation group. The LTP rate was much higher than rates reported in previous studies that included all HCCs located in the liver^[11,14,17-19]. Percutaneous RF ablation for subphrenic HCCs is difficult to adequately perform for several reasons. First, the poor sonic window resulting from the lung shadow makes it difficult to accurately target tumors with the electrodes. Second, all tumors were subcapsular HCCs in our study. In general, subcapsular HCCs are considered to be more difficult to treat with percutaneous HCC than nonsubcapsular HCCs because of the difficulty of placing an electrode and not being able to obtain enough ablative margin along the hepatic capsule.

In this study, patients who had undergone hepatic resection exhibited longer DFS compared to those who had undergone RF ablation. This result is in line with previous studies that compared DFS outcomes for hepatic resection and RF ablation for HCC^[20,21]. In our study, the estimated DFS rates at 1-, 3-, and 5-years were 85.0%, 70.9%, and 64.1%, respectively, for the hepatic resection group and 69.5%, 27.5%, and 18.3%, respectively, for the RF ablation group. In the previous study at our institution that compared RF ablation with hepatic resection for single HCC ≤ 3 cm located in the liver, the estimated DFS rate at 5 years was 61.1% for the hepatic resection group and 31.7% for the RF ablation group^[11]. The DFS rate for the hepatic resection group of this study was similar to our previous result. However, the DFS rate for the RF ablation group of this study was lower than our previous result. This difference can most likely be explained by the high LTP rate for the RF ablation group in this study.


According to previous studies, RF ablation was comparable to hepatic resection for



Number at risk										
Hepatic resection	80	68	58	53	51	46	28	19	9	7
RF ablation	63	53	37	24	21	18	9	2	2	0



Number at risk										
Hepatic resection	80	68	58	53	51	46	27	18	8	6
RF ablation	63	43	27	15	12	10	4	2	1	0



Number at risk										
Hepatic resection	80	78	76	71	69	66	43	30	12	7
RF ablation	63	61	57	46	41	37	21	9	7	4

Figure 2 Cumulative intrahepatic distant recurrence rates (A), disease-free survival rates (B), and overall survival rates (C). IDR: Intrahepatic distant recurrence; RF: Radiofrequency.

very early and early-stage HCCs in terms of OS^[22-24]. In our study, estimated OS rates for the hepatic resection group (97.5%, 92.3%, and 88.4% at 1-, 3-, and 5-years, respectively) appeared to be better than those for the RF ablation group (100%, 81.4%, and 68.7% at 1-, 3-, and 5-years). However, similar to previous studies, treatment modality was not an independent prognostic factor for OS according to multivariate




Figure 3 Local tumor progression rate in the radiofrequency ablation group. LTP: Local tumor progression.

analyses in our study.

Previous studies have reported comparable outcomes between RF ablation and hepatic resection in terms of long-term survival for patients with early-stage HCC. Based on these results, both hepatic resection and RF ablation are considered as curative treatment options for early stage HCC. Although treatment modality was not an independent prognostic factor for OS in patients with subphrenic HCCs, there were some differences in treatment outcomes between patients with subphrenic HCCs and nonsubphrenic HCCs that need to be considered when treatment modality is determined. First, the LTP rate after RF ablation was much higher for patients with subphrenic HCCs. Second, recurrent LTP was common in patients with subphrenic HCCs. In 12 (41%) of 29 patients who had LTP, multiple treatments were performed to control the LTP. Third, the peritoneal seeding rate for subphrenic HCCs was as high as 9.5% in the RF ablation group. Considering these unfavorable outcomes of RF ablation for subphrenic HCCs, it may be reasonable to preferentially consider hepatic resection as the first-line treatment for subphrenic HCCs rather than percutaneous RF ablation. Otherwise, laparoscopic RF ablation or combined TACE and RF ablation should be considered because these modalities can be more effective than percutaneous RF ablation alone in terms of local tumor control^[25-27]. However, this issue needs to be investigated further.

Our study has some limitations. First, because this is a retrospective study, the treatment groups were not randomized, and we could not exclude the possibility of selection bias. However, we analyzed the effect of treatment modality (hepatic resection *vs* percutaneous RF ablation) after controlling for potential compounding factors. Second, HCC was diagnosed based on clinical guidelines in most patients in the RF ablation group. Therefore, there was a possibility of false-positive diagnosis, which could affect the outcomes. Third, this is a single-center study. In general, the outcomes of both hepatic resection and RF ablation greatly depend on the expertise and experience of the operators. In addition, we only used the single straight type of RF electrode and US as a guiding modality. Using other types of RF electrodes or guiding modalities may result in different therapeutic outcomes. Therefore, care should be taken when generalizing our results to that from other institutions.

In conclusion, although OS was not significantly different between patients who had undergone hepatic resection or percutaneous RF ablation for HCCs abutting the diaphragm, DFS was better in the hepatic resection group, and LTP was as high as 46.6% at 5 years in the RF ablation group. Therefore, it may be reasonable that hepatic resection should be preferentially considered over percutaneous US-guided RF ablation as a first-line treatment for HCCs abutting the diaphragm.

Table 2 Univariate and multivariate analysis of prognostic factors for overall survival

Variable	Univariate analysis		Multivariate analysis	
	HR	P value	HR	P value
Group (RF ablation)	0.364 (0.179, 0.742)	0.005	0.676 (0.309, 1.482)	0.329
Age	1.025 (0.988, 1.063)	0.187		
Sex (female)	0.975 (0.439, 2.165)	0.95		
Etiology (hepatitis B virus) ¹		0.028		0.176
Hepatitis C virus	3.053 (1.240, 7.516)	0.01	2.180 (0.854, 5.566)	0.124
NBNC	2.292 (0.671, 7.831)	0.26	2.433 (0.686, 8.636)	0.232
Liver cirrhosis (absence)	1.834 (0.756, 4.446)	0.18		
Tumor size	1.505 (0.782, 2.899)	0.221		
α -fetoprotein	1.000 (1.000, 1.001)	0.345		
Platelet count	0.989 (0.982, 0.996)	0.003	0.991 (0.982, 1.000)	0.051
Alanine aminotransferase	1.006 (0.993, 1.019)	0.36		
Total bilirubin	1.031 (0.393, 2.704)	0.951		
Albumin	0.327 (0.148, 0.727)	0.006	0.485 (0.189, 1.244)	0.132
Prothrombin time (INR)	19.351 (1.354, 1798.539)	0.034	0.325 (0.002, 46.731)	0.657

The Cox proportional hazards model was used for univariate and multivariate analysis. The reference category for each categorical variable is provided in the square brackets in the first column.

¹Bonferroni correction was used owing to multiple comparisons. Numbers in parentheses represent the 95%CI. HR: Hazard ratio; RF: Radiofrequency; NBNC: Non-B non-C; INR: International normalized ratio.

Table 3 Univariate and multivariate analysis of prognostic factors for disease-free survival

Variable	Univariate analysis		Multivariate analysis	
	HR	P value	HR	P value
Group (RF ablation)	0.272 (0.174, 0.427)	< 0.001	0.352 (0.205, 0.605)	< 0.001
Age	1.036 (1.012, 1.060)	0.003	1.015 (0.987, 1.043)	0.306
Sex (female)	1.096 (0.650, 1.847)	0.732		
Etiology (hepatitis B virus) 1		0.726		
Hepatitis C virus	1.507 (0.789, 2.878)	0.31		
NBNC	1.121 (0.483, 2.601)	1		
Liver cirrhosis (absence)	1.526 (0.916, 2.544)	0.105		
Tumor size	0.976 (0.658, 1.448)	0.904		
α -fetoprotein	1.000 (0.999, 1.000)	0.391		
Platelet count	0.991 (0.987, 0.995)	< 0.001	0.998 (0.993, 1.004)	0.542
Alanine aminotransferase	1.008 (1.000, 1.017)	0.045	1.011 (1.003, 1.020)	0.006
Total bilirubin	0.995 (0.540, 1.833)	0.988		
Albumin	0.281 (0.168, 0.470)	< 0.001	0.481 (0.269, 0.860)	0.014
Prothrombin time (INR)	147.887 (13.992, 1563.115)	< 0.001	4.212 (0.195, 90.886)	0.359

The Cox proportional hazards model was used for univariate and multivariate analysis. The reference category for each categorical variable is in the square brackets in first column. ¹Bonferroni correction was used owing to multiple comparisons. Numbers in parentheses represent the 95%CI. HR: Hazard ratio; RF: Radiofrequency; NBNC: Non-B non-C; INR: International normalized ratio.

ARTICLE HIGHLIGHTS

Research background

Many studies have revealed that radiofrequency (RF) ablation is comparable to hepatic resection in terms of long-term survival for patients with early stage hepatocellular carcinoma (HCC). However, most studies have not taken into account the location of HCCs.

Research motivation

Our study attempted to analyze the effect of the subphrenic location of HCC on the long-term

therapeutic outcomes after hepatic resection and RF ablation.

Research objectives

To compare the long-term therapeutic outcomes between hepatic resection *vs* percutaneous RF ablation for HCCs abutting the diaphragm.

Research methods

A total of 143 Child-Pugh class A patients who had undergone hepatic resection ($n = 80$) or percutaneous RF ablation ($n = 63$) for an HCC (≤ 3 cm) abutting the right diaphragm were included. Therapeutic outcomes were compared.

Research results

Hepatic resection was an independent prognostic factor for disease-free survival (DFS) ($P \leq 0.001$; hazard ratio, 0.352; 95%CI: 0.205, 0.605; with RF ablation as the reference category); however, treatment modality was not an independent prognostic factor for overall survival (OS). The local tumor progression rate was 46.6% at 5 years for the RF ablation group.

Research conclusions

Although OS was not significantly different between patients who had undergone hepatic resection or percutaneous RF ablation for HCCs abutting the diaphragm, DFS was better in the hepatic resection group.

Research perspectives

Further studies with large sample size and multicenter prospective studies are needed to confirm the conclusion of this study.

REFERENCES

- 1 Bruix J, Sherman M; American Association for the Study of Liver Diseases. Management of hepatocellular carcinoma: an update. *Hepatology* 2011; **53**: 1020-1022 [PMID: 21374666 DOI: 10.1002/hep.24199]
- 2 Cho YK, Kim JK, Kim WT, Chung JW. Hepatic resection versus radiofrequency ablation for very early stage hepatocellular carcinoma: a Markov model analysis. *Hepatology* 2010; **51**: 1284-1290 [PMID: 20099299 DOI: 10.1002/hep.23466]
- 3 Wang JH, Wang CC, Hung CH, Chen CL, Lu SN. Survival comparison between surgical resection and radiofrequency ablation for patients in BCLC very early/early stage hepatocellular carcinoma. *J Hepatol* 2012; **56**: 412-418 [PMID: 21756858 DOI: 10.1016/j.jhep.2011.05.020]
- 4 Molinari M, Helton S. Hepatic resection versus radiofrequency ablation for hepatocellular carcinoma in cirrhotic individuals not candidates for liver transplantation: a Markov model decision analysis. *Am J Surg* 2009; **198**: 396-406 [PMID: 19520354 DOI: 10.1016/j.amjsurg.2009.01.016]
- 5 Chen J, Peng K, Hu D, Shen J, Zhou Z, Xu L, Chen J, Pan Y, Wang J, Zhang Y, Chen M. Tumor Location Influences Oncologic Outcomes of Hepatocellular Carcinoma Patients Undergoing Radiofrequency Ablation. *Cancers (Basel)* 2018; **10** [PMID: 30309001 DOI: 10.3390/cancers10100378]
- 6 Kang TW, Rhim H, Kim EY, Kim YS, Choi D, Lee WJ, Lim HK. Percutaneous radiofrequency ablation for the hepatocellular carcinoma abutting the diaphragm: assessment of safety and therapeutic efficacy. *Korean J Radiol* 2009; **10**: 34-42 [PMID: 19182501 DOI: 10.3348/kjr.2009.10.1.34]
- 7 Rhim H, Lim HK, Kim YS, Choi D. Percutaneous radiofrequency ablation with artificial ascites for hepatocellular carcinoma in the hepatic dome: initial experience. *AJR Am J Roentgenol* 2008; **190**: 91-98 [PMID: 18094298 DOI: 10.2214/AJR.07.2384]
- 8 Koda M, Ueki M, Maeda Y, Mimura K, Okamoto K, Matsunaga Y, Kawakami M, Hosho K, Murawaki Y. Percutaneous sonographically guided radiofrequency ablation with artificial pleural effusion for hepatocellular carcinoma located under the diaphragm. *AJR Am J Roentgenol* 2004; **183**: 583-588 [PMID: 15333339 DOI: 10.2214/ajr.183.3.1830583]
- 9 Kang TW, Rhim H, Lee MW, Kim YS, Choi D, Lee WJ, Lim HK. Radiofrequency ablation for hepatocellular carcinoma abutting the diaphragm: comparison of effects of thermal protection and therapeutic efficacy. *AJR Am J Roentgenol* 2011; **196**: 907-913 [PMID: 21427344 DOI: 10.2214/AJR.10.4584]
- 10 Kondo Y, Yoshida H, Tateishi R, Shiina S, Kawabe T, Omata M. Percutaneous radiofrequency ablation of liver cancer in the hepatic dome using the intrapleural fluid infusion technique. *Br J Surg* 2008; **95**: 996-1004 [PMID: 18581421 DOI: 10.1002/bjs.6058]
- 11 Kang TW, Kim JM, Rhim H, Lee MW, Kim YS, Lim HK, Choi D, Song KD, Kwon CH, Joh JW, Paik SW, Paik YH, Ahn JH. Small Hepatocellular Carcinoma: Radiofrequency Ablation versus Nonanatomic Resection--Propensity Score Analyses of Long-term Outcomes. *Radiology* 2015; **275**: 908-919 [PMID: 25688888 DOI: 10.1148/radiol.15141483]
- 12 Bruix J, Sherman M; Practice Guidelines Committee, American Association for the Study of Liver Diseases. Management of hepatocellular carcinoma. *Hepatology* 2005; **42**: 1208-1236 [PMID: 16250051 DOI: 10.1002/hep.20933]
- 13 Cucchetti A, Qiao GL, Cescon M, Li J, Xia Y, Ercolani G, Shen F, Pinna AD. Anatomic versus nonanatomic resection in cirrhotic patients with early hepatocellular carcinoma. *Surgery* 2014; **155**: 512-521 [PMID: 24439747 DOI: 10.1016/j.surg.2013.10.009]
- 14 Kim YS, Lim HK, Rhim H, Lee MW, Choi D, Lee WJ, Paik SW, Koh KC, Lee JH, Choi MS, Gwak GY, Yoo BC. Ten-year outcomes of percutaneous radiofrequency ablation as first-line therapy of early hepatocellular carcinoma: analysis of prognostic factors. *J Hepatol* 2013; **58**: 89-97 [PMID: 23023009 DOI: 10.1016/j.jhep.2012.09.020]
- 15 Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004; **240**: 205-213 [PMID: 15272045 DOI: 10.1097/SLA.0b013e3180313631]

- 15273542 DOI: [10.1097/01.sla.0000133083.54934.ae](https://doi.org/10.1097/01.sla.0000133083.54934.ae)]
- 16 **Ahmed M**, Solbiati L, Brace CL, Breen DJ, Callstrom MR, Charboneau JW, Chen MH, Choi BI, de Baère T, Dodd GD, Dupuy DE, Gervais DA, Gianfelice D, Gillams AR, Lee FT, Leen E, Lencioni R, Littrup PJ, Livraghi T, Lu DS, McGahan JP, Meloni MF, Nikolic B, Pereira PL, Liang P, Rhim H, Rose SC, Salem R, Sofocleous CT, Solomon SB, Soulen MC, Tanaka M, Vogl TJ, Wood BJ, Goldberg SN; International Working Group on Image-guided Tumor Ablation; Interventional Oncology Sans Frontières Expert Panel; Technology Assessment Committee of the Society of Interventional Radiology.; Standard of Practice Committee of the Cardiovascular and Interventional Radiological Society of Europe. Image-guided tumor ablation: standardization of terminology and reporting criteria--a 10-year update. *Radiology* 2014; **273**: 241-260 [PMID: [24927329](https://pubmed.ncbi.nlm.nih.gov/24927329/) DOI: [10.1148/radiol.14132958](https://doi.org/10.1148/radiol.14132958)]
 - 17 **Kang TW**, Lim HK, Lee MW, Kim YS, Rhim H, Lee WJ, Paik YH, Kim MJ, Ahn JH. Long-term Therapeutic Outcomes of Radiofrequency Ablation for Subcapsular versus Nonsubcapsular Hepatocellular Carcinoma: A Propensity Score Matched Study. *Radiology* 2016; **280**: 300-312 [PMID: [26824711](https://pubmed.ncbi.nlm.nih.gov/26824711/) DOI: [10.1148/radiol.2016151243](https://doi.org/10.1148/radiol.2016151243)]
 - 18 **Shiina S**, Tateishi R, Arano T, Uchino K, Enooku K, Nakagawa H, Asaoka Y, Sato T, Masuzaki R, Kondo Y, Goto T, Yoshida H, Omata M, Koike K. Radiofrequency ablation for hepatocellular carcinoma: 10-year outcome and prognostic factors. *Am J Gastroenterol* 2012; **107**: 569-577; quiz 578 [PMID: [22158026](https://pubmed.ncbi.nlm.nih.gov/22158026/) DOI: [10.1038/ajg.2011.425](https://doi.org/10.1038/ajg.2011.425)]
 - 19 **Lee DH**, Lee JM, Lee JY, Kim SH, Yoon JH, Kim YJ, Han JK, Choi BI. Radiofrequency ablation of hepatocellular carcinoma as first-line treatment: long-term results and prognostic factors in 162 patients with cirrhosis. *Radiology* 2014; **270**: 900-909 [PMID: [24475823](https://pubmed.ncbi.nlm.nih.gov/24475823/) DOI: [10.1148/radiol.13130940](https://doi.org/10.1148/radiol.13130940)]
 - 20 **Huang J**, Yan L, Cheng Z, Wu H, Du L, Wang J, Xu Y, Zeng Y. A randomized trial comparing radiofrequency ablation and surgical resection for HCC conforming to the Milan criteria. *Ann Surg* 2010; **252**: 903-912 [PMID: [21107100](https://pubmed.ncbi.nlm.nih.gov/21107100/) DOI: [10.1097/SLA.0b013e3181efc656](https://doi.org/10.1097/SLA.0b013e3181efc656)]
 - 21 **Vivarelli M**, Guglielmi A, Ruzzenente A, Cucchetti A, Bellusci R, Cordiano C, Cavallari A. Surgical resection versus percutaneous radiofrequency ablation in the treatment of hepatocellular carcinoma on cirrhotic liver. *Ann Surg* 2004; **240**: 102-107 [PMID: [15213625](https://pubmed.ncbi.nlm.nih.gov/15213625/) DOI: [10.1097/01.sla.0000129672.51886.44](https://doi.org/10.1097/01.sla.0000129672.51886.44)]
 - 22 **Nishikawa H**, Inuzuka T, Takeda H, Nakajima J, Matsuda F, Sakamoto A, Henmi S, Hatamaru K, Ishikawa T, Saito S, Nasu A, Kita R, Kimura T, Arimoto A, Osaki Y. Comparison of percutaneous radiofrequency thermal ablation and surgical resection for small hepatocellular carcinoma. *BMC Gastroenterol* 2011; **11**: 143 [PMID: [22204311](https://pubmed.ncbi.nlm.nih.gov/22204311/) DOI: [10.1186/1471-230X-11-143](https://doi.org/10.1186/1471-230X-11-143)]
 - 23 **Hong SN**, Lee SY, Choi MS, Lee JH, Koh KC, Paik SW, Yoo BC, Rhee JC, Choi D, Lim HK, Lee KW, Joh JW. Comparing the outcomes of radiofrequency ablation and surgery in patients with a single small hepatocellular carcinoma and well-preserved hepatic function. *J Clin Gastroenterol* 2005; **39**: 247-252 [PMID: [15718869](https://pubmed.ncbi.nlm.nih.gov/15718869/) DOI: [10.1097/01.mcg.0000152746.72149.31](https://doi.org/10.1097/01.mcg.0000152746.72149.31)]
 - 24 **Hasegawa K**, Makuuchi M, Takayama T, Kokudo N, Arai S, Okazaki M, Okita K, Omata M, Kudo M, Kojiro M, Nakanuma Y, Takayasu K, Monden M, Matsuyama Y, Ikai I. Surgical resection vs. percutaneous ablation for hepatocellular carcinoma: a preliminary report of the Japanese nationwide survey. *J Hepatol* 2008; **49**: 589-594 [PMID: [18620773](https://pubmed.ncbi.nlm.nih.gov/18620773/) DOI: [10.1016/j.jhep.2008.05.018](https://doi.org/10.1016/j.jhep.2008.05.018)]
 - 25 **Shibata T**, Isoda H, Hirokawa Y, Arizono S, Shimada K, Togashi K. Small hepatocellular carcinoma: is radiofrequency ablation combined with transcatheter arterial chemoembolization more effective than radiofrequency ablation alone for treatment? *Radiology* 2009; **252**: 905-913 [PMID: [19567647](https://pubmed.ncbi.nlm.nih.gov/19567647/) DOI: [10.1148/radiol.2523081676](https://doi.org/10.1148/radiol.2523081676)]
 - 26 **Peng ZW**, Zhang YJ, Chen MS, Xu L, Liang HH, Lin XJ, Guo RP, Zhang YQ, Lau WY. Radiofrequency ablation with or without transcatheter arterial chemoembolization in the treatment of hepatocellular carcinoma: a prospective randomized trial. *J Clin Oncol* 2013; **31**: 426-432 [PMID: [23269991](https://pubmed.ncbi.nlm.nih.gov/23269991/) DOI: [10.1200/JCO.2012.42.9936](https://doi.org/10.1200/JCO.2012.42.9936)]
 - 27 **Hirooka M**, Kisaka Y, Uehara T, Ishida K, Kumagi T, Watanabe Y, Abe M, Matsuura B, Hiasa Y, Onji M. Efficacy of laparoscopic radiofrequency ablation for hepatocellular carcinoma compared to percutaneous radiofrequency ablation with artificial ascites. *Dig Endosc* 2009; **21**: 82-86 [PMID: [19691779](https://pubmed.ncbi.nlm.nih.gov/19691779/) DOI: [10.1111/j.1443-1661.2009.00836.x](https://doi.org/10.1111/j.1443-1661.2009.00836.x)]

P- Reviewer: Aykan NF, He J, Huang C, Lambrecht NW

S- Editor: Ji FF **L- Editor:** A **E- Editor:** Song H





Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

