

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 43026

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Risk of cholangiocarcinoma in patients undergoing therapeutic endoscopic retrograde cholangiopancreatography or cholecystectomy: A population based study

Chi-Chih Wang, Ming-Chang Tsai, Wen-Wei Sung, Tzu-Wei Yang, Hsuan-Yi Chen, Yao-Tung Wang, Chang-Cheng Su, Ming-Hseng Tseng, Chun-Cha Lin

Match Overview

There are no matching sources for this report

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 26,100 条结果 (用时 0.35 秒)


Patients ever having had a **cholecystectomy** had a significantly lower **risk** of the **studied** malignancies. Conclusions: The **risk** of malignancy in the bile ducts, liver, or pancreas is elevated after **ERCP** in benign disease. However, ES does not seem to affect this **risk**.

Endoscopic Sphincterotomy and Risk of Malignancy in the Bile Ducts ...

[www.cghjournal.org/article/S1542-3565\(08\)00353-4/pdf](http://www.cghjournal.org/article/S1542-3565(08)00353-4/pdf)

 关于这条结果的详细信息  反馈

Endoscopic sphincterotomy and risk of cholangiocarcinoma: a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5063738/>  翻译此页

作者 : C Strömberg - 2016 - 被引用次数 : 2 - 相关文章

2016年9月14日 - In this **study**, we also found a decreased **risk** of malignancy in **patients** who had had a **cholecystectomy**. ... A secondary aim was to **study** the relationship between severe common bile duct stone exposure, which is a valid assumption for the entire **ERCP** cohort irrespective of ES, and malignancy in the biliary tract.

The role of endoscopic retrograde cholangiopancreatography in ...

<https://www.ncbi.nlm.nih.gov/pubmed/7797023> - 翻译此页

作者 : RA Erickson - 1995 - 被引用次数 : 130 - 相关文章

The role of **endoscopic retrograde cholangiopancreatography** in **patients** with ... The aim of this **study** is to use decision analysis to assess four different approaches to using **ERCP** in **patients** undergoing laparoscopic **cholecystectomy**. ... (2) selective preoperative **ERCP** for **patients** at high **risk** for choledocholithiasis, ...

缺少字词 : cholangiocarcinoma population



Risk of Cholangiocarcinoma in Patients Undergoing Therapeutic Endosc



全部

图片

新闻

购物

地图

更多

设置

工具

找到约 20,900 条结果 (用时 0.51 秒)

Google 学术 : Risk of Cholangiocarcinoma in Patients Undergoing Therapeutic Endoscopic Retrograde Cholangiopancreatography or Cholecystectomy: A Population Based Study

Diagnosis and management of primary sclerosing ... - Chapman - 被引用次数 : 882

Cancer surveillance in patients with primary sclerosing ... - Razumilava - 被引用次数 : 205

... and therapeutic retrograde cholangiography in patients ... - Wang - 被引用次数 : 123

Endoscopic sphincterotomy and risk of cholangiocarcinoma: a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5063738/> ▼ 翻译此页

作者 : C Strömberg - 2016 - 被引用次数 : 2 - 相关文章

2016年9月14日 - Results: A total of 69 925 patients undergoing ERCP from 1976 ... in therapeutic regime is that an increased number of young patients, with a long ... risk of malignancy in patients who had had a cholecystectomy. The primary aim of this population-based cohort study of all patients registered for an ERCP in ...

Endoscopic retrograde cholangiopancreatography and laparoscopic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3785632/> ▼ 翻译此页

作者 : JF Zang - 2013 - 被引用次数 : 19 - 相关文章

2013年9月28日 - National Institutes of Health ... Keywords: Cholecystectomy, Laparoscopic, Endoscopic, Therapy ... In this retrospective study, 156 patients with gallstones and CBD ... by the same surgical team at Taizhou People's Hospital (Taizhou, ... During the same period, two patients undergoing ERCP in group 1 and ...

Endoscopic Sphincterotomy and Risk of Malignancy in the Bile Ducts ...

[https://www.cghjournal.org/article/S1542-3565\(08\)00353-4/pdf](https://www.cghjournal.org/article/S1542-3565(08)00353-4/pdf) - 翻译此页

作者 : C Strömberg - 2008 - 被引用次数 : 20 - 相关文章

[全部](#)[图片](#)[新闻](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 21,000 条结果 (用时 0.52 秒)

Endoscopic sphincterotomy and risk of cholangiocarcinoma: a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5063738/> - 翻译此页

作者: C Strömberg - 2016 - 被引用次数: 2 - 相关文章

2016年9月14日 - Results: A total of 69 925 patients undergoing ERCP from 1976 ... in therapeutic regime is that an increased number of young patients, with a long ... risk of malignancy in patients who had had a cholecystectomy. The primary aim of this population-based cohort study of all patients registered for an ERCP in ...

Adverse events in older patients undergoing ERCP: a systematic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4423280/> - 翻译此页

作者: LW Day - 2014 - 被引用次数: 23 - 相关文章

2014年3月7日 - However, ERCP is not without risk, with a reported adverse event rate ... With the dramatic rise in the elderly population, it is anticipated that ERCP will be in greater ... Studies were assigned weights based on their sample size (number ... Efficacy and safety of therapeutic ERCP in patients 90 years of age ...

Endoscopic Sphincterotomy and Risk of Malignancy in the Bile Ducts ...

[www.cghjournal.org/article/S1542-3565\(08\)00353-4/pdf](http://www.cghjournal.org/article/S1542-3565(08)00353-4/pdf) - 翻译此页

作者: C Strömberg - 2008 - 被引用次数: 41 - 相关文章

Patients ever having had a cholecystectomy had a significantly lower risk of the studied malignancies. Conclusions: The risk of malignancy in the bile ducts, liver, or pancreas is elevated after ERCP in benign disease. However, ES does not seem to affect this risk.

(PDF) Endoscopic Sphincterotomy and Long-Term Risk of ...

https://www.researchgate.net/.../5373356_Endoscopic_Sphincterotomy_and... - 翻译此页

2018年8月1日 - We used data from population-based Danish health-care registries to ... The