

ALL IMAGES VIDEOS

162 Results Any time ▾

Hemophilic pseudotumor in two parts of the maxilla: Case ...

<https://www.researchgate.net/publication/304664493...>

This report describes a case of hemophilic pseudotumor of the maxillary alveolar ridge that occurred in a boy with mild hemophilia A (14% factor VIII activity).

Haemophilic pseudotumour in two parts of the maxilla: case ...

<https://www.birpublications.org/doi/full/10.1259/dmfr.20150440>

The nasopalatine duct is dislocated to the right side (arrow). The lesion mimics fibrous dysplasia or intraosseous vascular malformation. (d) On the sagittal contrast-enhanced CT scan with bone window setting, the cystic lesion A is adjacent to the intraosseous lesion B, and P is the pterygoid plate. ... report of case with review of literature ...

Cited by: 1 Author: Ah-Young Kwon, Kyung-Hoe Huh, Won-Ji...
Publish Year: 2016

Haemophilic pseudotumour in two parts of the maxilla: case ...

europepmc.org/abstract/MED/27359071 ▾

Jul 25, 2016 · Thus far, there have been only 31 reported cases in jaw bones and paranasal sinuses. Among them, the mandible is a more common site than the maxilla or paranasal sinuses. Here, we report a case of haemophilic pseudotumour in two parts of the maxilla.

Cited by: 1 Author: Ah-Young Kwon, Kyung-Hoe Huh, Won-Ji...
Publish Year: 2016

Pseudotumor of hemophilia in the mandible of a patient ...

<https://www.researchgate.net/publication/225282801...>

We report a case of hemophilic pseudotumor arising in the maxilla of a 46-year-old man with a history of hemophilia A. The initial symptom was painless swelling in the right buccal region.

(PDF) Pseudotumor of the Mandible as First Presentation of ...

<https://www.researchgate.net/publication/51121897...>

Pseudotumor of the Mandible as First Presentation of Hemophilia in a 2-Year-Old Male: A Case Report and Review of Jaw Pseudotumors of Hemophilia ... or hemophilia B. A review of the literature ...



Intraosseous venous malformation of the maxilla after enucleation <



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

11,300 Results

Any time ▾

[Haemophilic pseudotumour in two parts of the maxilla: case ...](#)

<https://www.birpublications.org/doi/10.1259/dmfr.20150440>

Here, we report a case of **haemophilic pseudotumour** in two parts of the **maxilla**. Contrast-enhanced CT showed an expansive and thinly corticated lesion with fluid attenuation at the **left anterior maxilla** which seemed like a **post-operative maxillary cyst**, ameloblastoma or odontogenic cyst.

Cited by: 2

Author: Ah-Young Kwon, Kyung-Hoe Huh, Won-Ji...

Publish Year: 2016

[Hemophilic pseudotumor in two parts of the maxilla: Case ...](#)

<https://www.researchgate.net/publication/304664493...>

Here, we report a case of **hemophilic pseudotumor** in two parts of the **maxilla**. Contrast-enhanced **computed tomography** showed an expansive and thinly corticated lesion with fluid attenuation at the...

[Haemophilic pseudotumour of the mandible: report of three ...](#)

<https://www.researchgate.net/publication/6363194...>

Here, we report a case of **hemophilic pseudotumor** in two parts of the **maxilla**. Contrast-enhanced **computed tomography** showed an expansive and thinly corticated lesion with fluid attenuation at the...

[Haemophilic pseudotumour of the mandible in a 5-year-old ...](#)

<https://www.researchgate.net/publication/45650273...>

Here, we report a case of **hemophilic pseudotumor** in two parts of the **maxilla**. Contrast-enhanced **computed tomography** showed an expansive and thinly corticated lesion with fluid attenuation at the...

[Pseudotumor of hemophilia in the mandible of a patient ...](#)

<https://www.sciencedirect.com/science/article/pii/S1079210411005701>

Feb 01, 2012 : The majority of **hemophilic pseudotumors** occur within soft tissues (**intramuscular**) and long bones of adult males. Fewer than 20 cases have been reported in the **maxillofacial region**. We report a rare case occurring in the **mandible** of a 14-year-old boy who presented with considerable expansion and displacement of teeth.

Cited by: 12

Author: Roger R. Thronsdon, David Baker, Patric...

Publish Year: 2012

<https://www.birpublications.org/doi/10.1259/dmfr.20150440>

Text-Only Report

国内版 国际版



Intraosseous venous malformation of the maxilla after enucleation



ALL

IMAGES

VIDEOS

4,320 Results

Any time ▾

Haemophilic pseudotumour in two parts of the maxilla: case ...

<https://www.birpublications.org/doi/10.1259/dmfr.20150440>

The aim of this **report** was to present a **case** of a haemophilic pseudotumour that occurred in two parts **of the maxilla**. An expansive and thinly corticated lesion with fluid attenuation at the left anterior **maxilla** mimicked a post-operative maxillary cyst, ameloblastoma or ...

Cited by: 3

Author: Ah-Young Kwon, Kyung-Hoe Huh, Won-...

Publish Year: 2016

Hemophilic pseudotumor in two parts of the maxilla: Case ...

https://www.researchgate.net/publication/304664493_Hemophilic_pseudotumor_in_two_parts...

This report describes a case of **hemophilic pseudotumor of the maxillary alveolar ridge** that occurred in a **boy with mild hemophilia A (14% factor VIII activity)**.

Haemophilic pseudotumour in two parts of the maxilla: case ...

europepmc.org/abstract/MED/27359071 ▾

Jul 25, 2016 · An expansive and thinly corticated lesion with fluid attenuation at the **left anterior maxilla** mimicked a **post-operative maxillary cyst**, ameloblastoma or odontogenic cyst radiologically. In addition, the thickened **left palatal process of the maxilla** seemed like fibrous dysplasia or **intraosseous vascular malformation**.

Cited by: 3

Author: Ah-Young Kwon, Kyung-Hoe Huh, Won-...

Publish Year: 2016

Haemophilic pseudotumour of the mandible: report of three ...

https://www.researchgate.net/publication/6363194_Haemophilic_pseudotumour_of_the...

We report a case of **hemophilic pseudotumor arising in the maxilla** of a 46-year-old man with a **history of hemophilia A**. The initial symptom was painless swelling in the right buccal region.